INDEX OF SURGICAL PROGRESS.

GENERAL SURGERY.

I. Treatment of Erysipelas. By Dr. W. OTTO (Rudolph Hospital, Vienna). In 1883 Barwell reported five cases of erysipelas very successfully treated by exclusion of air. For this purpose he used white lead paint. Dr. J. Breuer related to the author a favorable case similarly treated with liquor gutta-perchæ as an air-excluder. Otto uses a solution of 2 parts wax, 20 parts dryer and 100 linseed oil. With this he makes repeated applications over the affected part and a hand's breadth beyond. Five hospital cases are described and four dispensary patients mentioned. In all the temperature very soon fell, the pain stopped and no further extension occurred. The fact that erysipelas spreads through the upper layers of cutis may possibly depend on the accessibility of oxygen. [Local applications of lead paint, collodion, vaseline smeared on thick, etc., are esteemed by many American practitioners. *Reptr.*—*Wien. Med. Woch.* 1886. No. 43.]

II. A Contribution to the Pathogenesis of Actinomycosis of the Lung. By Dr. J. ISRAEL (Berlin). Israel's present case was that of a Russian driver, æt. 26, who had usually slept on straw or in hay lofts, and sometimes drank from the same trough as his horse.

After some pain in the left side of the chest in October, 1886, abscesses formed below the left nipple: these increased in number and developed ulcers. On admission in August, 1885, there was marked emaciation and considerable contraction of the left side of the thorax—the latter covered with abscesses and sores whose secretion contained abundant actinomycosis-granules. The muco-purulent sputum—at times tinged with blood—always contained the same. Death the end

of March, 1886, from diarrhœa, resulting from amyloid degeneration of the abdominal organs.

A single focal cavity in the lower part of the left upper pulmonary lobe, close to the anterior surface. From this the process had freely perforated the front wall of the chest.

In the actinomycotic pulmonary cavity was a lentil-sized foreign body which proved macroscopically, microscopically and chemically to be a fragment of a tooth.

This is the first positive of proof of the view which I. had previously advocated, viz., that pulmonary actinomycosis does not result from the inspiration of germs in the air but from the aspiration of germs from the oral or adjacent cavities, carious teeth evidently being one of the primary seats.

In the discussion, Roser (Marburg) noted an acute and a chronic form of actinomycosis. König (Göttingen) spoke of seeing dispensary cases frequently. Esmarch had seen ten cases with one death. The patients are almost always cattle men.—*Centbl. f. Chirg.* 1886. No. 24. Report of XV. German Surgical Congress. *Arch. f. klin. Chirg.* Bd. 34, Hft. I.

W. BROWNING (Brooklyn).

III. A Case of Actinomycosis of the Lung. By A. J. OCHSNER, M.D., (Chicago). Male, æt. 56, stock-raiser by occupation. After a history of severe pain in the antrum for some months, a spontaneous discharge of pus into the pharynx, with establishment of a permanent fistula, gave relief—this in 1878. Some of the discharge finds entrance into larynx during sleep, exciting severe cough. In 1880 antrum was trephined from the mouth, scraped and irrigated, the irrigation being continued daily for two years. In 1882, simultaneously with change of residence, Northern Mexico, and the territory between it and Colorado, the antrum closed and the general health was much improved. In 1885 pulmonary symptoms, consisting of suffocative sensations and cough began to trouble him. Expectorates mucus and blood, and in this sputum the characteristic fungus of actinomycosis is readily found by microscopic examination. His position is

stooping, the chest is full in front, and there is a decrease of motion on the left side, with dulness, roughened respiratory sounds and numerous mucous râles. Below the upper border of the fifth rib, and throughout the right side the sounds are normal.

The man has been engaged in raising, buying and selling, and handling large numbers of cattle for more than forty years. Among these animals there were many suffering from the disease known as lumpy-jaw, and it was the practice of the patient to cure the animals thus affected, by freely opening the abscess by crucial incision, extirpating as much as possible of the lump and introducing about one drachm of powdered arsenic into the cavity. Repeating this once or twice, usually effected a permanent cure.—*Chicago Med. Jour. and Exam.* Dec. 1886.

L. S. PILCHER (Brooklyn).

IV. A Case of Actinomycosis. Dr. WM. O'NEILL (Lincoln). A farmer, æt. 50, had been "dressing" calves affected with what he called "ring-worm." In a few weeks he was attacked with a pustular eruption on the left wrist and forearm, and a slight, apparently glandular swelling under the right ear. When (one month afterward) he first consulted Dr. O'Neill, there were also four or five smaller swellings in the neighborhood of the clavicles and one in the left axilla. Iodine and arsenic given internally, citrine ointment locally. Skin eruption almost disappeared in a few weeks, tumours slowing diminishing. But, after two months of treatment, tonsillitis and aphthous looking state of mouth.

Next month, mouth better, but two or three new enlargements appear; some of old ones larger. The one below the ear was as large as a small hen's egg, smooth, nearly painless, elastic, and slightly movable. Other smaller swellings, one beneath chin, one on back of head.

Now, sublimate and pot. iod. internally, iodide of ammon. ointment externally.

Next month, appearances much better, but unfortunately severe bronchitis set in. On its subsidence, the tumour, etc., are left much worse.

"The tumours in the angle of the neck had extended in all directions, more especially downwards. It had also suppurated, and two or three sinuses gave exit to a whitish purulent fluid, which frequently contained small yellowish particles. The sore generally presented a reddish granulating surface, but occasionally dark colored scabs would form on it. Several of the smaller tumours on the lower part of the neck had nearly disappeared, but the one under the chin was still large enough to be uncomfortable, and the tumour on the back of the head had become as big as a walnut, and, although unulcerated, was so painful that he could hardly sleep on his back. A small swelling had also formed under the left ear." The increase of temperature and pulse-frequency less than might have been expected.

At this stage iodide and bromide of ammonium were commenced. But, the tumours continuing to enlarge, the opinion of Mr. A. E. Barker was sought with a view to surgical, operative treatment.

Dr. O'Neill found the micro organisms "supposed to be peculiar to the disease."

Speaking of the calves with so-called "ring-worm," Dr. O'Neill says, "there are no lumps at present under the jaws, or in or about their mouths. Ring-worm began in them, as well as in several others, by the hair falling off, a reddening or inflammation of the skin, and the formation of ash-colored scabs, more or less thick, according to the severity of the disease in the part affected. The disease is caused by a fungus which can be destroyed by sulphur or by mercurial preparation, notably by corrosive sublimate." But Dr. O'Neill had not, up to the date of his paper, identified the fungus of the calf's disease with that of actinomycosis.

Up to recent years, actinomycosis has probably been confounded with scrofula and cancer, with the former in its early stages, and with the latter in its ulcerated stages.

About one-half of the published cases have been fatal. The disease may attack almost any organ in the body.

O'Neill thinks that iodine, arsenic and mercury have some power over the disease in its early stages.

But, after all, where actinomycosis appears in the shape of externa,

tumours, he thinks that the most expeditious and the best treatment would be *removal by surgical operation*.

Actinomycosis appears to have been first discovered by Italian veterinary surgeons and accurately described by Revolta in 1868. Bollinger rediscovered it, and named it in 1877, and, in the following year, Israel described the first case in man.

C. B. KEETLEY (London).

OPERATIVE SURGERY.

I. New Method of Amputation at the Ankle-Joint. By Prof. A. C. TAUBER (St. Petersburg, Russia). After having criticized

FIG. 1. STUMP PRODUCED BY TAUBER'S METHOD.

the methods of operation at the ankle-joint, introduced by Syme, Malgaigne, Roux, Sedillot, Le Fort and other surgeons, Prof. T. has demonstrated in the first congress of the Russian physicians his own method which is a modification of Pirogoff's. Fig. 1 shows a stump made according to Tauber's method. His operation Prof. T. performs as follows.