found empty. Hemorrhage from the lateral sinus compelled tamponing and this entailed a destructive process with hemiplegia and the patient died six weeks later; no necropsy.

5. BRAIN. MENINGES.

Herrick, W. W., and Parkhurst, G. M. MENINGOCOCCUS ARTHRITIS.
[American Journal Medical Sciences, 1919, CLVIII, No. 4, 473.]

After a brief review of the literature, report is made of twenty-eight cases of meningococcus arthritis occurring in an epidemic of 321 cases of meningococcic infection. In addition to these twenty-eight cases of meningococcic arthritis, there were twelve examples of serum arthritis. Three types of arthritis are described. The first, Type A, an acute polyarthritis involving wrists, knees, elbows, ankles, at times almost all the important joints. Its pathology is apparently hemorrhage into the synovia or periarticular tissues. The pain is severe and other inflammatory symptoms, except swelling, are marked. The duration is short and other metastatic complications of meningococcic infection, such as panophthalmitis, epididymitis, pneumonia are frequent. This early polyarthritis characterizes the severe infections and usually accompanies those cases with marked hemorrhagic rash, and makes the outlook somewhat grave. Of the twelve examples reported, four died—a mortality of $33\frac{1}{3}$ per cent.

The second type of arthritis, Type B, is quite different. It is monoarticular, usually involving the knee. The pathology is a purulent arthritis, the exudate containing demonstrable meningococci in many cases. The swelling is marked, but the other inflammatory symptoms are slight. The marked disproportion between the swelling and the other inflammatory signs is the striking feature of this type of arthritis. It occurs late in the course of the disease, is less often attended by other complications, is moderately prolonged and the prognosis favorable. The mortality in sixteen cases was 12.5 per cent. Aspiration of the exudate and local injection of serum are advisable when the exudate is large. The third type of arthritis, designated Type C, is the ordinary serum arthritis which does not require description. [Author's abstract.]

Stangl, F. H. MENINGITIS AND INFLUENZA. [Journal A. M. A., Oct. 4, 1919.]

The author notices in the reports of the pandemic of influenza references made to toxemia and to the symptoms of shock and meningitis, and quotes a number of authorities who have specially mentioned such conditions. He has, therefore, reviewed again the records utilized by Keeton and Cushman, with the addition of those which have accumulated since their report, 3,400 cases in all, and finds that nearly 1 per cent. of the total showed symptoms suggesting meningitis or cerebral involvement, ranging from slight neck rigidity and bilateral or lateral Kernig reactions to deep delirium and marked stiffness of neck, and in one

case to opisthotonus. Eight fatal cases, variously diagnosed as epidemic meningitis, uremia, tuberculous meningitis, apoplexy and polyarthritis, proved, on necropsy, to be influenza with marked lung involvement. The polyarthritis case is reported, the meningeal symptoms being most prominent. Another group of cases diagnosed as influenza-pneumonia or meningism, comprising twenty-one cases, including seven deaths, is tabulated. In all, rigidity of the neck was noted, the Kernig sign was positive in thirteen, and in six the Brudzinski sign was positive, and opisthotonos was present in one. The necropsy findings showed the usual influenzal lesions. The brain and its membranes presented only congestion and edema. The author summarizes as follows: "A severe toxemia and active delirium with definite meningeal manifestations such as is encountered in systematic infection and the acute exanthems occur in some patients suffering from influenza and influenzal pneumonia. The clinical picture in some instances closely simulates that of an actual meningitis or other intracranial causes of delirium and unconsciousness, and postmortem examination fails to reveal any inflammation of the brain or its membranes."

Caldwell, C. N. EPIDEMIC CEREBROSPINAL MENINGITIS. [Kentucky Med. Jl., Nov., 1919.]

Caldwell here reports upon 113 patients of whom he found seventy-two were of the type described by certain authors with which he was familiar as usual, nine chronic, ten abortive and twenty-two fulminating. Sixty-three had rigidity of the neck and twenty-nine had herpes; twenty-one had retention; an eruption was seen in sixty-one, and three were jaundiced; all had headache. These complications were observed: pneumonia, seventeen; eye complications, twelve; mastoiditis, one; nephritis, eight; metastatic abscesses, two; synovitis, two; pericarditis with effusion, one; empyema, one; deafness, two, and brain abscess, two. Total mortality 34 per cent., but ten of these patients had an intercurrent pneumonia of longer standing than the meningitis, one had an empyema and one had a brain abscess that was an extension from the ear.

Wegeforth, P., and Latham, J. R. LUMBAR PUNCTURE IN MENINGITIS. [A. M. Jl. Med. Sc., Aug., 1919.]

Infections of the meninges has been observed to occur following a lumbar puncture. Five such instances are cited by these authors in this paper. To prevent the possible accidental production of a meningitis following lumbar puncture it is recommended that careful consideration to the bacteriologic study of the blood be given before such punctures in acute diseases,, in the absence of definite signs of irritation of the central nervous system, lumbar puncture should be avoided unless it is first conclusively shown that the blood stream is free of infection;