what I considered a mild infection of hookworms, I gave him the usual treatment, which brought forth a large number of worms. He immediately began to improve and on the fourth day said he felt like a different person. The chancre quickly disappeared, as did all the other symptoms. One week later I gave him another treatment for the worms and he expelled a very few more. From this time on he rapidly improved and at the end of ten days he was as well as he ever was.

Did the hookworm infection retard the effects of the salvarsan? My suggestion is that the toxins produced by the worms either retarded or counteracted the effect of the salvarsan. I really think that the toxins only retarded the action of the salvarsan and that in a short time the salvarsan would have produced its effects.

SPIROCHETES IN THE MOUTH*

A. A. THIBAudeau, M.B., BUFFALO, N. Y.

In the routine examination of smears from the mouth, as carried out by students in the class-room, a considerable number were found, which, when stained by the Giemsa method, showed spirochetes morphologically resembling *Spirocheta pallida*. As some observers have considered it possible to make a diagnosis of syphilis by the finding of *Spirocheta pallida* in smears from suspicious sores in the mouth and pharynx, it was thought desirable to examine the smears from a number of normal mouths.

Smears were accordingly made from apparently healthy mouths, using the deposit at the junction of the gums with the teeth, and from the root of the tongue. After fixation in absolute alcohol they were stained by Giemsa's rapid method. Smears from each case were also stained by Loeffler's methylene-blue or an aqueous solution of gentian-violet.

Three main types of spirochetes were found:

1. The large *Spirocheta buccalis* with long flat curves, which is much thicker and wider than the other spiral forms found in the mouth, and which has fewer turns.

2. A moderate-sized spirochete, probably the "medium form" of *Spirocheta dentium* described by Hoffmann and Prowazek.¹ This organism stains a very faint purple by Giemsa. The coils of this spirochete, as found in mouth smears, are not so sharp or so regular as those of *Spirocheta pallida*, and the organism is thicker.

3. A small, very delicate spirochete, with sharp coils fairly closely applied was the form that most resembled *Spirocheta pallida*—probably the *Spirocheta microdentium* of Noguchi.²

This organism very frequently exhibited irregularities in the arrangement of the coils, seldom seen in *Spirocheta pallida*, but specimens were found in which the coils were regular throughout the organism. The spirochete, stained by Giemsa's rapid method, showed a coloration from a rose-red to a light purple, depending apparently on the degree of heat used and the length of time that it was exposed to the stain. Typical organisms of this type showed pointed ends. It stained but poorly with Loeffler's methylene-blue, and in about 30 per cent. of the cases, where present, could not be found in the control smears stained by this method. It is to be noted that while these organisms may not present all the characteristics of a typical *Spirocheta pallida*, there are usually in stained smears from definite syphilitic lesions, and in which typical *Spirocheta pallida* may be present, numbers of spirochetes, undoubtedly *Spirocheta pallida*, which are not typical in every respect, and which could not be distinguished in stained smears from this type of mouth spirochete.

I have examined smears from 149 mouths; in sixty-one, or 40.9 per cent. of these, spirochetes of the last type, which might be taken for *Spirocheta pallida*, have been found. In none of these cases did inquiry bring out a history of syphilitic

infection. The possibility of these organisms causing a mistake in diagnosis in a non-syphilitic ulceration or other non-syphilitic lesion of the mouth seems self-evident.

It has long been known that the differentiation of *Spirocheta pallida* from some of the mouth spirals is extremely difficult and even at times impossible. My investigation has shown that spirochetes closely resembling *Spirocheta pallida* were found in as many as 41 per cent. of normal mouths examined.

24 High Street.

A SIMPLE REFRIGERATOR FOR LABORATORY WORKERS

LOYD OSCAR THOMPSON, PH.B., M.D., LITTLE ROCK, ARK.

City Bacteriologist; Pathologist to St. Vincent's Hospital

Many physicians who do their own laboratory work, and even many of the smaller clinical laboratories, do not have refrigerator facilities, and it is often a problem to keep certain substances at a low temperature. This problem was solved recently in the Little Rock City Laboratory, when the ice-man failed to fill the refrigerator, by utilizing a vacuum bottle. We wanted to keep some guinea-pig blood at a low temperature over night to collect the serum for the Wassermann reaction. The vacuum bottle was filled about one-half full with ice-water and the test-tube containing the blood was lowered into it and the cork inserted. In the morning the temperature was 10 C., and the serum had completely separated.

This simple expedient may be used as above, for keeping complement, corpuscle suspension, antigen and patients' serum for the Wassermann reaction; for keeping urine, small samples of milk, etc.

606 State Bank Building.

Therapeutics

PULMONARY TUBERCULOSIS

In the treatment of this disease the following important points should not be forgotten. It is not sufficient to order fresh air, rest, proper diet and such medication as needed; more must be done. So much has been accomplished by fresh air, rest and food that the physician is too likely not to treat the patient as though he were seriously ill. While daily attention is not needed, unless the patient has considerable fever, a weekly study of his case and his symptoms and his signs are essential to the best treatment for him and for the promotion of as rapid recovery as possible, if he is to recover. To repeat, the diagnosis is not the only essential to the successful treatment of pulmonary tuberculosis. The patient must be studied weekly; physical examination must be made weekly to know exactly what is being accomplished by the treatment, or to ascertain why the treatment is not producing results. Toward this end the following items are suggestive, "lest one forget."

1. The cause of the patient's temperature should be sought. It may not be from the tubercle bacillus or the active tuberculous process alone. It may be from a coincident streptococcal infection, or the so-called double infection with a purulent bronchial exudate. There may be a pathologic condition of the throat and nose; there may be a digestive disturbance; there may be a genito-urinary disturbance; in fact, all other causes of high temperature must be eliminated before the decision is made that it is due to a more or less acute tuberculous process.

2. Continuous loss of weight may mean that the food is not sufficient, that the food is not properly digested or that there is some other source of loss to the individual than the tuberculous process. If the patient is a woman,

* From the Laboratory of Bacteriology, University of Buffalo.

¹ Read at the Annual Meeting of the American Association of Pathologists and Bacteriologists, April 4-5, 1912.

² Hoffmann and Prowazek: *Centralbl. f. Bakteriol.*, 1906, xii, 817.

³ Noguchi: *Jour. Exper. Med.*, 1912, xv, 81.