muscles were not only widely separated, but greatly atrophied. The entire abdominal wall was not over 1 cm. in thickness. As soon as the abdomen was opened, a little turbid, rather dark colored fluid poured out, and when the stomach was drawn out, a few flakes of lymph were found on it. In a moment the perforation was discovered. It was an aperture slightly oval, very sharply punched out, 5 mm. in its long diameter, situated precisely in the middle of the pylorus anteriorly. The whole pylorus was thickened, but not nodulated. Evidently it had been the seat of an old ulcer involving all the anterior wall. The ulcer was immediately closed by a continuous Lembert suture. The abdominal cavity was then flushed out with a large quantity of salt solution, care being taken to wash out between the diaphragm and the liver, and also down into Douglas' cul-de-sac, where a considerable amount of turbid fluid was found. The diaphragm was but little injected. The transverse colon was then turned up, a hole torn through the mesocolon, and a posterior gastro-enterostomy done by means of a Murphy button. The peritoneal cavity was then again most carefully flushed out till the fluid, as in the first instance, returned clear. A number of adhesions were found between the coils of the intestine and the anterior abdominal wall and the omentum and the abdominal wall in the lower half of the abdomen, but I did not think it wise to prolong the operation by interfering with them. She was placed in bed in Fowler's position (i. e., with head of the bed raised).

She made an uninterrupted recovery, her highest temperature being 100.4 F. On December 1, five days after the operation, her temperature reached normal. In consequence of the thinness of her abdominal wall, I kept her in bed for four weeks. She went home December 31, wearing a belt and able to walk about without trouble.

The recovery from so serious a lesion as a perforating gastric ulcer and so serious an operation as abdominal section and gastro-enterostomy in a woman past 70, is a sufficient reason, it would seem, for recording such a case. Her recovery was due, I have no question, largely, first, to the promptness with which Dr. Musser acted; and, secondly, to the fact that her stomach contained little beside albuminized water. Yet, in spite of this favorable condition of the stomach, there were a few flakes of lymph already present on the stomach. A culture was taken from the abdominal cavity as soon as it was opened, and this proved, after eight days, to be sterile. I do not know that Fowler's position influenced the recovery in this case very much, in view of the facts just stated, but it certainly, at least, did no harm.

I debated for a few moments whether I should do Finney's gastroduodenostomy by means of a horse-shoe-shaped incision extending from the duodenum into the stomach. I, however, finally decided against it on account of the great thickening of the whole anterior pylorus, and because I believed it would take longer than a gastro-enterostomy by the Murphy button. For the last reason also I decided against a gastro-enterostomy with simple suture.

Up to the present time, over two months after the operation, the button has not passed, but it has not produced any trouble.

To irrigate wounds or large cavities employ solutions at or slightly above body temperature, as cold interferes with repair.
—*Internat. Jour. of Surg.*

BOVINE AND HUMAN TUBERCULOSIS.*

D. E. SALMON, D.V.M.

WASHINGTON, D. C.

In the last two reports presented by this committee considerable space was devoted to the subject of tuberculosis, and particularly to that phase of it which relates to the transmission of the disease from animals to man. It was shown from a review of the facts obtained by different methods of inquiry that the clinical evidence, the evidence of statistics, and the evidence from experimental researches all favored the conclusion that animal tuberculosis was a factor in the causation of human tuberculosis, and it was held that sanitarians should advocate and enforce measures to prevent the transmission of the disease from animals to man. This question is so important and so much uncertainty has been caused by the paper presented at the British Congress on Tuberculosis by Koch, and by his subsequent address at the International Conference on Tuberculosis at Berlin, that it is deemed advisable to continue the discussion in the present report.

The fact that tubercular material from human subjects often failed to produce serious disease in cattle was observed by a number of the earlier investigators who experimented with such virus. It was the experiments and comparative studies of Theobald Smith, however, which attracted special attention to the difference in virulence shown by tubercle bacilli from human and bovine sources when inoculated on cattle. Smith also mentioned certain morphologic and cultural differences in bacilli from these two sources, and in the location and histology of the lesions in cattle produced by such bacilli. He did not conclude, however, that bovine bacilli could not produce disease in the human subject, but said: "It seems to me that, accepting the clinical evidence on hand, bovine tuberculosis may be transmitted to children when the body is overpowered by large numbers of bacilli, as in udder tuberculosis, or when certain unknown favorable conditions exist."

Koch, however, in his address at the British Congress on Tuberculosis, went far beyond this and maintained that "human tuberculosis differs from bovine, and can not be transmitted to cattle." As to the susceptibility of man to bovine tuberculosis, he said it was not yet absolutely decided, but one was "nevertheless already at liberty to say that, if such a susceptibility really exists, the infection of human beings is but a very rare occurrence." He emphasized this view in the following language: "I should estimate the extent of infection by the milk and flesh of tubercular cattle, and the butter made of their milk, as hardly greater than that of hereditary transmission, and I therefore do not deem it advisable to take any measures against it."

This conclusion was so radically different from the views of most experimenters, and so out of harmony with facts which had apparently been demonstrated by others, that it at once aroused opposition in the congress, followed by the adoption of dissenting resolutions, and led to numerous investigations in various countries. Koch's conclusions were based on his failure to produce tuberculosis in cattle and other animals by inoculating them with tubercular material of human origin, and his success in causing progressive and fatal tuberculosis in the same kinds of animals when inoculated with tubercular material of bovine origin. With such positiveness did

* Report of the Committee on Animal Diseases and Animal Food, presented by the chairman at the meeting of the American Public Health Association, at Washington, D. C., Oct. 27, 1903.

he hold to the constant and specific difference between the human and bovine bacillus that he promulgated an experimental method of discriminating between them. Speaking of the etiology of intestinal tuberculosis in man, he said: "Hitherto nobody could decide with certainty in such a case whether the tuberculosis of the intestines was of human or of animal origin. Now we can diagnose them. All that is necessary is to cultivate in pure culture the tubercle bacilli found in the tubercular material, and to ascertain whether they belong to bovine tuberculosis by inoculating cattle with them. For this purpose I recommend subcutaneous injection, which yields quite specially characteristic and convincing results."

These important and comprehensive conclusions followed from very few experiments on animals, and apparently no effort had been made to learn to what extent human tubercle bacilli might differ in their virulence for cattle or what grades of virulence there might be among bacilli of bovine origin. Vagedes had already shown that bacilli were sometimes present in human lesions which were as virulent as bovine bacilli, but his work was ignored by Koch. In the preceding report of this committee reference was made to the work of Ravenel, DeSchweinitz, Mohler, DeJong, Delépine, Orth, Stenström, Fibiger and Jensen, Max Wolff, Nocard, Arloing, Behring, and Dean and Todd, all of whom found tubercle bacilli in human lesions which proved to have about the same virulence for cattle and other animals as had the bacilli from bovine animals affected with tuberculosis. Since that report was presented Theobald Smith has reported the finding of a human bacillus virulent for cattle, and there has been made a preliminary announcement by Kossel of the results obtained by the German tuberculosis commission. The statement of Kossel should be given the most careful consideration, coming from a commission appointed on Koch's recommendation, and evidently inclined to favor his views as far as possible in its report.

Kossel stated that the commission had tested seven cultures of tuberculosis from cattle and swine, four of cattle and three of swine. Two of these cultures proved acutely fatal in cattle after 8 to 9 weeks; four of the cultures likewise produced a generalized tuberculosis, but which certainly had a more chronic course; while one of the cultures caused only an infiltration at the point of inoculation, with some caseous foci in the adjoining prescapular gland and in one of the mediastinal glands, and there was lacking the spreading of the tuberculosis over the entire body which they were accustomed to see after the injection of cultures of bovine tuberculosis. "Hence," says Kossel, "among bovine tuberculosis bacilli there can also occur differences with regard to the virulence."

The German commission also tested 39 different freshly made cultures from tuberculous disease in man. Nineteen of these cultures did not produce the slightest symptoms in cattle; with 9 others the cattle exhibited after 4 months minute foci in the prescapular glands which were mostly encapsulated and showed no inclination to progress; with 7 other cases there was somewhat more marked disease of the prescapular glands, but it did not go so far as a material spreading of the process to the glands next adjoining; there were four cultures, however, which were more virulent and caused generalized tuberculosis in the cattle inoculated with them.

It would appear, therefore, that hereafter even Koch must admit that it is impossible always to tell the source

of a culture of the tubercle bacillus by its effects when it is inoculated on cattle. One of the bovine cultures failed to produce generalized tuberculosis in cattle and some of the human cultures did produce this form of the disease in such animals; moreover, while some of the human cultures caused no disease at all, others led to the development of minute foci in the prescapular glands, and still others to somewhat more marked disease of these glands. There were, consequently, four degrees of virulence noted in these 39 cultures of bacilli from human sources, and three degrees of virulence in the 7 cultures from animal sources. Now, if we accept the views of Koch as to the specific difference between human and bovine tubercle bacilli, and that the human bacilli produce only localized lesions in cattle, while bovine bacilli produce generalized lesions in these animals, must we not conclude that the one non-virulent bovine culture was in reality of human origin and that the animal from which it was obtained had been infected from man? This is a logical deduction, but reverses the dictum laid down at London that human tuberculosis is not transmissible to cattle. Again, how are we to explain the human cultures of medium virulence? Are they human bacilli which for some unknown reason are increasing in virulence and approaching the activity of the bovine bacillus, or are they really bovine bacilli which have multiplied in the human body until their virulence has become attenuated? In whatever manner these questions are decided, it would seem that the findings of the German commission, instead of supporting Koch's views that we can decide with certainty by the inoculation of cattle as to the source of any given bacillus, really show that this method of diagnosis is extremely uncertain in the present condition of our knowledge.

It is definitely admitted that four of the human cultures caused generalized tuberculosis in cattle; but Kossel suggests that it might be possible that the bacilli in cases of human tuberculosis under certain circumstances could likewise attain a very high pathogenic activity for cattle without being for that reason bovine bacilli. Undoubtedly, the German commission is confronting the two horns of a dilemma, either one of which is fatal to the views of Koch as stated with such positiveness at London. If we accept this suggestion thrown out by Kossel, we must conclude that Koch was wrong in his assertion that human tuberculosis can not be transmitted to cattle, and thus with one blow we destroy the entire experimental support which he had for his argument before the British Congress on Tuberculosis. And if, on the other hand, we accept the conclusion which follows from the principle laid down by Koch for the discrimination between human and bovine bacilli, and which appears to be favored by Kossel, we must admit that bovine tuberculosis is an extremely important factor in the etiology of human tuberculosis. Of the 39 cases of human tuberculosis tested, 4, or over 10 per cent., were virulent for cattle and would be classified as of bovine origin; but these 4 cases were all found among the 16 cases of tuberculosis in children which the commission investigated; hence, it is plain that 25 per cent. of the cases tested of tuberculosis in children would by Koch's method be classified as of bovine origin.

In the Bureau of Animal Industry two distinct lines of experiments have been carried on, in order that one might serve as a check against the other. There has been, however, no discrepancy in the results. De Schweinitz, in the biochemic division, has isolated nine cultures from human tuberculosis. Two of these were

derived from human sputum, three from cases of generalized tuberculosis in adults, and four from cases of generalized tuberculosis in children. By comparing these cultures with a newly isolated virulent culture of bovine tuberculosis, there were found among them two cultures from children which were identical in their cultural and morphological characters with the bovine bacillus. These cultures also killed rabbits and guinea-pigs in as short a time as did the bovine bacillus. Hogs which were inoculated subcutaneously with these two cultures from children died of generalized tuberculosis. Two calves, weighing over 30 pounds each, were inoculated subcutaneously with these virulent human cultures and as a result developed generalized tuberculosis. A yearling heifer inoculated with one of the cultures showed generalized tuberculosis when killed three months after inoculation. Both the cattle and the hogs had been tested with tuberculin and found to be free from tuberculosis before the inoculations were made. It is important to observe in this connection that two out of four, or 50 per cent. of the cultures obtained from cases of generalized tuberculosis in children, proved virulent for cattle.

Mohler, working in the pathological division, has obtained three very virulent cultures of tubercle bacilli from the human subject. A goat inoculated subcutaneously with one of these cultures died in 37 days with miliary tuberculosis of the lungs involving the axillary and prescapular glands. This bacillus was obtained from the mesenteric gland of a boy. Of still greater interest is a bacillus isolated by Mohler from human sputum. A goat inoculated subcutaneously with a culture of this germ died in 95 days of pulmonary tuberculosis. A cat inoculated in the same manner died in 23 days of generalized tuberculosis. A rabbit similarly inoculated died in 59 days of pulmonary tuberculosis. Another rabbit inoculated with a bovine culture for comparison lived 10 days longer than the one inoculated with this sputum germ. Mohler also inoculated subcutaneously a 1-year-old heifer with a culture derived from the tubercular mesenteric gland of a boy 4 years of age. This culture was always refractory in its growth under artificial conditions, and the bacilli were short stubby rods corresponding in appearance with the bovine type. At the autopsy, held 127 days after the inoculation, the general condition was seen to be poor and unthrifty, and large, hard tumors were found at the points of inoculation. On the right side, the swelling measured $3\frac{1}{2}$ by 5 inches, and the corresponding lymph gland was $2\frac{3}{4}$ inches long by $1\frac{3}{4}$ inches in diameter. This gland contained numerous calcareous foci, one of these at the apex being an inch in diameter. The lesions on the left shoulder of the animal were very similar to those found on the right side, but the dimensions of the tumor were slightly less. The lungs presented an irregular mass of tubercular nodules, and 7 or 8 grape-like nodules were seen on the parietal pleura. Bronchial and mediastinal lymph glands contained numerous tubercular foci, and the pericardium, peritoneum, spleen and liver were also affected.

In order to throw some light, if possible, on the morphological constancy of the different types of tubercle bacilli, Mohler was requested by the chairman of your committee to make comparative studies of bacilli from various sources, and which had been passed through various species of animals, by making the cultures on dog serum after the method described by Theobald Smith. Some important results have been obtained.

One culture of human bacilli which had morphological and cultural peculiarities similar to those of the bovine bacillus, and which only produced local lesions in cattle, was passed through a series of 5 cats. It was then found to be completely changed in its morphological characters, the rods being elongated, slender, more or less beaded and entirely of the human type. But far from decreasing in virulence, as might be expected from their morphological appearance, this bacillus had so increased in its pathogenic activity that it now produced generalized tuberculosis in a cow. This cow was inoculated subcutaneously in front of each shoulder with 2 c.c. of a salt solution emulsion of the tuberculous omentum of the last cat of the series. The cow rapidly lost flesh, had a temperature of 104° F., with the point of inoculation and adjacent glands greatly swollen. The autopsy revealed generalized tuberculosis, involving the lungs, mediastinal glands, spleen, liver and kidneys. Tubercle bacilli of the bovine type obtained from the mesenteric glands of a sheep, hog and cow were similarly transformed in their morphological appearance after being passed through a series of cats and recovered on dog serum. These bacilli also increased in virulence, as the last cat of the series invariably succumbed in a shorter time than the first of the series.

These experiments and observations indicate that the types of tubercle bacilli are very inconstant, and that under suitable conditions they readily change both in morphology and in virulence. A similar conclusion was reached by other investigators in working with the avian and piscine types of tubercle bacilli several years ago, and was reasonably to have been expected with the human and bovine types.

It must be plain to all from these recent developments, that too much has been made of the slight differences in cultural characteristics, in morphology, and in virulence which have been observed in some cases in comparing the human and the bovine bacilli. The observations were interesting and it was important that they should be followed up until their significance was made entirely clear; but it was an almost unpardonable error, from a sanitary point of view, to promulgate sweeping generalizations, calculated to arrest and abolish important measures for preventing human tuberculosis, before the soundness of these generalizations had been established by a thorough course of experimentation.

When Koch said in the British Congress on Tuberculosis that he should estimate the extent of infection by the milk and flesh of tubercular cattle, and the butter made of their milk, as hardly greater than that of hereditary transmission, and that he therefore did not deem it advisable to take any measures against it, he went far beyond what was justified by any experiments or observations which he reported, and he did an immense amount of harm which will be manifested for years to come to those who endeavor to guard the human race from the dangers of animal tuberculosis. The researches to which your committee has alluded make these dangers more definite and certain than they have appeared before, and sanitarians should therefore most earnestly endeavor to counteract the erroneous and harmful impression which was made by Koch's address at London and his subsequent address at the International Conference on Tuberculosis at Berlin.

Now that the conclusions of the London address are shown to be incorrect and based on insufficient experimentation, it is well that some attention should be given to the Berlin address in which certain rules were laid

down for testing all clinical evidence before it can be accepted as indicating the transmission of tuberculosis from animals to man. While your committee agrees that it is advisable to scrutinize clinical evidence as carefully as possible before it is accepted as influencing the decision of such an important question, consistency demands that we should accept clinical evidence bearing on the transmission of bovine tuberculosis to man on the same terms that we accept evidence bearing on the transmission of tuberculosis from man to man. For instance, Koch said in his London address, "So the only main source of the infection of tuberculosis is the sputum of consumptive patients." How does he know this? Has he any evidence to support this conclusion which will stand the requirements which he has laid down for those who oppose his views with reference to the transmission of animal tuberculosis to man? If so, this evidence has not yet been presented. Your committee believes that the principal source of tuberculosis in man is infection from other tuberculous human subjects, but it also believes that a considerable proportion of the cases of tuberculosis in man is due to infection from animal sources; and in weighing the evidence by which we are to fix more rigorously the relation which these two sources of infection bear to each other, it is logical that we should apply the same rules of evidence in each case.

The rules which Koch formulated for testing the evidence as to bovine infection are given below, with such brief comments as your committee feels they deserve.

"1. Certain proof of tubercle, and where possible the primary focus must be supplied."

To this condition no objection is raised, but it should be remarked that the location of the primary focus, which is made so much of, is of little value in determining the origin of the infection. It has been clearly shown by various investigators, whose work has been referred to in the preceding reports of this committee, that tubercle bacilli may pass through the walls of the intestine without producing any lesion at the point of entrance, and that their first point of lodgement and multiplication may be the mesenteric glands, the liver, spleen or even the lungs. On the other hand, Koch has told us that the location of the primary lesion in the intestine is of no significance because human as well as animal bacilli may be swallowed and may be the cause of such lesions. It is, therefore, not at all apparent that this is an essential point.

"2. Other sources of infection must be excluded with certainty."

This condition appears to exclude all clinical evidence bearing on the source of tubercular infection. How is it possible to prove that any given individual has not been exposed to the bacilli of human tuberculosis? And, for the same reason, how is it possible to prove that any given individual has not been exposed to the bacilli of animal tuberculosis? It may be said that this consumptive person had habitually been in a room with another consumptive patient and was infected from that patient. But how can you prove that this person never ate any tuberculous meat, never partook of any tuberculous milk, never ate any butter containing the tubercle bacillus, never had opportunity to be indirectly infected from the hands of cooks or from table utensils which had been in contact with tuberculous meat, milk, or butter, and was never exposed to the infection scattered in so many ways by tuberculous animals? It is impossible to exclude with certainty all these sources of infection, and if we apply the condition under discussion to the evidence on

which Koch based the assertion that the main source of the infection of man is the sputum of consumptive patients, we would find no part of it that would stand the test. Now, is it logical to accept clinical evidence, as to the transmission of tuberculosis from man to man, by one set of rules, and to require an entirely different set of rules when considering evidence as to the transmission of the same disease from animals to man? Your committee is of the opinion that clinical evidence should be regarded as valuable in confirming experimental researches, or in indicating the probable facts where experimentation has been insufficient or is impossible, but that it is too much to expect that clinical evidence will be forthcoming as to the transmission of tuberculosis which will exclude all possible sources of error.

"3. In each case of alleged infection from milk affected with bovine tuberculosis the condition of the rest of the people who have taken the same milk should be borne in mind. These fellow consumers form to a certain extent a control experiment, and if of the numerous people who have drunk the suspected milk only a single one sickens, this weighs decidedly against the belief that this one person was infected by the common food."

This condition, also, is regarded by your committee as illogical and impossible of general application without discarding all clinical evidence on this question. If we apply it to the alleged cases of sputum infection, we find it is seldom, indeed, that the disease in more than one person can be traced with any probability to the same tuberculous patient, although scores of persons may have been exposed to that patient. Most people who escape tuberculosis have nevertheless been exposed many times to tuberculous patients, but this is not sufficient reason for concluding that others have not contracted the disease by similar exposure. Tuberculosis is a disease which is communicated with such difficulty among people under ordinary conditions of life, either by exposure to diseased persons or to tuberculous food, develops so slowly, and varies so much in the period which elapses before symptoms are observed, that it is only very rarely that groups of cases are observed which even appear to have originated from the same source. And yet, how erroneous it would be to exclude clinical evidence suggesting contagion because only one of those exposed to a certain consumptive had contracted the malady.

"4. The source of the milk should be attended to. Since in recent years it has become more and more evident that milk containing tubercle bacilli is yielded by such cows as suffer from tuberculosis of the udders, the general statement that some one has drunk milk from a cow suffering from tuberculosis no longer suffices to prove to us that bovine tuberculosis bacilli have really reached his digestive organs. It must be from a cow suffering from tuberculosis of the udder, and therefore a statement on this subject should not be wanting in a report on milk infection if it is said to be complete."

Your committee finds this statement of fact to be incorrect, since the tendency of recent investigations by competent persons is to make it more and more evident that cows with tuberculosis may yield milk containing tubercle bacilli when the udders present no signs of the disease. The argument is, therefore, antiquated. Of the many experiments that have been made to determine the proportion of tuberculous cows which yield infectious milk, the average results are about 15 per cent, while the cases of tuberculosis of the udder are about 2 per cent. In recent investigations made by Mohler of the Bureau of Animal Industry with 56 reacting cows, it was

found that 12 of these, or 21.4 per cent., at one time or another during the experiment gave milk which contained virulent tubercle bacilli. Undoubtedly cows with tuberculosis of the udder yield milk containing a larger number of tuberculosis bacilli than do those in which the udder remains unaffected and are, therefore, more dangerous, but it has been clearly shown that other tuberculous cows may yield virulent milk.

So much for the conditions laid down by Koch for testing clinical evidence bearing on the transmission of tuberculosis from animals to man. The effort to rule out all such evidence by applying impossible tests does not meet with our approval, but fortunately clinical evidence is no longer necessary for deciding the question. The subcutaneous injection of cattle with pure cultures of the tuberculosis bacillus from human sources, which, according to Koch, "yields quite specially characteristic and convincing results," has proved, in the hands of the German commission no less than in those of independent investigators, that bovine tuberculosis is communicable to man. It will require much work to decide with even approximate accuracy the proportion of human tuberculosis caused by animal infection; but the fact that 25 per cent. of the cases in children investigated by the German commission, and 50 per cent. of similar cases investigated by De Schweinitz showed by this test that they were caused by animal infection, is sufficient to convince us that measures should be taken and enforced at once to guard against infection from this source.

In concluding your committee desires to express its appreciation of the prompt, intelligent and indefatigable work of those scientific investigators who took up this question immediately after it was brought to the front at London, and who, in the comparatively short period of two years, have furnished the material for definitely settling it, so far as the principal contention is concerned.

THE CONDITION OF THE BLOOD IN PATIENTS SUFFERING FROM PULMONARY TUBERCULOSIS.*

JOHN M. SWAN, M.D.

Demonstrator of Osteology, University of Pennsylvania; Pathologist, St. Mary's Hospital; Instructor in Clinical Medicine, Philadelphia Polyclinic.

PHILADELPHIA.

The study of the blood in the patients who furnished the material for this paper was made possible by the kindness of Dr. Julius L. Salinger, who placed his tuberculosis ward in the Philadelphia Hospital at my disposal. The paper is based on the counting of the erythrocytes, the counting of the leucocytes, the estimation of the hemoglobin, the differential count of the leucocytes and the search for the iodine reaction of the leucocytes in twenty-five cases of pulmonary tuberculosis, and one case of extensive cavity formation in the lung due to bronchiectasis.

In studying the results of these blood counts I shall adopt the classification of Grawitz,¹ who divides the disease into three stages. A first stage, in which there is a beginning affection of the apex, without cavity formation; a second stage, in which there are symptoms of cavity formation with slight fever or without fever; and a third stage in which hectic fever is present.

FIRST STAGE.

There is one patient (26) who represents the first stage of consolidation of the apex, without cavity for-

mation. This patient showed a slight reduction of the erythrocytes to 3,980,000; a normal number of leucocytes, 10,000; and a moderate reduction of hemoglobin to 70 per cent. This patient improved very much under treatment, and was finally discharged without symptoms and with few physical signs. As his general condition improved there was a corresponding improvement in the blood picture; the erythrocytes rose to 4,580,000 and the hemoglobin increased to 82 per cent. With the improvement of the blood condition, the leucocytes rose to 16,600.

An examination of the differential leucocyte count of this patient shows that the relative proportion of the different cells was not changed. The treatment, under which the patient greatly improved, apparently caused an increase in the eosinophiles from 1.6 to 3 per cent.

SECOND STAGE.

There are eight patients (2, 4, 5, 6, 10, 14, 20, 22) representing the second stage of the disease. In these cases the erythrocytes were usually normal in number. In four instances they were present in greater numbers than 5,000,000 to the cubic millimeter, and in one instance they only numbered 3,680,000. The range of the erythrocyte counts was from 6,710,000, in a negro who had complained of cough for twelve years, to 3,680,000, also in a negro, who had been ill for two and one-half months, and who had albuminuria.

Seven of these patients showed a leucocytosis, varying from 34,000 to 11,120, and one showed a leucopenia of 7,100. The latter patient had carried a cavity in his right lung for a number of years and the disease was stationary.

In these patients the differential counts showed, as a rule, a percentage of polymorphonuclear cells above 80, the highest being 93.4. In three cases the polymorphonuclears formed 70.4 per cent., 73.6 per cent., and 78.8 per cent of the leucocytes, respectively. In one patient the first examination was made during an asthmatic paroxysm and showed 80.8 per cent of polymorphonuclear cells. A second examination was made about four weeks later, after the asthmatic paroxysms had ceased, and gave 78 per cent. of polymorphonuclear cells. Eosinophile cells were present in all of these patients except two, in proportions varying from 3 per cent. to 0.2 per cent. The two patients from whose blood these cells were absent left the hospital before a second examination could be made and their ultimate fate is unknown.

The hemoglobin varied from 90 per cent to 63 per cent., although all patients showed some pallor of the skin and the mucous membranes. Indeed, in some instances the percentage of hemoglobin was at first thought to have been erroneously estimated until more than one trial showed the reading to be accurate.

THIRD STAGE.

There are fifteen patients (3, 7, 8, 9, 11, 12, 13, 15, 16, 17, 18, 19, 21, 23, 25) who may be placed in the third stage of the disease—that of hectic fever. In these patients the erythrocytes were usually reduced in number, although a normal number of these cells was occasionally found. The counts ranged from 5,710,000 to 2,890,000, with an average of about 4,019,130. The highest count was obtained from a patient who was rapidly improving under treatment; indeed, he might be placed in the second stage of the disease but for the fact that a former count was made while he was having marked hectic fever. The leucocytes were usually in-

* Read before the Philadelphia County Medical Society.

1. E. Grawitz: *Deutsche med. Woch.*, 1893, No. 51, p. 1374.