narrowing with an infection of the urinary tract with the result of a stricture at one of these congenitally narrow places. That can be granted if all ureters are congenitally malformed, but we are not ready to grant this. If, however, there is an inflammatory tract or stone in the ureter or anything to cause trauma at one of the points of narrowing the result is stricture at that point.

Dr. Furniss has drawn attention to the relative frequency with which we see acute kidney infection following disease elsewhere—following operation, furunculosis, typhoid fever or acute infectious diseases, tonsillitis, etc.—and as I understand him, he says that most of the ureteral strictures are secondary to these infectious diseases elsewhere in the body. I agree with him. I wish to go further and state that some of these ureteral conditions occur without any infection, that is to say, they are lymphatic toxemias. They come from some distant focus in the body. In December, 1910, I showed before the Southern Gynecological and Obstetrical Association a series of cases of urethritis and ureteritis due to tonsillitis and I asked the members to bear the matter in mind and either disprove or prove my position. Since then I have had a sufficient number of these cases of chronic ureteritis to feel thoroughly convinced that my standpoint in that matter is right, and that a large number of these cases of stricture of the ureter come from toxins of the tonsils. Another thing rather pointing to the truth of that is the fact that a large percentage of cases are infected by the *Staphylococcus aureus* rather than by the colon bacillus, although the colon bacillus causes more than one-half of the infections of the urinary tract.

We do not know much about the treatment. No one has had enough experience yet to be able to state anything positive about it. Cases of stricture near the lower end of the ureter seem to have a tendency to have a long section of the ureter strictured, whereas, near the pelvic brim and nearer the kidney, the strictures are likely to be annular. This I think we shall be able to control by using bulb catheters and overstretching the strictures. I am getting some positive cures of these annular strictures simply by that treatment. With those having the long section involved near the bladder wall I believe it is necessary to go in extraperitoneally, cut them off above the seat of stricture and implant in the bladder. I have not had much success in dealing with them. Some can be dilated, but not all. I had one just last week and the only thing I could get in was a little probe-like searcher. After getting by the stricture, to get out again I had to bend the handle of my probe and push the Kelly clamp off over the probe and with one finger in the vagina pull as hard as I could. The stricture was so dense that it had grasped the end of the probe. Dilatation gives little hope. We shall have to cut the ureter off and place it either in the bladder or in the kidney.

Dr. H. D. FURNISS, New York: In my cases of stricture of the lower end of the ureter there has been little if any obstruction to the passage of small catheters, and most of them were diagnosed by radiography after the injection of collargol or argyrol. Otherwise we would have overlooked them except in the cases in which there was marked dilatation of the ureter above the stricture and retention of an appreciable amount of urine. Cases of renal pain with purulent urine following a few days after operation, I explain as being due to infection of the renal pelvis and ureter, with consequent swelling of the ureteral mucosa and obstruction thereby to the ureter. In these, pelvic lavage with silver nitrate has been successfully used. As Dr. Hunner states, obstruction of the lower end of the ureter does occur with pus-free urine. I think that the majority of these cases are in all probability due to lymphatic involvement of the vesical end of the ureter from a cystitis. Dilatation of the lower end of the ureter is easy, but as Dr. Hunner says, the results are unsatisfactory as the ureter will not remain dilated. I have a patient whose ureter requires dilatation every two weeks. When I return I shall inject argyrol, and make a radiogram to determine the exact site of the stricture and the amount of dilatation above. In this case I contemplate making a ureterovesical anastomosis on account of the unsatisfactory results from dilatation. The pain of the dilatation

itself is not severe, but that following three hours afterward and lasting three or four hours is; this is, I think, due to the obstruction of the ureter following the trauma. During the very painful period there is oliguria, which is followed later by a relative polyuria. Ureteral obstruction in women is common, and pyelography is the most accurate method of diagnosis.

A CASE OF SUSPECTED PULMONARY BLASTOMYCOSIS

JOHN H. SNOKE, M.D., AND E. J. STRICK, M.D.
SIO-KHE, CHINA

A Chinese woman, aged 36, came to the dispensary of the Neerbosch Hospital in March. She was so weak and short of breath that she was unable to walk alone. Our first impression was that this was a far-advanced case of tuberculosis. The patient was emaciated, anemic and very short of breath; she coughed almost constantly and raised large quantities of mucopurulent sputum. Both apices and the left lung were normal. There was an area of dullness about 5 cm. in diameter just above the right mammary region in the nipple line. Over this area were heard râles of nearly every description. Patient complained of no pain, but of weakness and an inability to sleep on the affected side on account of coughing. We examined the sputum as many as twenty times for tubercle bacilli but with constant negative results. We noticed, however, the constancy of certain spore-like bodies with double refractile capsules and granules. This led us to suspect blastomycosis. Up to this time, a period of about ten days, she had been taking emulsion of cod-liver oil and hypophosphites with guaiacol. Her fever ran from 101 to 103 and was more or less constant, except that there were evening remissions instead of the usual morning remissions. The pulse was 112 and rather weak. Under the cod-liver oil and guaiacol treatment there was no change. After we had decided that we were not dealing with tuberculosis but possibly with blastomycosis, we used copper sulphate as directed by Bevan¹ (quarter-grain doses, three times a day) and the result was marvelous. After twenty-four hours the temperature had dropped to 99, and in three days it was normal in the evening, with a fraction of a rise in the morning; but within a week both pulse and temperature were normal, the constant cough was much relieved, and the quantity of sputum greatly reduced. Its character, however, remained much the same. The patient was soon able to sleep on the affected side and gained in weight and strength very rapidly. We are sorry we have to report that some cough still continues and that we still find the blastomycetes in the sputum. Nevertheless the patient has gained so much in the three months of treatment that she is able to walk four miles to and from the dispensary.

Castellani² reports several cases which he calls bronchomycosis, the description of which answers quite closely to our own, but not entirely. He claims cures by the use of potassium iodid in large doses three times a day.

We have had ten cases of what seemed at first sight like tuberculosis, in the last three months, but the microscope showed tuberculosis in four cases, blastomycosis in five and one negative. It would seem therefore that blastomycosis is not rare in China.

The chief remedial agent employed successfully, says Ormsby,¹ is potassium iodid, first advised by Bevan. The drug nearly always produces marked results and in some cases entirely cures the disease. Doses as large as 600 grains a day are often required. It should be given in large dilution and should be continued for a long time or until no renewed activity occurs after its withdrawal. More recently Bevan has advised copper sulphate. In our opinion these drugs are only palliative, and as yet we have no cure.

1. Ormsby, Oliver: Blastomycosis, in *Lexer and Bevan's General Surgery*, 1908, p. 376. We refer those interested in the subject to an account of the pathology and history of the disease and a very complete bibliography appearing at the end of that article.

2. Castellani: *Philippine Jour. of Sci.*, July, 1910.