and hemorrhages continued. A movable ovarian cyst, the size of an egg, was discovered and operation was advised on account of the danger of torsion of the pedicle. This was easily effected, the unpleasant symptoms promptly disappeared, and the subsequent course of the pregnancy was uneventful.

Anatomy of Chronic Parametritis.—FREUND (*Centralblatt für Gynäkologie*, No. 43, 1902) presents the results of his recent studies bearing on this subject, which confirm his previous views with regard to the relation of this condition to certain reflex nervous symptoms. Sections through the diseased tissues show marked changes in the pelvic ganglia included within them. Besides being surrounded and compressed by the cicatricial tissue, their cells are shrunken, lose their normal shape, and the nuclei mostly disappear. The writer believes that this subject offers a fruitful field for investigation, and that various reflex neuroses usually referred to hysteria may be explained by these changes in the intra-pelvic ganglia.

Diagnosis of Genital Tuberculosis.—SELLHEIM (*Centralblatt für Gynäkologie*, No. 43, 1902) believes that the probable tubercular character of obscure pelvic disease in an individual may be inferred if it is present in other organs, or in members of the same family. If isolated nodules can be detected in the pelvic peritoneum, and especially at the uterine end of the Fallopian tube, the diagnosis is almost certain. The tube has a characteristic rosary-like feel, the nodules being of firm consistence.

Curettement of the uterine cavity and microscopic examination of the tissue removed should never be omitted in cases of suspected tuberculosis of the tubes and pelvic peritoneum, since in this way important information may be obtained bearing upon the questions of prognosis and treatment. If such a thorough examination is made the writer believes that in the majority of cases it is possible to make a diagnosis of genital tuberculosis.

He reports sixty-five cases observed at the Freiburg clinic in the course of eight years; twenty-eight of these were treated palliatively, with satisfactory result as regards the relief of symptoms and arrest of the disease; thirty-seven capital operations were performed, with equal success. Total extirpation of the uterus and adnexa is preferable.

Treatment of Pelvic Suppuration.—JUNG (*Centralblatt für Gynäkologie*, No. 43, 1902) reports 122 cases of laparotomy from the Greifswald clinic, with a mortality of 18 per cent., which he explains by the fact that these included the worst cases, small pus tubes and ovaries, with slight adhesions, being removed per vaginam. Another explanation of the high death rate was afforded by the virulence of the pus, which was found to be sterile in only 14 out of 78 cases in which it was examined bacteriologically. Streptococci were found twenty-six times, gonococci twelve times, colon bacilli six times, and tubercle bacilli thirteen times. Vaginal section and drainage are now practised at the clinic, the diagnosis being greatly aided by examination of the blood. In 21 recent cases the patients were cured by this method.