the papules were much larger. They were crescentic in shape and tended to become confluent; this eruption appeared in two patients and within twenty-four hours after the use of the serum in both instances. It caused little inconvenience. At the point of the introduction of the injection, local anesthesia was at first produced by means of ethyl-chlorid spray; this sometimes results in a deep and extensive ulceration about the point of introduction of the needle, as the local nutrition was interfered with in consequence of pressure after the freezing. After this experience we gave the patient a few whiffs of chloroform and discarded local anesthesia. I think that it is still too early to judge of the curative effects of the serum because of the great variability of the virulence of the epidemics. In the present epidemic the percentage of recoveries is high. But it seems to me that we are justified in using the serum because of the immediate primary change in the symptoms and the favorable turn produced evidently by the serum. There is the immediate relief of the headache, a decline in the fever, a marvelous clearing up of mental symptoms, a subsidence of the delirium, all caused by this serum, reminding one of the clearing up of the symptoms in diphtheria under the influence of the antitoxin.

DR. D. E. ENGLISH, Millburn, N. J.: Does Dr. Ladd give 30 c.c. of serum intraspinally at the first injection? Does he give the amount of serum that corresponds to the amount of cerebrospinal fluid removed? Or does he try to take away more fluid than he wishes to give of the serum? Is the subcutaneous dose larger than the intraspinal dose?

DR. O. M. GILBERT, Boulder, Colo.: I chanced to see a case of cerebrospinal meningitis in a specially severe epidemic in which the streptococcus was more in evidence than the *Diplococcus intracellularis*. The bacteriologist who made the examination first reported back that it was a case of pure streptococcus; but later the diplococcus was found. The giving of 20 c.c. streptolytic serum resulted in recovery. We had no Flexner serum. Eighty per cent. of our patients had been dying. Would not the streptolytic serum be of value when both germs are demonstrated?

DR. L. W. LADD, Cleveland: I have had little experience in the use of the Flexner serum in other than the meningococcus type of cerebrospinal meningitis. I have used it in several cases of tuberculous meningitis, but without any result whatever. In one case of meningitis in which streptococci and an unidentified streptothrix were associated I used the serum. The patient was moribund when seen and died three hours afterward. In Wheeling, West Virginia, I saw a case which I thought one of tuberculous meningitis. Lumbar puncture was performed and an increased amount of spinal fluid obtained, under considerable pressure, but examination of the fluid showed a normal cell picture and no micro-organisms could be found on coverslip or by culture. Following the puncture and giving of serum this patient recovered, but was not included in the reported series. I have used the antistreptococcus serum in one case only. This was the case of a child who had had otitis media followed by a mastoid infection and later complicated by meningitis. Lumbar puncture was performed and a very thick purulent fluid was obtained in which streptococci were present in large numbers. The antistreptococcus serum was used repeatedly and was followed by some apparent benefit, but death ensued in spite of the serum. The cases which I reported were all meningococcus infections. There were no mixed infections and with one exception the meningococcus was present in all. In answer to Dr. English's question I generally give the serum intraspinally, but sometimes both intraspinally and subcutaneously. When little fluid is removed by lumbar puncture I give more subcutaneous. The amount given intraspinally depends somewhat on the amount of fluid withdrawn. If a large amount is obtained, 30 c.c. are injected intraspinally. I follow the rule of withdrawing the fluid until it flows, drop by drop, as under normal conditions. Serious symptoms have followed the withdrawal of too much fluid at one time. In answer to Dr. Wells' criticism of my paper, I have purposely given statistics, in the first place including all cases irrespective of their condition when first seen and irrespective of the length of time which had elapsed after onset of the case, and then ruling out such

cases as were manifestly unfair tests of the efficiency of the serum. Surgeons have had great trouble in educating the medical profession to turn over to them their cases of appendicitis for operation, before symptoms of general peritonitis developed, in order that the result might be most favorable to the patient. The same necessity of early diagnosis and early use of serum is present in the treatment of this disease, and comparative statistics show this strikingly. Ruling out the moribund patients in those seen seventy-two hours or less after onset, 93 per cent. recovered. All of the patients, 5 in number, seen twenty-eight or less hours after onset recovered. Sometimes, of course, it is impossible to make a diagnosis as soon as desirable. With regard to the nature of the substances active in the serum, I do not think that Dr. Flexner has absolutely committed himself. It certainly is true that phagocytosis is stimulated, and in view of the fact that the meningococci after the use of the serum were reduced in numbers and stained poorly and that the patients treated cleared markedly in mentality, it seems reasonable to suppose that the serum is both bactericidal and antitoxic in its action.

DR. FRANK J. SLADEN, Baltimore: At a recent symposium on this subject before the American Pediatric Society Dr. Flexner reported an analysis of 380 cases in which the serum had been used. The mortality for 247, the number in this country, is about 27 per cent. At this time also Dr. Dunn of Boston reported 40 cases with 9 deaths, 22.5 per cent. mortality. Dr. Churchill of Chicago had a mortality of 22 per cent. for 9 cases. And Dr. Wilkinson of Washington, D. C., reported 3 deaths in 10 cases. The comparison of these statistics with those of the New York epidemic is striking. The mortality for 2,305 cases treated without serum was between 80 and 90 per cent. Dr. Flexner has committed himself, stating that he believes the power of the serum is chiefly bactericidal. He is a pathologist, and his opinion is based mainly on laboratory experiments. Clinically, however, the rapidity with which the symptoms and signs clear up impress us with the importance of the antitoxic property.

DR. C. F. HOOVER, Cleveland: One point brought out in this discussion should be appreciated by every physician, viz., the need of giving the serum as early as possible to anticipate the encephalomyelitis which results from meningo-arteritis which has lasted several days.

URETERO-VESICAL IMPLANTATION. A NEW METHOD OF ANASTOMOSIS.

WITH REPORT OF CASES.

R. L. PAYNE, JR., M.D.

NORFOLK, VA.

CASE 1.—*Patient*.—Annie G., aged 30, colored, was admitted to St. Vincent's Hospital, January 30, suffering with large bilateral pus tubes. The pus sacs were removed through an abdominal incision and drainage instituted. The wound healed up to a small sinus, when, on the fourteenth day following operation, urine began to flow from the sinus and a diagnosis of uretero-abdominal fistula was made. By the use of the cystoscope and ureteral catheters it was determined that the left ureter was the one affected, and two weeks later I performed the following operation:

Operation.—An incision was made from the lumbar region obliquely forward to the middle of Poupart's ligament, cutting down to the lumbar and subperitoneal fat behind and to the peritoneum in front. The peritoneum was then gently stripped inward from off the psoas muscle and iliac bone until the ureter was located at the bifurcation of the iliac artery. The ureter was deflected with the peritoneum. It proved to be very large and thickened and stripped from the membrane very easily. It was then freed down to the fistula tract, where the opening of the ureter proved to be very small and strictured. When stretched tight it lacked two inches of meeting the bladder, and I found that the injury to the ureter was behind the base of the broad ligament. The vesical stump was buried and hidden in a mass of inflammatory

exudate, and no attempt was made to free it. I then stripped the peritoneum backward from the bladder, loosened the bladder from its attachments on either side, and with the aid of a sound passed through the urethra into the bladder by an assistant, pulled the organ to the left and upward, thus elongating it into a diverticulum effect. The bladder was then fixed in this shape and sewed with catgut to the pelvic fascia both above and below the external iliac artery. This elongation upward of the bladder permitted a good approximation of the shortened ureter and the anastomosis was made by the method herein described. When the anastomosis was completed I found that too much of the ureter had been used for the anastomosis and that the duct was rather taut. I was afraid to leave it thus, fearing that the anastomosis might tear out, so I proceeded to lower the kidney a sufficient distance and fix it to relieve the strain. A small gauze drain was passed through the abdominal opening of the original sinus tract backward to the retroperitoneal point where the end of the ureter had been dissected out and another small drain placed from this point out through the loin space. Thus there was a gauze drain running straight through the patient, beginning in front just below the navel and emerging in the lumbar space behind, the whole drainage tract being extraperitoneal. The muscles and fascia were closed in layers by chromic gut, and the skin by interrupted silk sutures.

Postoperative History.—The patient was catheterized every four hours for the next five days, and at the end of this period she voided without much pain. A week later this difficulty had about disappeared, and it is interesting to know that a normal bladder can undergo such a transposition without interfering with the urinary function. The drain passed into the sinus tract was removed on the third day, and this portion of the drainage canal healed up in three more days. The drain in the loin, however, continued to discharge a seropurulent matter for about two weeks, and I thought that it was a leakage from the vesical stump of the divided ureter. This stopped, however, and with the exception of three small stitch abscesses the wound healed rapidly. The patient was walking around at the end of the fourth week and has had no trouble up to the present time.

Witzel was the first to elongate the bladder for a shortened ureter, and Baldy has since reported a case in which he sewed the bladder to the stump of the broad ligament. Several years ago Bovee successfully lowered the kidney in a dog for a shortened ureter but as far as I can find, the case herein reported is the first instance in which the procedure has been carried out on a human being.

Since Simon in 1869 first recommended nephrectomy for ureteral fistula, many prominent surgeons of the world have been striving to perfect operative means which would obviate the necessity for this radical measure.

Several varieties of uretero-ureteral anastomosis have been devised, none of which has proven entirely satisfactory or simple in the hands of the average good surgeon, and so the abnormal varieties of implantation into the colon, rectum and skin have been practiced in a limited number of cases.

It is certain that nephrectomy has been and is still too readily looked on as the proper panacea for the awful conditions resulting from ureteral fistula, and many a healthy kidney has been sacrificed without previous effort to cure the fistula by more conservative methods.

Of all the various plastic procedures for the relief of ureteral fistula, ureterovesical implantation is undoubtedly the best, for it reestablishes a natural state of things, cures the fistula and above all else conserves the kidney. So far as the kidney is concerned, the same can not be said of grafting into the bowel or on the

skin, for infection invariably follows and nephrectomy becomes a necessity.

Of the two latter measures bowel implantation is preferable to skin implantation, but since both are done, except in exstrophy, because of a shortened ureter, neither procedure is justifiable provided the pedicle of the kidney is long enough to permit of its being lowered.

Van Hook says that intestinal implantation is unjustifiable and Byron Robinson reports that in his cases of intestinal implantation in animals all the subjects died from peritonitis.

The objections to vesical grafting in cases of ureteral fistula have been that it could be applied only to those cases in which the injury is very close to the bladder; for a ureter severed at the base of the broad ligament, behind the uterus, or higher up, would not permit of the renal stump approximation with the bladder. It is true that the ureter is very elastic and hence amenable to much stretching, but traction largely reduces the possibilities of anastomotic union and at the same time tends to lessen the lumen of the duct and thus contributes a factor in the induction of hydronephrosis, the greatest danger of ureteral anastomoses. This source of danger can be overcome by lowering the kidney as was practiced in my case.

Fig. 1.—Ureter split and traction sutures passed.

The other objections to the various modes of ureterovesical implantation are, first, stenosis, which is apt to occur at the site of vesical grafting and, second, the danger of infection from ascending septic processes. Regarding these two points I believe that the method of anastomosis herein described and illustrated in Figures 1, 2 and 3, presents points in its favor which tend to obviate these results.

By the common method in vogue the severed ends of the ureter is united flush with the bladder wall, mucous membrane to mucous membrane and when the process of cicatrization takes place a resulting contraction and stenosis must follow. To correct this, several operators have followed the plan of Van Hook, splitting the ureter on one side only, flaring the end and thus uniting the ureter into the bladder.

By the method adopted by myself the ureter is split up on two sides, thus forming separate flaps and when these are drawn down into the bladder by traction sutures and fixed on each side of the vesical incision there is a resulting ureteral opening which is larger than the caliber of the tube and around which there are no

sutures or united mucous margins to undergo cicatrization and subsequent contraction. The union of these flaps to the mucous membrane of the bladder was at first a matter of doubt, but I can assert that in the cases in which I have used this method the flaps do unite firmly and at the same time atrophy and smooth off to the bladder wall. Most probably the fibrous union begins in an erosion around the traction sutures but in any event firm union does take place, for with the cystoscope I demonstrated not only the fibrous fixation of the flaps but a large funnel-shaped ureteral opening with no tendency toward contraction. These cystoscopic examinations were made at intervals of three weeks to three months following operation.

(Thus I believe that I have a method of anastomosis in which the possibilities of stenosis are very small, thereby eliminating the great danger of all forms of ureteral implanation, whether into bowel, skin, bladder or vagina—resulting hydronephrosis being the common sequel of all forms of experimental ureteral grafting.

Fig. 2.—Traction sutures passed through incision and out through bladder wall ready for tying.

How infection takes place following a ureteral grafting I do not think any one knows. The common theory is that it is due to the loss of the valve-like opening of the normal ureter and the substitution of an orifice which is patent, thereby allowing regurgitation of infected urine from the bladder. I believe that regurgitation does not account for these infections, but that infection is due to the stenosis following the method of anastomosis, thus forming obstruction to the free discharge of urine into the bladder.

In support of this assumption I present two cases of ureteral fistula in which the severed end of the ureter was markedly strictured, the ureter dilated and thickened, and a pronounced pyuria present. The strictured ends were cut off, the anastomosis into the bladder completed by the flap method herein described, giving a wide patent orifice and in three weeks' time the pyuria in each case had entirely disappeared.

Granting, however, that this regurgitation does take place after ureteral grafting, I do not believe that there is any valve to a normal ureteral orifice which prevents this happening, for Pozzi, Modkinsky and Desnos report cases of leakage of urine from the bladder through the vesical stump of a divided ureter, Hartman reports leakage following a nephrectomy and I have reason to believe that the same thing happened in one of my cases. Certainly there are thousands of cases of ascending renal infection in which the ureters and bladder are anatomically normal. Polk did a successful bladder implantation of ureter following resection for cancer. A second operation for return of cancer revealed a dilated and thickened ureter which he considered due to constriction at the point of anastomosis.

I believe that infection of the ureter and renal pelvis of the affected side is always present in ureteral fistula and since this condition cleared up rapidly in my cases after anastomosis by the flap method, I am led to believe that regurgitation plays a small rôle, if any, in ascending infections following ureterovesical grafting. If I believed otherwise, then the method of anastomosis herein described should be condemned, for with the resulting wide opening ample chance for regurgitation is offered.

Free drainage is the great surgical principle in the relief of all infections and no ureter or kidney already infected can be rid of this condition while stenosis exists at the ureteral orifice.

Fig. 3.—Cross section of anastomosis complete; (a) traction suture tied; (c) vesico-ureteral sutures tied.

While the urine is being poured into the bladder the organ is in a quiescent state and it empties itself by a contraction of its musculature, which is composed of an inner and outer longitudinal and a middle circular layer. Now if the site of ureteral implantation be made at the summit of the bladder to further prevent regurgitation, and the ureter passed directly through the bladder wall, hence through this mesh-like arrangement of muscle fibers, it seems reasonable to conceive that vesical contraction would produce a valve-like constriction of the ureter, thereby preventing any forcible regurgitation. This procedure was followed in my cases.

Several operators have endeavored to imitate the normal oblique course of the ureter by folding the bladder around the ureter. This was first done by Witzel, after the manner of folding a handkerchief around a pencil (Fig. 5). If this method is used it can be readily seen how, when vesical contraction takes place,

instead of contracting with a valve-like action the walls of the ureter are held apart and regurgitation favored.

Some operators, originally Witzel, claim for this last procedure the advantage that it further anchors the ureter to the bladder and hence prevents any tendency for sutures to tear out at the site of anastomosis. This seems very commendable, but I believe that the two flaps produced by the method described herein, when united to the bladder, result in a sufficient fixation of the ureter to bladder to afford strong union.

If there is any doubt in the mind of the operator regarding the union of these flaps he can readily resect a little of the mucous membrane on either side of the vesical incision, thus securing approximation of connective tissue to connective tissue. The use of the traction sutures further facilitates the anastomosis by affording a quick simple adjustment and firm fixation of the ureter while the anastomosis is completed with the vesico-ureteral sutures. Kelly attained this end by grasping the end of the ureter with long forceps passed through the urethra into bladder. Markoe used a small catheter passed up the ureter, across the bladder and out the urethra, while Kaysar held his ureter in place by long sutures passed through the end of the ureter and drawn out through urethra, fixing them to a weight or the external dressing.

Fig. 4.—Cystoscopic view of anastomosis healed.

As regards the method of approach to the ureter I think that the extraperitoneal route incurs not only considerably less risk to the patient but affords a freer access to the ureter and plenty of space through which to work. Baumm was the first to do an extraperitoneal operation, while Tauffer, Markoe, Bazy and Navarro employed the transperitoneal route. Albarran made his approach by the hypogastric route while Delbet used the L-shaped incision over the sacrum advocated by Kraske.

In reviewing the literature I have been able to collect seventy-seven cases of ureterovesical implantation with thirteen deaths, giving a mortality of 16.08 per cent. Of these seventy-seven cases, thirty-eight were intra-peritoneal operations with eight deaths, a mortality of 21 per cent. Eleven cases were extraperitoneal with one death, a mortality of 9.09 per cent., while in the remaining twenty-eight cases there were four deaths and the method of approach undetermined. Thus statistics point with greater favor to the extraperitoneal route.

I make the lumbo-ilio-inguinal incision beginning in the loin one-half inch below the last rib and at the outer margin of the erector spina muscle. This runs obliquely downward parallel with the last rib to a point one inch inside the superior iliac spine and is then prolonged downward parallel with Poupart's ligament by cutting with scissors while the peritoneum is separated with the forefinger.

Although this is a long incision, the important structures cut are very few. Posteriorly the subcostal artery and vein and last dorsal nerve should escape, and cutting forward the only other structures severed besides the muscles are the lateral cutaneous branch of the last dorsal nerve and the superficial and deep circumflex iliac arteries. The hypogastric branch of the ilio-hypogastric nerve which runs parallel with the wound should escape cutting.

The shock following the extraperitoneal operation is

Fig. 5.—Witzel's method.

small compared to that of any intra-abdominal work and the fact that by the intra-abdominal route the peritoneum is cut both in front and behind, that the sympathetic nerve supply is vastly disturbed, combined with the imminent possibility of infection, should always make the extraperitoneal operation the one of election. In conclusion I report a second case in which the patient was operated on successfully by the method described.

CASE 2.—Patient.—Miss S., aged 26, white, was referred to me by my father. She had had a panhysterectomy done on her five months previous, and five days following the operation urine began to flow from the vagina. I found the left ureter discharging into the left vaginal fornix and a vagina so small that it was impossible to do any sort of plastic operation through this channel.

Operation.—The lumbo-ilio-inguinal incision was made, and the abdominal flap together with the peritoneum was turned inward. The ureter was located at the crossing of the iliac artery, and since hysterectomy had previously been done I had no difficulty in freeing the duct down to the vagina. It had

been tied about the cervix uteri and hence I had a full-length ureter to transplant into the bladder. The orifice of the ureter was markedly strictured and the whole tube thickened and dilated. The ureter was split upward about one and one-half centimeters, making two lateral flaps, two chromic gut sutures were passed and the bladder incised at the point of selection. Boldt advises distention of the bladder with water to aid in selecting the point of anastomosis, but I think that this contributes towards the possibility of infection from leakage through the incision. In my cases the bladder was emptied and then cut on the point of a male sound, while an assistant made the bladder pouch at the desired point. The incision was about one and one-half centimeters long, and the edges at once grasped with forceps and held up. The needles carrying the traction sutures were then passed down through the incision and made to pierce the bladder from within out and at a point distant from the margin of the incision equal to the length of the ureteral flaps. These traction sutures were then tied and the anastomosis completed by four vesico-ureteral sutures. The wound was closed in layers and healed by first intention.

Postoperative History.—Before the operation the patient was voiding daily from her bladder about sixteen ounces of urine, and the first twenty-four hours after operation she passed thirty-four ounces. On the fourteenth day the patient was out of bed and the pyuria which previously existed had entirely cleared up at the end of three weeks. Five and six months, respectively, have elapsed since these operations, and the results up to the present time have been perfect.

Clinical Notes

ETHMOIDAL PERFORATION OF THE INNER CORNER OF THE EYE AND INVOLVEMENT OF THE ANTRUM OF HIGHMORE.*

CULLEN F. WELTY, M.D.
SAN FRANCISCO.

Patient.—C., male, aged 70, single, had always been healthy.

History.—Two weeks before consulting me he noticed a swelling over the right eye, which increased in size daily (Fig. 1). The pain and temperature also increased from day to day. Temperature was present, 104. He had never noticed a discharge from his nose. He has good sense of smell and no odor in the nose.

Examination of the Nose.—Right side, large roomy nose. Middle turbinate, perfectly healthy; middle meatus, roomy and free from pus. Olfactory fissure was free from pus. Probe puncture of the antrum of Highmore brought a large quantity of offensive and cheesy pus.

Treatment.—In order to cleanse the pus cavity, the incision had to be carried as far back as the ear. From the center of this incision another one was made at right angles extending almost to the vertex. A fistula leading to the upper, and another to the lower lid was opened at the dependent point, and gauze introduced through the fistula. Very careful search was made for the bony fistula, but it could not be found. Further operative procedure was postponed until this communication was found.

Subsequent History.—In three weeks the various incisions had healed with the exception of a small sinus that continued to discharge just above the outer angle of the eye. A canula was introduced and the wound irrigated with sterile water which came through the nose. On further search for the bony fistula it was found to be posteriorly, and above, and on the inner side of the eye. I wish to call attention to this bony fistula occurring so far back, and again to the fact that the fistula discharged at the outer corner of the eye, rather than

the inner. I am not familiar with a similar case on record. This fistula was irrigated for about four weeks, but the treatment failed to bring about a cure.

Second Operation.—The Killian operation was recommended and done. The frontal sinus was full of polypi and pus. The anterior and postethmoid cells were a mass of polypoid tissue. Sphenoid cavity was full of polypi.

Result.—Convalescence was rapid. However, some pus continued to come from the sphenoid cavity and a second curettage was necessary which effectually cured the discharge. A short time after the radical operation was done on the antrum of Highmore. The outer incision closed by interrupted sutures

Fig. 1.—Showing patient before operation.

Fig. 2.—Showing patient after operation.

(Fig. 2). The antrum of Highmore was completely filled with polypi, granulation tissue and cheesy pus, which was very offensive. There had been such extensive destruction of bone that the antrum extended to the median line of the hard palate. The bone that remained of the hard palate was paper thin.

I wish to call especial attention to this case as I do not find a similar one on record in Zuckerkandl's work on "Anatomy and Pathology of the Accessory Sinuses of the Nose." Neither does Hajek.

The floor of the nose was also paper thin; was removed and at present the floor of the nose is the paper-thin bone of the under surface of the hard palate.

* Read in the Section on Laryngology and Otology of the American Medical Association, at the Fifty-ninth Annual Session, held at Chicago, June, 1908.