ARTERIAL HYPERTENSION*

FREDERICK M. ALLEN, M.D.

NEW YORK

The great importance of arterial hypertension or the conditions associated with it, as a widespread and apparently increasing cause of disability and death, is a self-evident incentive to investigation. At the same time it must be recognized that our knowledge of the subject is in a very confused state, as respects its phases of etiology, mechanism and treatment. Just as with diabetes, the literature contains almost every conceivable variety of opinions and suggestions. The first task of an investigator to choose is what seems significant and valuable among this mass of conflicting material, and the second is to build concrete facts with accurate methods in a clinical field which is still largely a morass of tradition and imagination.

THEORIES REGARDING THE CAUSES OF
HYPERTENSION

More or less restriction of salt has been employed for many years by some observant clinicians in the treatment of the various forms of nephritis. The first scientific foundation of both theoretical and practical importance is afforded by the work of the French school, beginning with Widal's demonstration of the rôle of the chlorids, particularly in the causation and treatment of edema, and with the application of this principle to hypertension by Ambard,¹ Combe,² Laufer³ and others. Similar results were obtained in arteriosclerosis by Bayer;⁴ but with only a few exceptions the tests with nephritis in Krehl's clinic were negative or doubtful, and the general German experience was opposed to a strict relationship between retention of chlorids and elevation of blood pressure.⁵

In American literature, salt and fluid restriction in hypertension is mentioned casually and incompletely if at all, and the protein intoxication theory is undoubtedly the dominant one. Treatment consists chiefly in low protein diets, the elimination of supposed toxins, or the artificial reduction of pressure by drugs, bleeding, electricity, and the like. Mental and bodily rest is advised to a degree which largely terminates usefulness in life, and many conservative practitioners refrain from any serious attempt to reduce the pressure, and devote themselves to keeping the patient as comfortable as possible, with resignation as to the results.

There are various reasons for this confusion and pessimism. In the German work the cases were not always well chosen and the restriction of salt was generally inadequate, though this work does furnish evidence against any invariable association of hypertension and gross chlorid retention. On the other hand, whether or not protein has any influence on blood pressure, it is impossible to accept Ambard's assertion that it is harmless in nephritis. Ambard considers fluid restriction unnecessary, and in adopting milk diet he comes back to the identical therapeutic program used

by the followers of the intoxication hypothesis. The fluid restriction of Laufer, who accepts von Noorden's limit of a liter daily without reckoning the important additional quantities contained in the food, is too lax for the worst cases. Likewise a much stricter exclusion of salt than represented by milk diet is essential for results in really severe cases. These inaccuracies are responsible for an unnecessarily high percentage of cases of absolutely or relatively irreducible hypertension in French, German, American and all other records, and also for the unnecessarily frequent and early deaths from apoplexy, cardiac failure, pulmonary edema and convulsions (misnamed uremic). For the same reason there have been no differences in therapeutic results sufficiently marked to establish any one theory to the satisfaction of all parties.

RESTRICTION OF WATER AND SALT

It would be wrong to ignore the unknown and uncontrollable factors in hypertension and the unsolved problems which they present. But any practitioner who will conform his treatment to one reasonable and definite condition, namely, the necessity of the organism to force a filtrate of water and dissolved substances through a damaged and partially blocked glomerular filter, can readily demonstrate for himself the compensatory element in the hypertension, by observing the fall in pressure and the relief of some attendant symptoms when proper diet reduces the quantity of this filtrate to a minimum. The opposite treatment under the intoxication hypothesis, through unsuitable diets and attempts at "flushing out," entails much harm. In some hypertension cases the mere reduction of the overload of salt and water brings relief which is comparable to the benefits of diet in diabetes.

Fuller laboratory study, particularly by blood analyses, contributes new information of theoretical, prognostic and therapeutic importance. Among the nephritic cases treated in our clinic to date, there have been twenty examples of marked hypertension. Most of these were treated in private hospital status, but four were outpatients. The observations are necessarily empiric, and are subject to modification from wider clinical experience, and particularly to more fundamental and decisive interpretations when animal experiments become available. Numerous data of these cases, function tests according to Ambard's principles, and details of individual histories and results, must be reserved for a more complete publication at a later date.

It may suffice here to state that the patients' ages ranged from 25 to 69 years, with the usual predominance of those past 40. All the cases were of chronic type. Their severity was mostly too great to yield to mere rest and hospital care, and some were known as highly severe cases which had proved intractable to a variety of customary treatments for several years. Bleeding was used temporarily in three critical cases, and brief experiments have been tried with some other measures; but in general the patients were not kept in bed or confined to the house but were treated entirely by diet.

To establish the general principle, the effects of sudden dosage with salt or water in suitable cases are illustrated by two experiments.

In the first experiment (Table 1), attention may be called to the absence of real retention of either salt or water (contrary to the French reports), and to the

* Read before the American Society for Experimental Pathology, Cincinnati, Dec. 30, 1919.

* Preliminary communication, based on a clinical study in collaboration with Drs. J. W. Mitchell and J. W. Sherrill.

1. Ambard, L.: Thèse de Paris, 1905; *Semaine méd.* 26: 361, 1906.

2. Combe, A.: *Monatsschr. f. Kinderh.* 4: 13, 82, 1905.

3. Laufer, R. J.: *Compt. rend. Soc. de biol.* 56: 249, 1904.

4. Bayer, R.: *Arch. f. exper. Path. u. Pharmakol.* 57: 162, 1907.

5. Loeb, A.: *Deutsch. Arch. f. klin. Med.* 85: 348, 1905-1906.

Brodski, J.: *Berl. klin. Wchnschr.* 43: 906, 1906; *Deutsch. Arch. f. klin. Med.* 93: 310, 1908. Löwenstein, C.: *Arch. f. exper. Path. u. Pharmakol.* 57: 137, 1907.

remarkably high plasma chlorid figures⁶ and the elevation of blood pressure accompanying these.

The second experiment (Table 2) was performed on a different patient. The plasma chlorid percentage was not high, and fell lower with the water drinking, while the blood pressure rose markedly.

The first patient received salt, but showed no effect from a corresponding test with water alone. Because of the close relations between salt and water, it is

TABLE 1.—SALT TOLERANCE TEST, SEPT. 25, 1919

Time	Water Given, C.c.	Urine			Plasma Chlorids		Blood Urea		Blood Pressure	
		Vol., C.c.	Sp. Gr.	NaCl %	NaCl,* Gm.	Mg. per 100 C.c.	Mg. per 100 C.c.	Sys- tolic	Dias- tolic	
8 a.m.	200	...			...	...		202	104	
9 a.m.	200	65	1.011	0.72	0.47	...		...	...	
10 a.m.	200	410	1.006	0.30	1.21	...		...	...	
11 a.m.*	200	305	1.007	0.34	1.03	672	19.6	200	100	
12 m.	200	120	1.008	0.53	0.53	672	19.6	198	94	
1 p.m.	200	160	1.015	1.15	1.84	684	19.6	194	98	
2 p.m.	200	105	1.016	1.21	1.27	676	...	224	100	
3 p.m.	200	120	1.016	1.30	1.55	704	...	242	110	
4 p.m.	200	210	1.007	0.90	1.88	...	...	266	110	
5 p.m.	200	500	1.007	0.44	2.20	...	...	238	100	
6 p.m.	200	250		0.60	1.49	648		210	100	
6 p.m. to 8 a.m.	550	830	1.009	0.41	3.40	644	19.6	182	80	
Total	2,750	3,275			16.87					

* Ten grams of sodium chlorid were given at 11 a. m.

difficult to distinguish accurately between their effects, and space requirements necessitate leaving other tests to a later publication. The rise in pressure in both cases was accompanied as usual by the subjective symptoms of headache, dizziness, weakness and general malaise. Precautions were taken so that no patients suffered harm from such tests, but in severe cases there is danger that the reckless administration of either salt or water may cause alarming symptoms and even death. The accuracy of the tests is often interfered with by a protective diarrhea which also prevents dangerous disturbances.

The results of a short test may be negative in some patients or at some stages, and bad effects may become

TABLE 2.—WATER TEST, NOV. 19, 1919

Time	Water Given, C.c.	Plasma Chlorids		Blood Urea		Blood Pressure	
		Mg. per 100 C.c.	Mg. per 100 C.c.	Mg. per 100 C.c.	Mg. per 100 C.c.	Sys- tolic	Dias- tolic
8:30 a.m.	...	571	30.2	...	...	215	130
8:40 to 9:20 a.m.	1,000	...	...	...	...	...	...
9:25 a.m.	...	...	...	...	...	226	128
9:20 to 9:50 a.m.	600	...	...	...	...	...	...
9:55 a.m.	...	...	...	...	...	245	130
10:00 to 11:10 a.m.	1,400	...	...	...	...	...	...
11:15 a.m.	...	...	...	...	...	238	140
11:20 a.m. to 12:20 p.m.	800	...	...	...	...	...	...
12:30 p.m.	...	543	23.7	...	...	254	120
2:25 p.m.	...	...	...	...	...	226	125
4:00 p.m.	...	...	...	...	...	188	100
5:00 p.m.	...	...	...	...	...	190	100
9:30 a.m.	...	...	...	...	...	200	120

evident only after some days. The effects of salt are in general more powerful and obvious, as demonstrated by the success of Ambard and others in numerous cases. A very close restriction of water may be impracticable at the beginning of treatment, because of the salt stored in the patient's body and the inadvisability of imposing any real suffering from thirst. Also in many cases no rigid fluid restriction is necessary, and moderate quantities (from 1 to 3 liters daily) may be actually beneficial for flushing out salt or nitrogen. But the prolonged combination of excessive fluid and

little salt possibly involves osmotic and eliminative difficulties which are responsible for the anasarca, pulmonary edema and cardiac failure sometimes described in connection with long continued salt-poor diet.

No rigid classification, either anatomic or functional, of the cases listed in Table 3 was attempted, partly because the experience is insufficient, and partly because every case of nephritis is in some respects an individual problem, in which tests of the efficiency of different functions should be made and individual treatment prescribed accordingly. Different tests will give different classifications, and there are presumably types of cases not shown in this series. The phenol-sulphonphthalein excretion is given here as one indication of renal action, but the complete publication will include other function tests. The data shown are otherwise limited to those which seemed most essential for treatment and prognosis. The order of cases is not merely consecutive or accidental, but forms a series based on these data.

TABLE 3.—EFFECTS OF DIETETIC TREATMENT IN TWENTY CASES OF HYPERTENSION

Case No.	Phenolsulphonphthalein Excretion in 2 Hrs., %	Before Treatment				After Treatment			
		Plasma Chlorids, Mg. per 100 C.c.	Blood Urea, Mg. per 100 C.c.	Blood Pressure, (Pressure*) Sys- tolic	Dias- tolic	Plasma Chlorids, Mg. per 100 C.c.	Blood Urea, Mg. per 100 C.c.	Blood Pressure, (Pressure*) Sys- tolic	Dias- tolic
1	...	624	29.2	208	...	569	28.0	166	98
2	54.5	640	27.1	(300)	...	...	...	...	...
3	54.0	620	21.8	240	170	588	20.0	134	70
4	42.8	610	19.8	210	110	541	24.5	140	80
5	36.0	620	20.5	240	110	586	21.2	160	95
6	...	670	23.5	220	140	590	10.4	166	75
7	46.0	608	32.2	195	92	556	23.5	178	98
8	48.0	613	30.7	220	98	591	26.9	162	80
9	...	611	32.6	220	70	570	28.5	110	80
10	41.3	612	47.0	220	110	576	16.2	130	66
11	4.0	556	69.4	212	104	555	31.4	154	80
	(625)	...	...	...	...	556	52.6	156	80
12	30.2	664	94.4	210	120	596	26.4	172	100
13	...	659	104.5	198	100	545	62.5	90	60
14	5.0	648	134.0	214	170	537	42.2	140	100
15	...	597	27.1	190	92	597	...	108	70
16	38.2	569	36.4	190	90	595	14.6	165	100
17	...	534	16.7	210	100	553	56.2	180	95
18	27.4	576	55.0	235	145	543	23.7	170	98
19	12.1	548	58.2	(305)	...	...	...	...	...
	...	...	...	215	170	505	18.0	180	130
20	64.8	565	21.0	234	160	559	19.4	180	120

* Parentheses in the systolic pressure column indicate that the pressure in these cases was repeatedly near 300 m.m., but the actual pressure on beginning dietetic treatment was 215. In Case 19 this reduction had been accomplished only by heavy bleeding in another hospital.

CASES OF PURE HYPERTENSION

The first cases are more or less typical examples of so-called pure hypertension. One of them (Case 2) had been observed for over a year, and occasional urinalyses, even when the blood pressure was as high as 300 m.m., never showed albumin. In recent months, with more frequent urinalyses, traces of albumin have sometimes been found. The others of this group have traces of albuminuria. Omitting individual peculiarities, it may be said that this group is characterized by activity of excretion and apparent absence of retention in tests with nitrogen, water and even salt. The kidney retains power to concentrate, and both the blood urea and the urea function according to the Ambard principle may be fully normal. But the chlorid threshold, whether determined mathematically or by salt privation, is high, and these patients purchase their apparent efficiency of salt excretion at the price of abnormally high blood salt and blood pressure. Extraordinary strictness of salt-free diet may be necessary to control these, and the therapeutic success corresponds to the restriction in salt intake. This single feature some-

6. For chlorid method, see Donleavy, J. J.: J. Biol. Chem. 37: 551 (April), 1919.

times makes the difference between health and invalidism. While protein and water create no symptoms, on general principles they should be limited to some extent. Without assuming that all possible kinds of cases are represented in this list, and without overlooking other abnormalities revealed by sufficiently thorough study, it can be stated that pure hypertension, so far as it has occurred in this series, has been essentially a salt nephritis.

HYPERTENSION WITH KIDNEY IMPAIRMENT

As all standards of kidney function are to some extent arbitrary, this group passes by gradual transition into the second and larger group, in which there is obvious impairment of several kidney functions, as indicated by retention of urea and other products, impaired excretion of urea and chlorids by Ambard's laws, more or less fixation in the chlorids and specific gravity of the urine, and especially chlorid retention with or without evident edema.

The degree of impairment in the nitrogen economy calls for corresponding restriction of protein, and influences the prognosis. But the hypertension is affected only through the salt and water functions. Careful fluid restriction is necessary as a rule, but attention should be given to the nitrogen retention thus entailed in certain toxic cases. This is an additional reason for limiting the protein intake, though it is well known that the percentage of blood urea is not an infallible index of danger. Even though chlorid retention be far greater than in the first group and the other signs more evident as stated, the actual chlorid threshold may be lower and the actual salt restriction required to control the pressure may be less stringent in the second group.

It is well known that retained salt may be diffused through edema fluid without raising the percentage in the blood. Patient 11 thus had much edema, and as this diminished, the chlorid percentage in the blood rose as shown in parentheses. With this proviso, the great majority of these cases had high chlorids in the blood plasma, and it is probable that the same will hold for the majority of all hypertension cases. But in approaching Group 3, we find a small group of hypertension patients with continuously low blood chlorids and low chlorid threshold, and often with normal or actually low blood urea.

HYPERTENSION WITH DIABETES OR SEVERE NEPHRITIS

The six closing cases of this table may be divided into two groups of three each. The first group includes three cases (15, 16 and 17) of diabetes. In them the prognosis is good with thorough treatment of the diabetes, sometimes with the aid of salt and water restriction and sometimes without it. Though there are presumably rare cases of intractable hypertensive nephritis associated with diabetes, there is generally the strong probability that proper treatment, associated with reduction of the blood sugar, will reduce the hypertension.

The last three cases of the table are nondiabetic, and have proved the most difficult and intractable of the entire series. Limitation of salt, protein and fluid, reducing the twenty-four hour urine volume below 300 c.c., has given symptomatic relief and more or less complete checking of retinitis, but the influence on the pressure has on the whole been slight and the reduction shown in the table has not been permanent. Notwith-

standing the innocent looking blood analyses, according to function tests these are cases of severe nephritis, and the worst element in their prognosis from the standpoint of hypertension seems to be the absence of a high blood chlorid before treatment.

LIMITATIONS IN TREATMENT

Certain precautions or limitations should be mentioned with regard to the practical application of these principles in treatment.

First, a diet which is reasonably satisfying and at the same time sufficiently poor in salt is not so easy to arrange as it may appear. Both physician and patient may sometimes believe conscientiously that strict abstinence from salt has been maintained, when analyses will show the blood chlorids undiminished and an output of from 5 to 10 gm. in the twenty-four hour urine. Such analyses are important for controlling the treatment, and no pressure is rightly called irreducible unless the daily chlorid excretion has been reduced to practically zero.

Second, some quantity of salt is considered indispensable in the permanent ration. This is stated as about 2 gm. for normal persons, but the limits for nephritics have never been determined. On the one hand, it is not certain that the intake need be as great when the output is obstructed. On the other hand, there may be danger in withdrawing salt from certain nephritics, especially with nitrogen retention, who seem to need much more than 2 gm. of salt for its diuretic action to ward off uremia. As a rule, the outstanding effect of undue salt privation has been weakness. It is necessary, therefore, to work out salt rations for individual nephritics which may range from almost zero to a considerable quantity.

Third, restriction of salt or fluid, like every dietetic treatment of organic disease, is only palliative. Success is not as uniform as in the treatment of diabetes, and complete or partial failures are more numerous. When the first results are more or less satisfactory, there is still reason to fear that the same patients may pass into a hopeless condition in the course of months or years. Furthermore, the most effectual relief of the hypertension demonstrates strongly that the essential trouble is more than mechanical. Reduction of the blood pressure nearly or completely to normal may still leave the patient an invalid, either from weakness due to the strict salt privation which is necessary to hold down the pressure, or still worse, if salt is given so as to allow the pressure to go up. They may also be subject to attacks of hypertension and accompanying symptoms from little or no evident cause. The greatest defect of the French work is the disposition of those authors to treat the entire question as a mere mechanical matter of salt retention. Obviously, even the mechanics of hypertension remains largely unsolved, and the fundamental condition thus far baffles both research and treatment.

Notwithstanding these limitations, the benefit of suitable chlorid restriction for hypertension is still great, in making the patients more comfortable, diminishing the danger of apoplexy, and possibly checking the progressiveness of the disorder. So many of these patients give a history of copious use of salt that this may be a possible contributory factor. Granting a primary infectious or toxic etiology of the renal or vascular damage, the prolonged irritation by the excess of salt and the wear and tear of the hypertension itself

may well be regarded as a cause of progressive aggravation. By far the best and easiest results are obtainable in the earliest cases, when the hypertension is intermittent rather than continuous. Early diagnosis is therefore important, and in advance of the sphygmomanometer it may be hoped that chemical tests, such as for the blood chlorids and chlorid threshold (the normal values for which are still undecided), may give advance warnings to guide prophylaxis.

Such studies may also throw some light on the origin and the control of arteriosclerosis and some other disorders. One of the most interesting branches of the investigation has concerned ophthalmic disorders, and with the cooperation of several oculists, evidence has been gathered indicating that many cases of retinitis are due chiefly or solely to chlorids and are benefited by chlorid restriction. More cases of retinitis and of glaucoma are needed for this study. The promising study of salt metabolism will probably soon be extended more widely by those who are in position to undertake parallel determinations of various salts and to develop the more fundamental problems through animal experiments. The present investigation, which has had to be self-supporting to date, has been limited and superficial for lack of these things. The clinical facts collected, however, warrant urging that unfounded theories be discarded, the widespread abuse of drugs, particularly nitrites and iodids, stopped, and the possibilities of modern laboratory methods utilized for the guidance of treatment.

5 East Fifty-First Street.

INFECTION BY THE BROAD TAPEWORM, *DIPHYLLOBOTHRIUM LATUM* *

M. W. LYON, JR., PH.D., M.D.
SOUTH BEND, IND.

While cases of infection by the broad tapeworm have been rather frequently reported in the United States in recent years, and most of them, like the present one, have occurred in immigrants who have acquired the infection abroad, yet because of the probability of the permanent establishment of this parasite in this country, it is desirable to record each case with as many details as possible.

The worm here recorded was casually handed to me in the early part of January of this year. The dark spots in each mature segment had been noticed, but the true nature of the parasite had not been fully appreciated. The specimen had been washed from the feces with which it was expelled by the patient. The head is lacking, although many of the immature segments immediately following the head are present. The specimen is in several pieces, and the combined length of them laid out without traction on a table is 6,230 mm. All the segments appear in excellent condition, and none appear atrophic or degenerated.

The patient presented almost no symptoms.

REPORT OF CASE

B. N., man, aged 39, Russian Jew, machinist, born at Bielostok, Poland, where he resided until he was about 19 years of age, spent the next six years, including the time of the war between Russia and Japan, and while with the Russian army, at various places in Siberia, such as Omsk, Tomsk, Irkutsk and Yakutsk. During these six years he ate much

uncooked fish. Although much fish had been eaten during his youth at Bielostok, he thinks it was well cooked by his mother. After the six years spent in Siberia he returned to Bielostok, and soon after came to the United States. In this country he spent one year in New York, six years in Chicago, and six in South Bend, Ind. He first knew he had a tapeworm eight years before, while he was living in Chicago, his knowledge being based on the fact that he passed fragments from 12 to 18 inches in length. These fragments appeared to him well preserved, and he says that they showed motility. He found segments only at the time of defecation. He never noticed any segments passed while he was in the Old World, but he suggested that, being younger, perhaps he was a careless observer and did not pay so much attention to himself. The patient said that he had always been well and had never consulted a physician until he wished the worm expelled in the latter part of December, 1919, and except when being examined for life insurance. He passed the insurance examination successfully. About three months before seeking treatment for tapeworm he had obscure abdominal distress; and he had another attack at the time he sought treatment. He consulted Dr. C. C. Terry of this city, by whom he was given aspidium. A few days later, when he had recovered from the effects of medication, the worm was submitted to me for identification. I was unable to get the patient for a blood examination until nearly two weeks after the worm had been expelled. Blood examination then revealed: erythrocytes, 5,148,000; leukocytes, 6,400; hemoglobin, 85 per cent., Sahli; polymorphonuclears, 56 per cent.; small lymphocytes, 28 per cent.; large lymphocytes, 7 per cent.; large mononuclears, 2 per cent.; transitionals, 3 per cent.; eosinophils, 3 per cent.; basophils, 1 per cent. The patient has the appearance of being in perfect health, and looks strong and robust.

COMMENT.

There can be little doubt that this patient became infected with a broad tapeworm while living at Bielostok, which is one of the general regions where such worms are common, or in Siberia, where he knows he ate uncooked fish. He may have acquired it in this country, but that is very doubtful.

That the patient is and has always been robust shows that an anemia is not necessarily caused by the presence of this parasite. The fact that he passed well preserved and easily recognized segments is rather unusual¹. The absence of anemia, and the good condition of the segments passed as well as of the entire worm as subsequently passed, are in accord with the views that diseased or disintegrating worms are the chief cause of the anemia due to the absorption of products from disintegrating segments.²

This case has afforded an excellent opportunity for the parasite to become permanently established in this region of the United States. For the past six years many thousands of eggs must have entered the sewerage system of South Bend, very soon to reach the St. Joseph River and a short time later Lake Michigan. In both river and lake, hatching larvae must have been able to parasitize various fishes.

1. Brumpt, E.: Précis de parasitologie, Ed. 2, 1913, p. 249.
2. Brumpt: Précis de parasitologie, p. 263.

Probabilities as to When the First Birth Will Occur.—According to Knibbs, statistician for Australia, the probability of a nuptial first birth occurring in less than one year after marriage is 0.4946 for age 15 of the wife; 0.7770 for age 18; 0.9176 for age 21; the maximum, or 0.9771, for age 25; 0.9075 for age 30; 0.6748 for age 35; 0.3245 for age 40, and 0.0622 for age 45. The foregoing statistics are based on the supposition that the husband is approximately the same age as the wife. If he is younger or older than the wife, the probability will be affected accordingly.

* From the laboratory of the Clinic.