Clinical Notes :

MEDICAL, SURGICAL, OBSTETRICAL, AND THERAPEUTICAL.

CASE OF TUMOUR OF MESENTERY.

By J. K. HAWORTH, M.D., B.S. DURH.,

RESIDENT MEDICAL OFFICER, ROYAL VICTORIA INFIRMARY,
NEWCASTLE-ON-TYNE; LATE CAPTAIN, R.A.M.C., S.R.; LATE
SURGEON-SPECIALIST, EIGHTH LUCKNOW DIVISION.

THIS case, I think, sheds a little more light on the interesting and rare conditions of mesenteric tumour.

While doing duty as surgical specialist to 8th Lucknow Division, Rifleman H. G., 1st Garrison Battalion, Royal Irish Rifles, aged 28 years, was sent to me from Cawnpore on August 12th, 1917, with a lump in his abdomen. He had very few symptoms, but the presence of this tumour seemed to worry him. On examination he was found to have a tumour about the size of a large tangerine orange, situated in the right iliac region, below and to the right of the umbilicus; it felt hard and circumscribed, and it was movable all over the abdomen. He had no difficulty with his bowels, and apart from his tumour he felt fairly well. A diagnosis of mesenteric cyst was made.

Operation.—The abdomen was opened by a vertical incision in the right semilunar line below the umbilicus over the tumour. A mass, yellow in colour, was seen in the mesentery rather nearer to its posterior than anterior attachment. The cyst was in a favourable position with regard to vessels and was enucleated. The mesentery was sutured and abdomen closed; other enlarged glands in the mesentery were felt and the case was considered to be tuberculous in origin. On examination the tumour consisted of a thick firm homogeneous capsule containing semi-solid caseous material and debris, no signs of any hairs or material suggesting a dermoid. It was unilocular. There was no minute pathological examination of the wall or contents. Recovery was uneventful.

My case is almost identical with the cases reported by Mr. E. L. Pearce Gould,¹ except that I was able to make out other glands in the mesentery; this fact, I think, adds weight to Pearce Gould's supposition that these conditions are tuberculous in origin.

THE BACTERIOLOGY OF COLITIS.

METHOD OF INVESTIGATION FROM INTESTINAL MUCUS
INSTEAD OF FÆCES.

By H. L. LYON-SMITH, M.D. DURH.

THE presence of mucus in the stools is the chief point in the diagnosis of colitis, as mucus is not a normal constituent of intestinal evacuations. In taking the history of a case patients are usually found to be ignorant of the presence of mucus, with the exception of more severe cases where they have noticed slimy and abnormal motions. The majority of physicians have been content to order a specimen of the fæces to be sent to the bacteriologist, who usually receives for examination purposes a small portion of a solid stool, from which he is expected to prepare a vaccine of the most conspicuous organisms. It is the method which is required for distinguishing which foods have been digested or otherwise, but is too gross a one for the diagnosis of the bacterial factors in colitis. The malady is not of the fæces, but of the intestinal mucous membrane, and the closer we get to the latter in our researches the more likely we are to discover the real nature of the pathogenic organisms concerned in the ætiology.

The Method Described.

It has been the practice of the writer for the last seven or eight years to ask the patient to have a Plombière douche, and from the second washing flakes of mucus can be collected. The rectum is first washed out with a pint of warm water and the faecal discharge thus obtained discarded. A second irrigation is then made with two or three pints of plain warm water, slowly injected, and allowed to remain in the intestines about six minutes, whilst the patient turns from side to side in order to separate the adhering mucus. This second wash-out is received in a clean receptacle, and search is then made for flakes of mucus, which are transferred to a sterile specimen bottle. When received these flakes are poured out into a large Petri dish half full of sterile normal saline and examined against a black background in order to obtain some

idea of the character of the mucus. Under macroscopic examination these may be found in thin, pale brown, soft flakes, or, in other cases, thick, dark brown, tenacious masses, and in severe cases a complete cast of the intestine may be found. To search for amœbæ, a portion of the membrane may be broken off and examined under a cover-glass on a warm slide. Other fresh films are made and stained.

Slide 1.—Stained by carbol-thionin, which shows up, clearly, capsulated bacteria, such as enterococci, *B. mucosus capsulatus*, pneumococci, or the longer chains of *Streptococcus mucosus capsulatus*. It will also show if there are any ova of intestinal parasites.

Slide 2.—Stained for T.B.

Slide 3.—Gram-stained, gives a differential estimation of the relative proportions of the Gram-negative and Gram-positive bacteria. In most specimens of mucus from cases of colitis the latter exceed the former in numbers.

For cultural purposes a small piece of the mucus is teased off with platinum needles, put into a tube of sterile broth, incubated at 37° for an hour, then well shaken up, and loops of this broth are subcultured on various media at the discretion of the worker. The writer prefers to use: (1) ordinary agar tubes; (2) glycerine agar; (3) freshly made blood-agar tubes; (4) bile-salt-agar plates, a very convenient preparation for making these being the "Solmedia" preparation of MacConkey's bile-salt agar, issued by Baird and Tatlock.

The freshly made blood-agar media enable one to pick out those small, powerfully hæmolyzing colonies of streptococci which are extremely pathogenic and frequently the chief causes of dangerous anæmias. In one case of pernicious anæmia accompanied by colitis, a vaccine of hæmolytic streptococci was the principal constituent of an antigen which proved most effective, as shown by the differential blood counts. This occurred three years ago, and the patient is still in good health.

The advantages of this method of examination over that in general use—namely, investigation of the solid fæces—must be evident, as it is possible for the bulk of the fæces to be evacuated without having ever been in contact with the inflamed intestinal walls, so that the pathogenic organisms responsible for the colitis may easily escape detection.

A CASE OF FATAL INTERNAL LACERATIONS WITHOUT EXTERNAL WOUNDS.

By WALTER E. MASTERS, M.D. BRUX., M.R.C.S.,
L.R.C.P.

THE case here described is probably not unique, but it is significant that there should have been such severe internal injuries without as much as a bruise about the abdomen or chest; it is also remarkable that the boy should have lived for one and a half hours with the splenic vessels torn off.

On April 19th, 1920, a healthy young adult African native fell down a mine-shaft for a distance of 150 ft. to the bottom. He was brought to the Prestea Hospital and seen by me immediately, about half an hour after the injury. He had a laceration of the dorsum of the left foot, minor bruises of the legs, arms, and head of an insignificant nature. Blood was coming from his penis, due to a rupture of the urethra near the prostate, resulting from a blow on the scrotum during the fall. The bladder was intact. The foot wound was being dressed when the writer was suddenly called away. The patient had said that he was not in much pain, and turned himself over while being dressed. The writer returned in three-quarters of an hour and the boy was dead, having lived for one and a half hours after the injury.

A post-mortem examination was carried out. There were no abrasions or bruises of any kind on the trunk, externally or internally. On opening the abdomen three pints of red blood escaped. The spleen, four times its normal size, was badly lacerated near the hilum, the splenic vessels were completely torn off, but the spleen was not separated from its normal bed. The cardiac end of the stomach was lacerated for four inches, and the stomach contents were free in the abdomen. There was hæmorrhage into the pancreas and great omentum. Death was due, of course, to internal hæmorrhage.

The rarity of the circumstances and the vitality of the subject seem to me to justify publication of the case.

Prestea, Gold Coast.

ERRATA.—In the article by Drs. Dreyer and Burrell on the Vital Capacity Constants that appeared in our last issue some corrections arrived after we had gone to press. On p. 1213, near the middle of the first column, "No. 11 (see Table IIA)" should read "No. 11 (see Table IB)"; and the lines commencing a few lines lower should read: (1) No. 15, (2) No. 55, (3) No. 139, (4) No. 157, (5) No. 168, (6) No. 176, (7) No. 178, (8) No. 196 (see tables), (9) No. 115 (returned — 9.4 per cent., early good case which did well), (10) No. 164 (returned — 4.2 per cent., a patient with indefinite signs and good condition, but in whom T.B. were found).

¹ Brit. Jour. Surg., vol. iii., No. 9.