PRE-PALEOZOIC DECAY OF CRYSTALLINE ROCKS NORTH
OF LAKE HURON

BY ROBERT BELL

ASSISTANT DIRECTOR OF THE GEOLOGICAL SURVEY OF CANADA

(Read before the Society December 28, 1893)

CONTENTS

	Page
Relation of Archean to Paleozoic Rocks.....	357
Evidence of pre-Paleozoic Decay.....	358
Where best displayed	358
"Ovens" or Pits of Decay.....	358
Rocks of the Huronian Belt	359
Ancient Erosion	359
Typical Erosion Surface	359
Conditions affecting Erosion.....	361
Contacts of Archean with Paleozoic Rocks.....	362
River and Lake Channels due to Rock Decay.....	364
Absence of Paleozoic Deposition and its Significance	366
Effect of glacial Denudation.....	366

RELATION OF ARCHEAN TO PALEOZOIC ROCKS.

In all parts of Canada where the Paleozoic strata come in contact with the underlying Archean rocks the latter appear to pass beneath them with very much the same contour or slope as that of their modern surface. It appears evident that these ancient stratified or laminated rocks have been tilted to their present high angles, and that their included intrusive masses, whether of granite or greenstone, and even most of their dikes, have occupied their present positions relatively to each other before the commencement of Paleozoic time, excepting in cases where faulting has taken place. Not only is this so, but at that early date they appear to have been worn down by denudation to something like their present superficial aspect.

What has become of the vast amount of *débris* resulting from this denudation? This is a problem of American geology which has not

yet received much explanation. Between the time of the folding of the Huronian rocks and the deposition of the earliest fossiliferous beds there must have been an interval much greater than is commonly supposed. Throughout the vast region of Canada occupied by the Archean rocks the attitude of the Huronian and Laurentian strata generally approaches the vertical, and their surface has generally been cut down to nearly a horizontal outline, which is only slightly raised above the sealevel.

Lower Silurian strata, especially limestone, unaltered and full of fossils, may be seen in many places, resting almost horizontally upon the up-turned and denuded edges of the crystalline rocks, and yet the former contain but few fragments of the older terranes, and they occur only at the very contact, showing that the latter had a hard and naked surface when the Paleozoic rocks were being deposited upon them.

EVIDENCE OF PRE-PALEOZOIC DECAY.

Where best displayed.—The evidence of pre-Paleozoic decay of the ancient surface is more discernible upon the surfaces of masses of intrusive granite than upon gneiss or other foliated rocks. The reason of this seems to be that in the bottom of the deep sea, where the eating away of the rocks appears to have taken place, the granite was more susceptible to disintegration than the others. Possibly this susceptibility may have been increased by a somewhat higher temperature in the intrusive granite.

"Ovens" or Pits of Decay.—The northwestern portion of George island and the greater part of the township of Rutherford, at the northwest extremity of Georgian bay, consist of red granite. Wherever the surface of this rock has been protected from glacial action it is found to be eaten into hollows in the form of round and sack-like pits and small caverns. The latter generally occur on steep slopes or in perpendicular faces of the rock, which, however, do not rise to any great height in this vicinity. The floors of these caverns are usually flat, as they are formed by the lower sides of horizontal joints, while the roofs are arched like ovens. Whatever may have been the agency which excavated these ovens, it always worked inward from the granite face and upward from the floor formed by a joint, the rock of the latter not being affected, but remaining as sound here as elsewhere along the joint-plane.

One of these caverns, which occurs in the western part of Killarney village, opposite to George island above mentioned, and which is shown in figure 1 of plate 15, the reproduction of a photograph, has almost the form and dimensions of the clay ovens used by the French Canadians for baking bread, while other caverns in the neighborhood bear more or less resemblance to this one; hence they have received the local name of

FIGURE 1.—“OVEN” IN GRANITE, WEST END KILLARNEY VILLAGE.

FIGURE 2.—ERODED GRANITE SURFACE AT KILLARNEY VILLAGE.

PRE-PALEOZOIC ROCK DECAY.

"ovens," which may also be adopted for convenience as their geologic designation. The oven shown in this figure measures 5 feet deep, 8 feet wide and $2\frac{1}{2}$ feet high. Two men can crouch inside of it. The door or entrance is 18 inches high and $2\frac{1}{2}$ feet wide, which are also about the proportions of the door of a baking oven of the above dimensions.

The pits, which occur mostly upon the nearly horizontal surface of the granite, differ from glacial kettle-holes in being shallower, seldom quite circular and in having rough walls. The sack-shaped ovens are most numerous upon sloping surfaces. In nearly every case the fundus of these hollows is pointed up-hill, arched over by a roof having a thin edge of rather harder granite than the general mass, and is in the form of a segment of a circle. The granite around these pits and caverns is all bare and exposed to the weather, but it shows no decay or disintegration below the immediate surface. Figure 2 of plate 15, the reproduction of a photograph taken at Killarney village, represents this condition. In some places near by it is partially smoothed and striated, as if the glacier had touched it very lightly and planed off little spots here and there, the fine striæ running south 30° west, but at half a mile to the eastward the granite is completely moutonné and the grooves run south 40° west.

ROCKS OF THE HURONIAN BELT.

The portion of lake Huron between the main north shore and the Manitoulin chain of islands is called the North channel. The quartzites, schists, etcetera, of the great Huronian belt, standing nearly on edge, strike very regularly about east and west for many miles along the north shore and through numerous islands of this channel. A few miles west of La Cloche an intrusive mass of red granite, having a nearly circular form and a diameter of about four miles, interrupts the run of the quartzites, etcetera, which, however, continue their regular east-and-west course on either side of it. On Benjamin island the hills in the central part of the granite intrusion rise to a height of 160 feet above lake Huron. Resting on the southern flanks of this island are small areas of Black river limestone near the level of the lake, while larger flat-lying portions of the same formation, constituting Amedros, Clapperton and Hook islands, approach the granite closely to the east, south and west.

ANCIENT EROSION.

Typical Erosion Surface.—Benjamin island, which is about two miles long, consists of hummocks of granite of various heights up to 160 feet, and these present bluffs toward all points of the compass. In many

places on this island, where the rock surface has been sheltered from glacial action, whether level, sloping or perpendicular, it is eroded into pits, hollows, hummocks and knobs and has a very rough appearance. These surfaces bear a strong resemblance, on a large scale, to the pitted exterior of an *aërolite*. This is well shown in figure 1, a drawing from a photograph, which represents a granite surface on Benjamin island.

The hollows sometimes take the form of long but not deep caves, undermining the perpendicular faces of steps in the rock which rise at intervals of ten to twenty feet from one horizontal joint to the next above. At other times they are sack-shaped cavities, extending from a few inches to several feet into the rock. As on George island, already referred to,

FIGURE 1.—Eroded Granite Surface near the north End of Benjamin Island.

this form is very common on steeply sloping surfaces, and in these cases the upper side of each cavern is generally covered with an arch which thins away toward the front (figure 2).

In half a dozen cases observed on Benjamin island, the pits at various heights from the level of the lake up to 100 feet or more were filled *in situ* with horizontal layers of the fossiliferous Black river limestone like that which overlies the granite at lower levels. The limestone entered into all the little irregularities of the pits and adhered firmly to their walls, thus helping to retain its hold in these cavities during the time the present surface has been exposed to the weather. The little patches constitute veritable, if minute, inliers of the Black river formation. The limestone probably originally filled all the inequalities in the granite

FIGURE 1.—PIT IN GRANITE CONTAINING ARENACEOUS LIMESTONE IN SITU.

FIGURE 2.—GAP EXCAVATED ALONG JOINTS IN GRANITE TRANSVERSE TO THE COURSE OF GLACIATION.

ROCK DECAY, DEPOSITION AND GLACIATION.

surface on which it was deposited, but along with the great mass of the formation was removed by denuding agencies, mostly prior to the glacial epoch.

Figure 1 of plate 16 represents some arenaceous limestone remains *in situ*, in one of the pits, on a steeply sloping granite surface at the south end of Benjamin island. The eroded surface of the rock on this island also occasionally exhibits curious low, circular ridges three or four feet in diameter and three or four inches in height, weathered out of the granite (figure 3). Each circle is surrounded by a slight depression, while the space within has the form of a very shallow basin with a small mound rising in the center. The granite of the mound and the annular

FIGURE 2.—Section of eroded Surface of Granite with Lower Silurian Limestone resting upon it.

ridge are rather finer grained than the surrounding mass. They are apparently of concretionary origin.

Conditions affecting Erosion.—Under what conditions did the ancient erosion above described take place? The facts above presented, as well as others about to be mentioned, would appear to indicate that it occurred at the bottom of the deep sea, for it is known that rock-surfaces do sometimes become pitted or dissolved under deep water. If this erosion had taken place on land we should have evidence of deeper decay in the substance of the rock, while the subsidence of the land to permit the deposition of the fossiliferous limestone upon it would be indicated by detrital shallow-water deposits, such as sandstones and

conglomerates. The occurrence of pure limestones in the pits and ovens of Benjamin island, in one case in the face of a perpendicular bluff, as

FIGURE 3.—Low, circular Ridge, four Feet in Diameter, on weathered Surface of Granite.

well as of the nearly horizontal beds of greater extent upon the sound though uneven surface of the granite, such as might exist at great depths in the sea, but not upon the land, shows that the rock had become thus eroded and was destitute of other covering when the limestone began to be deposited upon it. The sea bottom then rose to a more moderate depth, so that an abundant fauna found a suitable habitat and the formation of the limestone commenced. The area now occupied by the North channel of lake Huron might at that time have formed

a quiet arm of the sea, with hills of Huronian rocks to the north, as at present, and similar hills to the south, which have since sunk to lower levels, but of which traces remain.

CONTACTS OF ARCHEAN WITH PALEOZOIC ROCKS.

Along the junction of the Archean and Paleozoic rocks occurring between the foot of lake Ontario and the head of Georgian bay the actual contact of the Potsdam sandstone and Black river limestone with the Laurentian gneiss may be seen in a great number of places. The surface of the gneiss, which can be observed to pass under the nearly horizontal beds of the newer rocks, is generally rough or angular, quite hard and fresh looking, as if it had never been exposed to weathering on land. The points and promontories of these flat-lying strata which stretch northward from the general line of the basal geographic boundary of the Silurian rocks in this region lie on the lowest ground or in the orographic depressions of the southern margin of the Archean area, where they found shelter from glacial erosion or preglacial decay. It is doubtful if either these or higher formations ever extended over a large part of the Archean area to the northward, as is commonly supposed, although judging from the vast amount of their ruins which have been carried south they must have been much more extensive before the glacial epoch

than at present. The middle Silurian inlier in the great orographic depression of lake Temiscaming on the Ottawa, which will be again referred to, is another example of these newer and undisturbed strata occupying ground much below the general level of the surrounding country.

On the north shore of lake Huron, between Killarney village and Spanish river, and again on Grand Manitoulin island, in the vicinity of Shéguenda and westward of that village, the Lower Silurian strata rest almost horizontally upon the flanks of bold bare ridges of quartzite, the stratification of which stands nearly vertically. The angles of slope of the ancient rock-surfaces exposed to the weather along the crests of these ridges are continued, as far as can be observed in sections, under the newer rocks forming their flanks, so that the former represent only the upper parts of ridges, once much higher, but now partly buried under the strata which accumulated around them in Silurian times. A high ridge of quartzite runs along the north side of Frazer bay, and a similar ridge, but less elevated, forms the northwest side of Killarney bay. The summits and sides of these ridges are, for the most part, thoroughly smoothed and striated by glaciation, but near their southern bases occasional spots are found which have escaped the action of the ice and which show evidence of great antiquity, the flinty rock being worn into pits and hollows, but with no sign of granular or textural disintegration. A broken fringe of Black river limestone skirts the base of the quartzite range on the northwest side of Killarney bay, and the erosion of the spots referred to may be of pre-Paleozoic date, having been preserved till now by a thin limestone covering, or they may represent portions of the surface as it existed immediately before the glacial epoch which have escaped the smoothing action of the ice, but this is not so likely.

In the neighborhood of the quartzite ridges only those beds of the newer rock which are in contact with the old formation contain angular fragments of the rocks on which they lie. The layers containing this débris are usually of no great thickness and the fragments themselves are of small size; but close to Shéguenda, to the north and south of the village, are pretty thick strata composed of sharply angular pieces of quartzite of all sizes packed closely and confusedly together, with only enough calcareous cement to consolidate them. This locality is about twenty miles south of the main Huronian area left uncovered by Silurian strata and the water of lake Huron.

On La Cloche island and elsewhere in the vicinity numerous ridges and domes of Huronian quartzite protrude through the flat-lying Silurian limestones and shales, so that the former resemble masses which have been forced up through them from below, whereas they are only higher portions of the old sea bottom, around which the horizontal beds

of the much newer rocks were deposited. The quartzites appear to have undergone no change in texture, structural attitude or even surface contour since a period vastly antedating Silurian time. Such facts as these indicate pretty clearly that the physical features of the older parts of our continent had their origin at a very remote period.

RIVER AND LAKE CHANNELS DUE TO ROCK DECAY.

The writer has elsewhere* shown that many of the long, straight valleys in the Archean regions of Canada, now often occupied by straight river stretches, by long, narrow lakes, and by inlets of the larger lakes, are due to the decay and removal of wide greenstone dikes or of parallel dikes, together with the belts of rock between them. When the depressions along these decayed dikes are not overspread by water they form valleys, more or less filled up with drift, so that the greenstone in the bottom may be only exceptionally exposed to view. Copious springs frequently issue from the drift in these valleys.

The long, narrow and straight inlets of the northern part of Georgian bay have had their origin along the courses of dikes of this class. The continuation of each of these channels out into the bed of the lake in front of it is distinctly marked by a straight line of deeper soundings, having the same direction as the fiord itself. Collins, Key, Henvey and Byng inlets, each about twelve miles long, but averaging only 200 or 300 yards in width, are good examples of these fiords. Among the more conspicuous of the long, narrow and nearly straight valleys of the Archean region which have been formed in the manner just described may be mentioned that of Onaping lake, 30 miles long, north of lake Huron; Long lake, 52 miles long, north of lake Superior, and Sepiwek lake and Nelson river below it, 96 miles long, north of lake Winnipeg. The greenstone of the dike or dikes along the course of these channels may be seen only on islets and points or in patches adhering to the country rock on either side. In the case of Long lake an immense greenstone dike emerges from its southern extremity and runs into the hills in the same bearing as the central line of the lake.

The course of the Mattagami river, the central branch of the Moose, is guided in its northward trend for a distance of 160 miles from the head of lake Kenogamisse by a number of long dikes of greenstone from 200 to 1,000 feet wide, all having nearly the same direction.† In a typical section of this part of the river we find that the central portion of the dike is coarsely crystalline and more deeply eroded than at the sides and

* Reports Geol. Survey Canada, 1870, p. 331; 1875, p. 315; 1878, p. 15 CC; Report Bureau of Mines Ontario, 1891, p. 76; Bull. Geol. Soc. Am., 1890, p. 300.

† Report of R. Bell: Geol. Survey of Canada for 1875, p. 315.

that it has given rise to the main channel of the stream. The gneiss has been altered and shattered for some distance from the walls of the dike, and this has also suffered decay and erosion and now forms a parallel channel on either side of the central one. Between them the finer grained, hard and undecayed greenstone constituting the outer portions of the dike rises up in the shape of ridges and chains of islands, so that the river flows as a main central channel more or less separated from the smaller lateral ones.

Lake Temiscaming, on the Ottawa, like the Montreal river, which enters its southern extremity, appears to follow the course of a great decayed dike or set of parallel dikes. It lies in a narrow depression which cuts across the general strike of the Archean rocks of the region, and its surface is probably 500 feet below the average elevation of the surrounding country. Its width is only from a mile to two miles, and it has a length of thirty-five miles, but the channel is continued in Deep river, the name given to the section of the Ottawa immediately below it. Temiscaming is an Ojibwé word meaning "deep lake," and it is well named, for in one part it measures, according to Mr A. E. Barlow, 1,800 feet in depth. The bottom is covered very unequally with drift, which has been pushed into it by glaciers at different times, and the deepest part of the bottom may still be a considerable height above the sound greenstone below this filling. If we suppose this height to be 300 feet, then we have a total of 2,600 feet as representing the average height of the surrounding country above the bottom of the excavation. The surface of the lake is 612 feet above the sea, so that the bottom of the water is about 1,200 feet, and that of the whole rock excavation may be 1,500 feet or more below this level. This may, therefore, be regarded as a very deep cutting across a country, the general aspect of which is that of a mammillated plateau with few great inequalities.

The islands toward the north end of the lake consist of limestone of the Niagara formation, which also rises in cliffs at its extremity and extends northward a considerable distance beyond the lake in the same orographic depression, while the Upper Ottawa river falls over the side of this valley by a rapid descent from the plateau to the eastward.

This valley therefore existed before the date of the Niagara formation, and it is probable that under the limestone just mentioned may be found older members of the Silurian system. Greenstone dikes which so frequently cut the Archean rocks of northern Canada have never been found to traverse the overlying Silurian, and we are therefore warranted in supposing that the valleys which mark the courses of the decayed dikes among the former class of rocks were mostly formed before the deposition of the Paleozoic strata.

ABSENCE OF PALEOZOIC DEPOSITION AND ITS SIGNIFICANCE.

The greater part of the Archean area referred to probably never received any deposit of Paleozoic rocks upon it, so that although numerous long and deep valleys, such as those which have been referred to, existed since an earlier period, no evidence of such rocks may ever be found in the larger number of them.

The depressions already mentioned as filled with outliers and projecting tongues of Potsdam sandstone and Black river limestone along the northern boundary of the Paleozoic formations between Georgian bay and the foot of lake Ontario also go to prove, as we have seen, that the surface of the Archean rocks had been reduced to something like its present level and aspect before these beds were laid upon them.

EFFECTS OF GLACIAL DENUDATION.

In a former paper* the writer described the effects of glacial denudation in forming valleys and water channels along the courses of ancient lines of crushing and subsequent decay in the crystalline rocks north of lake Huron. A similar effect has been produced in this region, where groups of parallel joints run close together, with a considerable breadth of more solid rock on either side, and where the comparatively thin walls between them have become softened by the long continued penetration of surface water. It is probable that the deep decay along these joints and lines of fracture took place in pre-Paleozoic times. The disintegrated rock along them was removed by glacial erosion with almost equal facility, no matter what their course may have been, even where the resulting channels are narrow and lie transversely to the direction of the excavating force. A good example of this, on a small scale, is shown in figure 2 of plate 16, which is taken from a photograph of a gap in the granite on George island, at the northwestern extremity of Georgian bay.

*See Report of the Ontario Bureau of Mines, 1892.