cultures from both the mesenteric and thoracic glands being positive in every animal. With the tubercle bacilli growths were more difficult to obtain, and it is obvious that rats and mice destroy these bacilli much more quickly than they do John's bacillus. If one is dealing with a bacillus that develops preferably in a special part of the anatomy of the animal used, then it will be found in this situation as is the case with John's bacillus. An emulsion of John's bacillus inoculated into the peritoneal cavity of a rabbit or mouse will, in some of the animals, eventually produce disease of the intestines and mesenteric glands, where a large number of bacilli may be found, while in the thoracic glands they may or may not be rare. One has here, however, an entirely different factor coming into play—that is to say, the question of the site of the animal body most suitable for the growth and multiplication of the particular bacillus. The place of predilection for the development of a bacillus has often little or no relation to the site of the primary distribution immediately after inoculation.

The inoculation of a certain quantity of oil into the peritoneal cavity to prevent any absorption of septic material has been recommended and has been practised fairly extensively, especially by continental surgeons. A few experiments bearing on this question were also performed, oil being inoculated from 15 minutes to 24 hours before the injection of the bacilli. Although the number of experiments in this direction have not been numerous, they demonstrate clearly that the oil does not prevent the escape of bacteria from the peritoneal cavity and their absorption by the thoracic glands, and it is doubtful if it in any way hinders the dissemination of the bacilli.

Incidentally, the above experiments point to a method by which acid-fast bacilli may be isolated from contaminating micro-organisms and recovered in pure culture in a relatively short space of time, and this means was utilised to isolate John's bacillus from the putrid gut of a sheep infected with John's disease. Tubercle bacilli were also isolated in the same way from emulsions of cultures purposely contaminated.

The time necessary for the escape of infectious material from the peritoneal cavity is thus often only a question of minutes, and it is obvious that even during an operation if the general peritoneal cavity is accidentally contaminated from the rupture of a localised abscess, &c., ample time is given before the end of the operation for the thoracic glands to become infected, and if these glands are penetrated there is a danger of the infection becoming generalised. These facts may explain the beneficial effect of Fowler's position after abdominal operations.

The expenses of these researches were partly paid for by a grant from the London University (Henry Dixon Fund), and I have also to thank the Royal Society for the grant which has enabled me to purchase the animals used in these experiments.

THE USE OF ARSENICAL COMPOUNDS FOR SYPHILITIC DISEASE OF THE NOSE AND THROAT:

A NOTE ON 16 CASES.

By HAROLD L. WHALE, M.D. CANTAB., F.R.C.S. ENG., ASSISTANT SURGEON, THE METROPOLITAN EAR, NOSE, AND THROAT HOSPITAL.

I HAVE treated and followed the history of these 16 patients as follows: 12 in Mr. W. D. Harmer's and Mr. F. A. Rose's clinics at St. Bartholomew's Hospital, 1 in my clinic at the Metropolitan Ear, Nose, and Throat Hospital, and 3 in private patients.

The number of doses given.—These 16 patients have had 43 injections, varying from 1 per patient to 9 (two cases).

The relative advantage of different preparations.—12 injections were intravenous, "606"; 1 intramuscular, "606"; 26 intravenous, neo-salvarsan, "914"; and 4 intramuscular emulsion of "Allenburys" hyposol. All the cases were adults, and the dose was always the equivalent of 0.6 gm. of "606." When more than one injection has been given to any patient, sometimes the second injection was of a different drug from the first. As regards the clinical results and Wassermann tests obtained, in this small series no one preparation appears to be better than another. The advantage

of hyposol emulsion is the extreme simplicity of giving it. For "914" may be claimed a very low toxicity. Not a single injection of the 26 given caused a sufficient general reaction to incapacitate the patient for a day, and the drug is given with justifiable confidence in the out-patients' clinic. Against "914" it may be said that many more injections are required to produce a clinical cure or a negative Wassermann reaction; this is illustrated in the tables given below. Some patients weary of the repeated injections, and defeat their cure by refusing to reattend hospital. After an intravenous injection the vein presumably becomes thrombosed and obliterated, either completely, or partially by a concentric phlebitis obliterans; and it is often found that on the next occasion another vein must be used. In 2 cases out of this series I was obliged, after several injections, to do an open dissection. The advantages and disadvantages of "606" are the converse of those of "914."

The interval between injections was one week, unless there were some local or general reaction, when it was two weeks.

The frequency of Wassermann tests.—As a rule the blood was tested after every second injection, except when a sudden change in the clinical picture indicated this test at other times. The blood was tested 48 hours after the injection.

Adjuvant treatment.—Between injections some of the patients had mercurial inunctions, others potassium iodide; the former yielded the better results.

The relation of Wassermann's tests to the clinical findings.—This is not exact. It is true that in none of these cases has a positive Wassermann reaction changed to negative without the patient being cured; moreover, if several doses are required the clinical cure will be complete sooner than the blood test becomes negative. But the converse does not hold good; and when after several injections the patient becomes cured of symptoms and progressive signs, but the Wassermann reaction remains positive, the injections have been stopped pending further recurrences of disease, which in this small series have not supervened during three months. The improvement of the general condition is sometimes most marked, more particularly when there has been some general reaction. In one case the red cell count rose, with two injections of hyposol, from 3,500,000 to 5,200,000 per c.mm.

The diseases treated may be classified as follows:—

Inflammation or Necrosis of Bone or Cartilage.

No.	Bone or cartilage affected, and lesion.	Drug and number of injections.	Result.
1 ...	Ethmoid necrosis.	... 8, of "914."	{ Separation of sequestrum; disease arrested.
2 ...	Perforated hard palate (old-standing).	... 1, of "606."	Unaffected.
3 ...	Perforated hard palate, 3 years' duration, ulceration now active.	... 3, of "606."	{ Improved, partly healed.
12 ...	Perforated hard palate, one week's history.	... 1, of "914."	{ Cured, healed over.
4 ...	Gummata of septum nasi and frontal bone.	... 2, of "606."	Cured.
5 ...	Gumma of septum.	... 7, of "914."	"
6 ...	Gummata of skull and of epiglottis.	... 1, of "606."	"
10 ...	Periostitis of nasal bone.	... 2, of "606."	Improved.

Laryngitis with no Gross Lesion.

No.	Injections.	Result.
1	9, of "914."	Cured.
5	9, of "914."	"
9	1, of hyposol.	Uncertain.
11	2, of "606."	Improved.

Gummatous Laryngitis, Nodular or Pachydermatous.*

12	1, of "914."	Improved.
13	2, of "606," then 1, of "914."	"
14	2, of "914."	Unaffected.

* As a result of the absorption of excrescences in the anterior commissure or on one cord Nos. 12 and 13 were subsequently able to phonate "a," which they had been previously unable to do.

Inflammation of Pharynx without Ulceration.

2	1, of "606."	Cured.
5	7, of "914."	"
7	2, of "914."	"
9	1, of "914."	Improved.
15	1, of "914."	Uncertain.
16	1, of "606," then 1, of hyposol.	Cured.

Inflammation of Post-Nasal Space without Ulceration, but with Exudation of Mucus or Mucopus causing Deafness.†

3		3, of "606."		Deafness cured.
7		2, of "914."		" "
15		1, of "914."		Uncertain.

† In all these cases of deafness the tuning-fork tests had previously shown middle-ear deafness; no labyrinthine cases are included in this series.

Maxillary Sinusitis.‡

8		2, of hyposol.		Much improved.
---	-------	----------------	-------	----------------

‡ The improvement in this case of maxillary sinusitis, however, may have been due to the concurrent irrigation.

Conclusions.—Salvarsan and its allies offer a fair prospect of arresting tertiary syphilitic disease of the bones or cartilages of the nose and throat, syphilitic laryngitis, and non-ulcerative inflammations of the pharynx and nasopharynx. When repeated injections are not contra-indicated, either by technical difficulties or by unwillingness on the part of the patient, neo-salvarsan is the safest preparation to use.

Harley-street, W.

EARLY DIAGNOSIS OF A CASE OF LEPROSY MUCH ASSISTED BY THE X RAYS.

BY THOMAS MILLER, M.D. MANITOBA.

On Feb. 18th a Chinaman, aged 22 years, came to me complaining of general weakness and inability to work. Questioning through an interpreter elicited merely a history of shooting pains in the left leg, but nothing more definite. On examination of the feet and legs one was at once struck by the shortness of the left great toe, in which there was only the proximal phalanx to be felt, and that half the normal size. The nail was half the usual size and very much retracted. The extremity of the second toe was markedly bulbous. There was no history of ulceration or abscess to account for the loss of the distal phalanx, and it was found that an area of the skin corresponding to the first and second toes and the distal halves of the metatarsal bones on both

FIG. 1.

Left foot : advanced changes.

dorsal and plantar surfaces was slightly thickened and anæsthetic to pain and temperature. There was also a scar, three quarters of an inch in length, on the plantar surface of the great toe at the metatarso-phalangeal articulation, of which no history could be obtained.

A tentative diagnosis of leprosy was made, although no other signs of the disease could be found. The hands are normal.

Two days later Dr. E. S. Hasell, of the Royal Provincial Jubilee Hospital, Victoria, obtained X ray photographs of both feet, and these show most interesting alterations in the bones, confirming the diagnosis of leprosy. Of the proximal

FIG. 2.

Right foot: changes commencing.

phalanx of the left great toe only half remains. The distal phalanx of the second toe has almost disappeared and the distal phalanges of the other toes are also undergoing rarefaction, their shadows being more transparent than normal. The same changes are seen in the heads of the metatarsals. The changes in the right foot are similar, but less advanced.

The left great toe is said to have been decreasing in size during two years, but the patient was unaware of the existence of anæsthesia until I directed his attention to it. Hence, I conclude that the earliest sign of leprosy may be alterations in bones which are made manifest by skiagraphic examination, and therefore venture to put this case on record, as any assistance in the early diagnosis of this so serious disease is of great importance and may help other physicians in difficulty.

Victoria, British Columbia.

INTERNATIONAL COMMITTEE FOR POST-GRADUATE MEDICAL EDUCATION.—A series of lectures on balneology and balneotherapeutics, arranged under the auspices of the International Committee for Post-graduate Medical Education, will take place at Karlsbad from Sept. 28th to Oct. 4th. The list of lecturers includes, among others, Professor Dr. Albu (Berlin), Dr. Dautwitz (St. Joachimsthal), Professor Dr. Falta (Vienna), Privy Councillor Professor Dr. Fleiner (Heidelberg), Professor Dr. Fronz (Vienna), Councillor Professor Dr. Glax (Abazzia), Privy Councillor Professor Dr. Goldscheider (Berlin), Councillor Professor Dr. Grödel (Nauheim), Professor Dr. Hoffmann (Düsseldorf), Councillor Professor Dr. von Jaksch (Prague), Professor Dr. Kionka (Jena), Professor Dr. R. Kutner (Berlin), Professor Dr. Lühje (Kiel), Professor Dr. Mohr (Halle), Professor Dr. Plehn (Berlin), Professor Dr. Riehl (Vienna), Councillor Professor Dr. Schauta (Vienna), Privy Councillor Professor Dr. A. Schmidt (Halle), and Professor Dr. Strauss (Berlin). The lectures may be attended by medical practitioners on payment of a registration fee of 10 kr. for the entire series. For programme of lectures and cards of admission application should be made to the Kuramt of Karlsbad.