TRANSACTIONS

OF

THE LINNEAN SOCIETY.

I. Report on two Botanical Collections made by Messrs. F. V. McCONNELL and J. J. QUELCH at Mount Roraima in British Guiana. By N. E. BROWN, A.L.S., and others. (Communicated by Sir WILLIAM THISELTON-DYER, K.C.M.G., F.R.S., F.L.S., Director of the Royal Botanic Gardens, Kew.)

(Plates I.-XIV.)

Read 15th March, 1900.

CONTENTS.

	Page
Introduction. By I. H. BURKILL, M.A., F.L.S.	1
Enumeration of the Plants collected.	8
I. Spermatophyta. By N. E. BROWN, A.L.S. (the Orchidaceæ by R. A. ROLFE, A.L.S.)	18
II. Pteridophyta. By C. H. WRIGHT, A.L.S.	77
III. Bryophyta: Musci. By V. F. BROTHERUS, Ph.D.	88
Hepaticæ. By F. STEPHANI	93
IV. Thallophyta. By G. MASSEE, F.L.S.	101

INTRODUCTION.

By I. H. BURKILL, M.A., F.L.S.

IN the autumn of 1894 Messrs. F. V. McConnell and J. J. Quelch first visited Roraima: they were there again in the autumn of 1898, and on both occasions not only encamped higher on the slopes of the mountain than any previous traveller, but they spent nights on the summit in order that the time given to collecting its flora and fauna should be uninterrupted. The collections of plants were transmitted to Kew, and both are enumerated in this paper.

In 1894 the journey to Roraima commenced (October 16) at Kwaimatta, a village on the open savannah to the south-east in latitude $3^{\circ} 50' N$. The Ireng was crossed at the Karona Falls, the Kotinga at Sokoking, a ford not named on the older maps, but situated a little to the south of the meeting of the Kotinga and Karakanang streams. Thence the way led into the Arabapu Valley (November 3), on the north and west sides of which rise the slopes of Roraima. The return journey was commenced on November 11, and the route taken crossed the Kotinga nearer its source at Orinidoui (apparently the "Orinidouk" of the old maps), and the Ireng again at the Karona Falls.

In 1898 the journey was commenced (August 20) on the lower Mazaruni River. The great bend of this river, where the Peaiman Falls occur, was avoided by a portage from the Kurubung Creek to the upper Mazaruni, which was again left by the Kako River; the Kako was paddled up until a point was reached about twenty miles north-east of Roraima, and thence, leaving the canoes, a track through dense forest was taken until, in the second week of October, the south-eastern face of the mountain was reached. Roraima was left again at the end of October, and the way retraced to the lower Mazaruni.

Three days in 1894, nine in 1898, were spent in making collections upon the broken plateau which forms the summit of Roraima.

VISITORS TO RORAIMA.

	Year.	Month.	Height reached.	Route to Roraima.	Return Journey.
R. H. SCHOMBURGK	1838	Nov.	Base of Mountain.	From S. over Humirida Mtns.	Retraced steps.
R. H. and RICHARD SCHOMBURGK. }	1842	End Oct. to Dec. 5. End Oct. to Nov. 22.	Base of cliff on S. side.	ditto.	{ At first S., then to W. of Roraima. Retraced steps.
KARL APPUN	1864	End Jan. to end Feb.	Base of cliff on E. and S. sides.	From N. over Merume Mtns.	Towards S.E. to Pirara.
C. B. BROWN and J. G. SAWKINS.	1869	Early Feb.	Upper slopes.	From S.E.	
J. W. BODDAM-WHETHAM and M. McTURK.	1878	March.	ditto.	From N. over Merume Mtns.	Retraced steps.
D. BURKE	1881		ditto.		
SIEDEL	1884	Feb. to Apr. and Nov.	ditto.	From N.	
E. F. IM THURN and H. I. PERKINS.	1884	Nov. to Dec.	Summit from the S. side.	From R. Potaro to Ireng, then to Roraima from S.E.	Retraced steps.
E. KROMER	1886	Nov.	ditto.		
DRESSEL	1887	Oct.	ditto.	From N. over Merume Mtns.	
F. V. McCONNELL and J. J. QUELCH.	1894	Nov.	ditto.	From S.E.	To S.E. by a some- what parallel track.
F. V. McCONNELL and J. J. QUELCH.	1898	Oct.	ditto.	From N. over Merume Mtns.	Retraced steps.

Appearance and Structure of Roraima, and its Geographical Relations to the Mountain-systems of South America.—Im Thurn and Perkins, who made the first ascent on December 18, 1884, reached the top under unfavourable conditions: they had ascended that day from their base-camp at 5400 feet, and only reached the summit in the afternoon, when clouds almost invariably envelop it; they had no means of remaining there, and were forced to return after a very short stay of three hours. Perkins speaks of these clouds as impenetrable (Proc. Roy. Geogr. Soc. vii. 1885, p. 532)*; at all events they limited the range of exploration, and the report was made that “the vegetation on the summit is extremely scanty and insignificant, there being no trees, only small bushes from three to six feet in height, growing at long intervals.” It has been reserved for McConnell and Quelch to ascertain that there exist in favoured places trees which attain a height of no less than forty feet.

Of the ascents of Dressel and Kromer I can glean no more than is in a brief notice in ‘Timehri,’ 1887, p. 330. Neither can have remained on the summit for more than a few hours.

The features of this summit are:—Blackened elevated ridges; irregularly terraced and rugged; winding gullies which drain away the superabundant rain through shallow pools; patches of sand in their shelter with isolated small bushes; clumps of vegetation lodged in the most protected spots; piles of wind-cut rocks without a sign of plant-life, rising into pinnacles; no soil; the keen northerly wind, and mists which gather at least after noon on almost every day in the whole year. The highest pinnacle rises to 8740 feet; the deepest gully may be 400 feet below it.

The known sides of the mountain which face north-east and south-east, produce rocky savannah up to 5000 or 5400 feet, the rocks so numerous that Appun likens the country to a grave-yard; then forest to 7100 feet, dense below but stunted and more open above; and from this altitude rise the precipitous cliffs, which give to the whole the appearance of a vast fortress with a tree-clad glaciis slope.

These precipices were deemed impossible of ascent by the travellers they had baffled until Whitely indicated, and Im Thurn showed practicable, the one known ledge leading to the summit.

The rock is a quartzose sandstone interbedded with diorite. The topmost and thickest bed of diorite comes to the surface where the dense forest is; the lower beds make a series of terraces down the lower slopes. These beds lie perfectly horizontal without folding or other sign of any great disturbance. Diorite lends itself to the formation of a soil †, the sandstone does not.

Such sandstones as constitute Roraima have a very wide extension in South America; they pass eastward into Surinam; Duida is made of them; they form the bed-rock of the llanos of Venezuela, and appear in the Caripe mountains to the east of Caracas; in Brazil the area occupied by them is immense.

* Cf. Im Thurn in ‘Timehri,’ 1885, p. 41.

† The Indians prepare their provision-grounds on the diorite (see Brown & Sawkins, Reports, p. 17), and those living under the south-eastern face of Roraima have placed them far up the sister-mountain of Kukuenaam, where the diorite comes to the surface (Quelch, in ‘Timehri,’ 1895, p. 164).

All this sandstone area belongs to the oldest land of South America: against the edge of it the Andes were heaped up—newer land which grew to its present form in the tertiary period, and formed a link of high ground round the head of the then existing Amazon valley.

It is significant to us that the chief mountain-systems of South America outside the Andes reach very similar heights and similarly stand more or less parallel to the nearest coast, whence the trade-winds bring an abundance of rain. These mountain-systems are three: (1) the Coast Andes of Venezuela, of direct eruptive origin, but continued eastward from Caracas in the lesser sandstone mountains of Caripe; (2) the Parime mountains with Roraima at the eastern end and Duida at the western end; and (3) the mountains of South Brazil formed of schists tipped at a high angle.

*Heights * north of the Equator.*

Coast Andes.		Parime Mountains.	
Niaguata	9125 feet.	Roraima	8740 feet.
Silla of Caracas	8741 „	Duida	8278 „
Tamaya	8052 „	[Peaks near Duida estimated at 10,000 feet.]	

*Heights * in South Brazil.*

Itatiaia	8999 feet.	Itambe	5960 feet.
Organ Mountains	6609 „ and more.	Serra da Piedada	5874 „
Serra da Caraça	6411 „	Itacolumi	5760 „

Unlike in geological structure the three systems are unlike as well in their relation to the Andes. The Casiquiare, by uniting the Orinoco and Amazon, encircles the Parime mountains, and the undulating country of little elevation to the west of it effectually separates this system from the Andes; the plains of Matto Grosso, &c., wherein rise within a few miles of each other the Paraguay and Madeira, to flow the one north, the other south, separate the Brazilian mountains and the Andes; but the Venezuelan coast-range is most intimately bound to the Andes proper through the Cordillera of Merida.

Spanish settlers in the New World soon came to recognize a belt on the mountains of the tropics suited to their needs and for the growth of their food-plants. They called it the temperate land—“tierra templada,”—and the range of its mean annual temperature may be set down as 15°–20° C. (59°–77° F.). Above the “tierra templada” is the “tierra fria,” below it the “tierra caliente.”

The limits of these belts depend on exposure. In the Venezuelan mountains, according to Sievers (‘Venezuela,’ p. 26), the “tierra fria” extends from about 7200 feet upwards; among the Great Andes it sometimes commences as high as 10,000 feet; on Roraima

* The heights of the Coast Andes are taken from Sievers’s ‘Venezuela’ (Hamburg, 1888), pp. 277, 278; those of Duida and neighbouring peaks from Sir Robert Schomburgk’s narrative in Proc. Roy. Geogr. Soc. x. 1840, p. 245; and those of the Brazilian mountains, Itambe excepted, from Liais, ‘Climats, Géologie, Faune et Géographie botanique du Brésil’ (Paris, 1872), pp. 45–49. Most astounding are the erroneous statements published regarding the altitudes of the Brazilian mountains; for instance, the Serra dos Pyrenas, estimated previously to reach 9700 feet, proves to be no more than 4543 feet (Cruls, ‘Relatorio da Comissão exploradora,’ Rio, 1894, p. 26).

it descends presumably into the forest-belt or to the base of the cliffs *. Roraima bears the most eastern patch of "tierra fria" upon the northern side of the Amazon.

Some hundreds of feet above the commencement of the "tierra fria" is the limit of trees. There are reasons for believing that both the Coast Andes and Roraima just reach this limit; they do not distinctly rise above it.

The Flora.—Passing on to a consideration of the nature of the flora of Roraima I have given in the table on pp. 8–16 as complete a list as is now possible of the species found on the mountain above 5000 feet, and we have in it:

239 Spermatophyta, of which 121 (50·6 per cent.) are endemic.

88 Pteridophyta, of which 16 (18·2 per cent.) are endemic.

63 Bryophyta, of which 15 (23·8 per cent.) are endemic.

11 Thallophyta, of which 3 (27·3 per cent.) are endemic.

The proportion of endemic Spermatophyta may seem large, but does not exceed that on record for some of the mountains of Mexico. Of far greater interest is the number of endemic genera. The law that mountains by their isolation and extension, as well as by their latitude, produce endemic genera, is illustrated by their number in the ranges of Cis-equatorial South America; thus, there are eleven among the Spermatophyta on Roraima, and only two on the Coast Andes, which are comparatively small and not isolated; but I am aware of no fewer than thirty-six in the extensive Andes of Colombia, including with them the Cordillera of Merida.

The endemic genera of Roraima are enumerated on p. 7. They belong to as many orders. One of them—*Heliamphora*—has no kindred in South America, but belongs, like *Cyrilla*, to a North-American group; all the rest have more or less close allies in genera of the South-American continent. *Ledothamnus*, however, deserves further remark because it is one of the very few Ericaceæ with ericoid leaves which exist in the New World.

* This belief is based chiefly upon the nature of the flora. With regard to temperature the following may be added:—Hann ('Handbuch der Climatologie,' Stuttgart, 1883, p. 152), after quoting Boussingault's estimate for the Andes of the tropics that 57° C. in mean annual temperature is lost for every 100 metres ascended, and Humboldt's for Mexico and Colombia of 53° C. for every 100 metres, adds: "as a general rule for the tropics one may allow 58° C. for the amount of heat lost in every 100 metres ascended." This is equivalent to 4° F. for every 100 feet. As the mean annual temperature of Demerara is 81°, we obtain for Roraima these figures:—65° F. at 4000 feet, 60° at 5400, 53° at 7100, 47° at 8600, and 46° at 8740 feet. The following are all the recorded observations of which I am aware:—

	Night.	6–8 A.M.	Midday and afternoon.
Summit, Quelch in 1894	47° F.	49–52° F.	64°·5 F.
„ im Thurn			54° (in mist).
About 7000 ft., Appun	50°		
About 6400 ft., Quelch	50°		
About 5400 ft., R. Schomburgk . .	52°		
„ „ „ im Thurn	48°		
At the base, R. Schomburgk	58° and below	62°	{ 87° (shade).
„ „ „ „	58°·5		{ 100° (sun).
„ „ Brown „		59°	
„ „ Boddam-Whetham . .	Below 60°		
„ „ Quelch	54°		

The endemic genera of the Coast Andes are *Euosmia* and *Caracasiasia*. *Euosmia*, Humb. & Bonpl., belongs to the Rubiaceæ, and was found by Humboldt and Bonpland in the mountains of Caripe; it is a little obscure and needs re-examination. *Caracasiasia*, Szyszy. (*Vargasia*, Ernst) is a genus of the Marcgraviæ, with two species found near Caracas; it is nearly related to the genus *Ruyschia*, found in the West Indies and from Guiana to Peru.

Following the list of endemic genera on p. 7 are three lists to be considered together. All are of genera represented in the tierra fria of the Andes: the first contains seven found on Roraima, throughout the Andes to Chili, and in South Brazil; the second contains fourteen found on Roraima and in the Northern Andes, but reaching neither Chili nor the mountains of South Brazil; the third list contains thirteen found in the Andes, which have passed along the Coast Andes to Niaguata or the Silla of Caracas*, but have not yet been found on Roraima.

These lists illustrate two points:—in the first place, they indicate how the Coast Andes belong essentially to the Andes proper,—and the community of genera extends to species in *Cinchona*, *Acæna*, *Cardamine*, and *Berberis*; in the second place, that Roraima and the mountains of South Brazil only have in common among woody plants montane genera so widely Andine as to pass from Colombia to Chili. One is then led to suspect that the mountains near Rio de Janeiro obtained their Andine shrubby genera from the south-west, and that in their case the plains of Matto Grosso have been a very similar barrier to immigration as the wooded hills on the west of the Casiquiare have been to the Parime mountains.

Six more genera are named in a last list; they are common to Roraima and to the mountains of South Brazil, without reaching Chili; but they all descend to low levels in the Province of Alto Amazonas, and obviously are not montane in a restricted sense.

Other genera which cannot be regarded as truly montane, contribute montane forms to the flora of Roraima. The most notable of these is *Abolboda*, of which genus *A. Sceptrum* is by far the largest species.

Saxofredericia regalis is, like *Abolboda Sceptrum*, the largest of its genus. *Lisianthus Elizabethæ* and another species collected under this name by Appun are among the largest-flowered of these gentians; and *Utricularia Humboldtii* so impressed Sir Robert Schomburgk by its showy blossoms as to cause him to name the place where he first saw it the El Dorado Swamp, after the treasure-city Raleigh and others sought in Guiana.

im Thurn has suggested a resemblance between the floras of Roraima and the Brazilian mountains (Trans. Linn. Soc., ser. II. Bot. ii. (1887), p. 257), but the similarities he notes are much more due to similarities of climate than to any evidently intimate relationship between them in times past.

A very rough estimate of the number of genera of Spermatophyta in the flora of British Guiana places the total at 1070. The orders most strongly represented are:—Leguminosæ 86, Orchidaceæ 66, Rubiaceæ 54, Compositæ 44, Gramineæ 43, Euphorbiaceæ 31, and Melastomaceæ 28. On Roraima above 5000 feet, the order

* Ernst in Journ. Bot. x. 1872, p. 261; in Bull. Soc. Bot. France, xxii. Rev. Bibl. p. 239; in 'Idea general de la Flora de Venezuela' in 'Estudios sobre la Flora y Fauna de Venezuela' (Caracas, 1877), pp. 212-235; and elsewhere.

Endemic Genera (2 occur on the Kaieteur Savannah).

<i>Heliamphora</i> (Sarraceniaceæ).	<i>Notopora</i> (Vacciniaceæ).
<i>Leitgebia</i> (Violaceæ).	<i>Ledothamnus</i> (Ericaceæ).
<i>Blepharandra</i> (Malpighiaceæ).	<i>Connellia</i> (Bromeliaceæ).
<i>Crepinella</i> (Araliaceæ).	<i>Nietneria</i> (Liliaceæ).
<i>Chalepophyllum</i> (Rubiaceæ).	<i>Everardia</i> (Cyperaceæ).
<i>Quelchia</i> (Compositæ).	[<i>Enterosora</i> (Filices)].

*Genus found at either end of the Parime Mountains.**Stegolepis* (Rapateaceæ).*Genera of the Andes (as regards S. America) extending from the North to Chile, also to Roraima and the Mountains of South Brazil.*

<i>Drimys</i> (Magnoliaceæ).	<i>Gaultheria</i> (Ericaceæ).
? <i>Momina</i> (Polygalaceæ).	<i>Pernettya</i> (Ericaceæ).
<i>Weinmannia</i> (Saxifragaceæ).	<i>Puya</i> (Bromeliaceæ).
<i>Relbunium</i> (Rubiaceæ).	

Genera of the Northern Andes (as regards S. America), not overpassing Bolivia southwards, reaching Roraima, but not South Brazil.

<i>Chetolepis</i> (Melastomaceæ).	<i>Sophoclesia</i> (Vacciniaceæ).
<i>Centronia</i> (Melastomaceæ).	<i>Sphryspermum</i> (Vacciniaceæ).
<i>Monochaetum</i> (Melastomaceæ).	<i>Befaria</i> (Ericaceæ).
<i>Viburnum</i> (Caprifoliaceæ).	<i>Grammadenia</i> (Myrsinaceæ).
<i>Psammisia</i> (Vacciniaceæ).	<i>Scaphosepalum</i> (Orchidaceæ).
<i>Cavendishia</i> (Vacciniaceæ).	<i>Lepanthes</i> (Orchidaceæ).
<i>Vaccinium</i> (Vacciniaceæ).	<i>Tofieldia</i> (Liliaceæ).

Genera extending from the Andes along the Coast Andes of Venezuela to the Neighbourhood of Caracas, but not yet found on the Parime Mountains.

<i>Berberis</i> (Berberidaceæ).	<i>Sedum</i> (Crassulaceæ).
<i>Cardamine</i> (Cruciferae).	<i>Cinchona</i> (Rubiaceæ).
<i>Arenaria</i> (Caryophyllaceæ).	<i>Galium</i> (Rubiaceæ).
<i>Adesmia</i> (Leguminosæ).	<i>Espeletia</i> (Compositæ).
<i>Acæna</i> (Rosaceæ).	<i>Gaylussacia</i> (Vacciniaceæ).
<i>Potentilla</i> (Rosaceæ).	<i>Muehlenbergia</i> (Gramineæ).
<i>Escallonia</i> (Saxifragaceæ).	

Genera found on Roraima and the Mountains of South Brazil, avoiding the Chilean Andes, but occurring at low levels in Alto Amazonas, &c.

<i>Remijia</i> (Rubiaceæ).	<i>Hedyosmum</i> (Chloranthaceæ).
<i>Ladenbergia</i> (Rubiaceæ).	<i>Masdevallia</i> (Orchidaceæ).
<i>Stiftia</i> (Compositæ).	<i>Stenoptera</i> (Orchidaceæ).

of the Orchidaceæ comes first, Melastomaceæ second, Compositæ third, and Rubiaceæ and Gramineæ bracketed fourth.

It is unfortunate that all the plants collected on the upper parts of Roraima have been gathered in the last three months of the year. February brings to Roraima a few bare trees (Appun, 'Unten den Tropen,' ii. p. 226); March brings copious Gesnerads into flower, at least at the foot of the mountain (Boddam-Whetham, 'Roraima'); and April is a month producing more flowers than November (Siedel *vide* im Thurn in Trans. Linn. Soc., ser. II. ii. p. 250); but I am unable to record anything collected above 5000 feet during these months.

In conclusion, my thanks for help are due to Messrs. McConnell, Quelch, and C. B. Clarke, to my colleagues, and also to Professor T. McKenny Hughes and Dr. William Pollard.

*The Flora of Roraima above 5000 feet, as now known, with the Geographical
Distribution of each Species.*

		N. Brazil.	S. Brazil, Paraguay.	Colombia to Bolivia.	Venezuela and Trinidad.	W. Indies.	Central America.	Elsewhere in America.	Old World.
MAGNOLIACEÆ.									
<i>Drimys granatensis</i> , Mutis	Ledge.	..	*	*	*	..	*	S.	
SARRACENIACEÆ.									
<i>Heliamphora nutans</i> , Benth.	to Summit.								
VIOLACEÆ.									
<i>Sauvagesia erecta</i> , Linn.	to Ledge.	*	*	*	*	*	*		*
<i>Leitgebia Imthurniana</i> , Oliver	Ledge & Summit.								
POLYGALACEÆ.									
<i>Polygala glochidiata</i> , H. B. & K.	to 5400 ft.	*	*	*	*	*	*		
" <i>hygrophila</i> , H. B. & K.	to 5400 ft.	*	*	*	*	*	*		
" <i>longicaulis</i> , H. B. & K.	to 5400 ft.	*	*	*	*	*	*		
" <i>variabilis</i> ?, H. B. & K.	to 5400 ft.	*	*	..	*	*	*		
<i>Monnina cacumina</i> , N. E. Br.	Summit.								
GUTTIFERÆ.									
<i>Moronobea intermedia</i> , Engl.	at 5400 ft.								
<i>Clusia palmicida</i> , Rich.	to 6000 ft.	*	..	..	*				
TERNSTREMIACEÆ.									
<i>Marcgravia</i> sp.	at 5400 ft.								
<i>Bonnetia Roraimæ</i> , Oliver	Summit.								
<i>Archytæa multiflora</i> , Benth.	to 6000 ft.								
MALVACEÆ.									
<i>Sida linifolia</i> , Benth.	to 5400 ft.	*	*	*	*	*	*		Africa.
MALPIGHIACEÆ.									
<i>Byrsonima crassifolia</i> , H. B. & K.	to 5400 ft.	*	*	*	*	*	*		
<i>Heteropteris oleæfolia</i> , Griseb.	to Upper slopes.								
<i>Tetrapteris rhodopteron</i> , Oliver	to 5400 ft.								
" sp.	to 5400 ft.								
RUTACEÆ.									
<i>Ravenia ruellioides</i> , Oliver	Upper slope.								
ILICACEÆ.									
<i>Ilex retusa</i> , Klotzsch	to Summit.								
" <i>apicidens</i> , N. E. Br.	Summit.								
" <i>vacciniifolia</i> , Klotzsch	to 6000 ft.								

		N. Brazil.	S. Brazil, Paraguay.	Colombia to Bolivia.	Venezuela and Trinidad.	W. Indies.	Central America.	Elsewhere in America.	Old World.
CYRILLACEÆ.									
	<i>Cyrilla brevifolia</i> , N. E. Br.								Summit.
LEGUMINOSÆ.									
	<i>Swartzia</i> sp.								at 5000 ft.
	<i>Cassia insignis</i> , N. E. Br.								to Upper slopes.
	„ <i>polystachya</i> , Benth.								to Upper slopes.
	<i>Pithecolobium ferrugineum</i> , Benth. ..								to Upper slopes.
ROSACEÆ.									
	<i>Rubus guianensis</i> , Focke								to 7100 ft.
	„ <i>urticifolius</i> , Poir.	*	*	*	..	..	*		to 7100 ft.
SAXIFRAGACEÆ.									
	<i>Weinmannia guyanensis</i> , Klotzsch ..								to Summit.
	„ <i>elliptica</i> , H. B. & K. ..	..	..	*	..	..	*		to 6000 ft.
	„ <i>fagaroides</i> , H. B. & K.	..	..	*					Summit.
DROSERACEÆ.									
	<i>Drosera communis</i> , St. Hil.	*	*	*	*	*	*		to Summit.
MYRTACEÆ.									
	<i>Myrtus myricoides</i> , H. B. & K.	..	..	*	*	..	*		to Summit.
	„ <i>roraimensis</i> , N. E. Br.								Summit.
	„ <i>stenophylla</i> , Oliver								Ledge.
MELASTOMACEÆ.									
	<i>Microlicia bryanthoides</i> , Oliver								to Summit.
	<i>Cambessedesia Roraimæ</i> , Rich. Schomb.								to 6000 ft.
	<i>Siphanthera microlicioides</i> , Cogn. ..								to 5400 ft.
	<i>Macairea aspera</i> , N. E. Br.								Upper slopes.
	<i>Tibouchina fraterna</i> , N. E. Br.								Upper slopes.
	„ <i>lasiophylla</i> , Cogn.								to 5400 ft.
	„ <i>aspera</i> , Aubl.	*	*	*	*				to 5400 ft.
	<i>Chætolepis amisandra</i> , Naud.								to Ledge.
	<i>Marcetia taxifolia</i> , DC.	*	*						to 7100 ft.
	„ <i>juniperina</i> , DC.	..	..	*	*				Summit.
	<i>Monochætum Bonplandii</i> , Naud.	..	..	*	*				to Summit.
	<i>Meriania</i> sp.								at 6000 ft.
	<i>Centronia crassiramis</i> , Triana								at 5-50 ft.
	<i>Miconia tinifolia</i> , Naud.	..	..	..	*				to Summit.
	„ <i>holosericea</i> , DC.	*	*	*	*				Upper slopes.
	„ <i>Fothergilla</i> , Naud.	*	*	*	*	*	*		to 5400 ft.
	<i>Tococa aristata</i> , Benth.	*	..	*	*				to 6000 ft.
	„ <i>guyanensis</i> , Aubl.	*	*	*	*				to 6000 ft.
	„ <i>nitens</i> , Triana	*	*	..	*				to 6000 ft.
	<i>Clidemia capitata</i> , Benth.	..	..	..	..	..	*		to 6000 ft.
PASSIFLORACEÆ.									
	<i>Passiflora</i> , 2 (possibly 3) species.								about 5400 ft.
BEGONIACEÆ.									
	<i>Begonia towarensis</i> , Klotzsch	..	..	*	*	*	*		at 5400 ft.
ARALIACEÆ.									
	<i>Didymopanax rugosum</i> , N. E. Br. ..								Summit.
	<i>Sciadophyllum coriaceum</i> , March. ..								at 5400 ft.
	„ <i>umbellatum</i> , N. E. Br.								Upper slopes.
	<i>Crepinella gracilis</i> , March.								Summit.
CAPRIFOLIACEÆ.									
	<i>Viburnum glabratum</i> , H. B. & K. ..	..	..	*	*	*	*		to 7100 ft.
RUBIACEÆ.									
	<i>Remijia Roraimæ</i> , K. Schum.	*							to 6000 ft.
	„ <i>truniflora</i> , Benth.	*							to 6000 ft.
	„ <i>densiflora</i> , Benth.								to 6000 ft.
	<i>Ladenbergia Lambertiana</i> , Klotzsch	*							to 6000 ft.
	<i>Chalepophyllum speciosum</i> , N. E. Br.								Summit.

		N. Brazil.	S. Brazil. Paraguay.	Colombia to Bolivia.	Venezuela and Trinidad.	W. Indies.	Central America.	Elsewhere in America.	Old World.
RUBIACEÆ (cont.).									
<i>Sipanea pratensis</i> , Aubl.	to 5400 ft.	*	*	*	*	*			
<i>Coccocypselum canescens</i> , Willd.	to 5400 ft.	*	*	*	*	*	*		
<i>Retiniphyllum laxiflorum</i> , N. E. Br. .	Upper slopes.								
<i>Psychotria concinna</i> , Oliver	to Summit.								
" <i>crassa</i> , Benth.	to 6000 ft.	*	..	..	*	*			
" <i>inundata</i> , Benth.	to 6000 ft.	*	*						
" <i>Imthurniana</i> , Oliver	to Ledge.								
" 2 spp.	to 6000 ft.								
<i>Declieuxia chiococcoides</i> , H. B. & K. .	to 5400 ft.	*	*	*	*				
<i>Cephaelis axillaris</i> , Sw.	to 5400 ft.	..	..	*	..	*			
<i>Relbunium</i> sp.	to 5400 ft.								
COMPOSITÆ.									
<i>Vernonia chreticifolia</i> , Benth.	to 6000 ft.	*							
" <i>remotiflora</i> , Rich.	to 6000 ft.	*	*						
<i>Eupatorium conyzoides</i> , Vahl	to 5400 ft.	*	*	*	*	*	*	N. & S.	
" <i>roraimeense</i> , N. E. Br. .	Summit.								
" <i>fuscum</i> , N. E. Br.	Summit.								
<i>Mikania pannosa</i> , Baker	at 5400 ft.								
<i>Baccharis Vitis-idaea</i> , Oliver	to Summit.								
" sp., near <i>B. cassinea-</i> <i>folia</i>	Summit.								
<i>Heterothalamus densus</i> , N. E. Br. .	Summit.								
<i>Gnaphalium purpureum</i> , Linn.	to 5400 ft.	*	*	*	*	*	*	N. & S.	*
<i>Verbesina guianensis</i> , Baker	to 5400 ft.								
<i>Pectis elongata</i> , H. B. & K.	to 5400 ft.	*	*	*	*	..	*		
<i>Quelchia conferta</i> , N. E. Br.	Summit.								
<i>Erechtites hieracifolia</i> , Rafin.	to 5400 ft.	*	*	*	*	*	*	N. & S.	Introd.
<i>Stiffia condensata</i> , Baker	to Summit.								
" <i>Connellii</i> , N. E. Br.	Summit.								
LOBELIACEÆ.									
<i>Centropogon levigatus</i> , A. DC.	to Ledge.	*							
" <i>surinamensis</i> , Presl ..	to 5400 ft.	*	*	*	*	*	*		
VACCINIACEÆ.									
<i>Psammisia guianensis</i> , Klotzsch	at 6000 ft.								
" <i>formosa</i> , Klotzsch	at 6000 ft.								
" <i>coriacea</i> , N. E. Br.	Summit.								
" 2 spp.	to 5400 ft.								
<i>Notopora Schomburgkii</i> , Hook. f. .	to 5400 ft.								
<i>Cavendishia</i> sp. (<i>Thibaudia nutans</i> , Klotzsch)	at 6000 ft.								
<i>Vaccinium puberulum</i> , Klotzsch ..	to 6000 ft.								
" <i>roraimeense</i> , N. E. Br. .	Summit.								
<i>Spherospermum Roraimeæ</i> , Klotzsch	to 6000 ft.†								
<i>Sophoclesia</i> sp.	Ledge.								
ERICACEÆ.									
<i>Pernettya marginata</i> , N. E. Br.	Summit.								
<i>Gaultheria setulosa</i> , N. E. Br.	Summit.								
" <i>cordifolia</i> , H. B. & K. (<i>G.</i> <i>Roraimeæ</i> , Klotzsch)	to 5400 ft.	..	..	*	*	..	*		
<i>Ledothamnus sessiliflorus</i> , N. E. Br. .	to Summit.								
<i>Befaria guianensis</i> , Klotzsch	to Ledge.								
" <i>Imthurnii</i> , N. E. Br.	Summit.								
" <i>Schomburgkii</i> , Klotzsch	to 6000 ft.								
MYRSINACEÆ.									
<i>Myrsine</i> sp.	to 6000 ft.								
<i>Cybianthus</i> sp.	at 6000 ft.								

† If synonymous with *Hughesia guianensis*, Klotzsch & Rich. Schomb.

		N. Brazil.	S. Brazil, Paraguay.	Colombia to Bolivia.	Venezuela and Trinidad.	W. Indies.	Central America.	Elsewhere in America.	Old World.
MYRSINACEÆ (cont.).									
<i>Grammadenia lineata</i> , Benth.	to 5400 ft.								
<i>Ardisia Quelchii</i> , N. E. Br.	Summit.								
SAPOTACEÆ.									
<i>Chrysophyllum emarginatum</i> , Klotzsch †	at 6000 ft.	*							
<i>Lucuma rigida</i> , Mart. & Eichl.	at 5400 ft.								
APOCYNACEÆ.									
<i>Mandevilla angustifolia</i> (<i>Echites angustifolia</i> , Benth.) ..	to 6000 ft.								
„ <i>glabra</i> , N. E. Br.	Upper slopes.								
ASCLEPIADACEÆ.									
<i>Nephradenia linearis</i> ?, Benth.	to 5400 ft.	*	..	..	*				
LOGANIACEÆ.									
<i>Antonia ovata</i> , Pohl	to Upper slopes.	*	*						
GENTIANACEÆ.									
<i>Curtia tenuiflora</i> (<i>Schuebleria tenuiflora</i> , G. Don)	to 5400 ft.	*	*	..	*	..	*		
<i>Lisianthus uliginosus</i> , Miq.	to Upper slopes.	*	*						
„ <i>Elisabethæ</i> , Griseb.	to Ledge.								
„ <i>Inthurnianus</i> , Oliver ..	Summit.								
„ <i>Quelchii</i> , N. E. Br.	Summit.								
SCROPHULARIACEÆ.									
<i>Melasma</i> ? <i>spathaceum</i> , Oliver	Upper slopes.								
<i>Beyrichia scutellarioides</i> , Benth.	to Upper slopes.	*	*	..	*				
LENTIBULARIACEÆ.									
<i>Utricularia Humboldtii</i> , Rob. Schomb.	to Ledge.								
<i>Utricularia nervosa</i> ?, Web.	Upper slopes.	*	*						
„ <i>Campbelliana</i> , Oliver ..	to Upper slopes.								
„ <i>alpina</i> , Jacq. (<i>U. montana</i> , Jacq.)	to Upper slopes.	..	..	*	..	*			
„ <i>Quelchii</i> , N. E. Br.	Summit.								
„ <i>roraimensis</i> , N. E. Br. ..	Summit.								
<i>Genlisea roraimensis</i> , N. E. Br.	Summit.								
BIGNONIACEÆ.									
<i>Tabebuia Roraimæ</i> , Oliver	to Upper slopes.								
VERBENACEÆ.									
<i>Lippia Schomburgkiana</i> , Schauer ..	to Upper slopes.	*							
LABIATÆ.									
<i>Hyptis lantanæfolia</i> , Poit.	to 5400 ft.	*	*	*	*	*	*		
POLYGONACEÆ.									
<i>Coccoloba Schomburgkii</i> , Meissn.	to 5400 ft.								
PIPERACEÆ.									
<i>Peperomia reflexa</i> , A. Dietr.	to Upper slopes.	*	*	*	*	*	*	..	*
„ <i>tenella</i> ?, A. Dietr.	to Upper slopes.	..	..	..	*	*			
„ sp.	at 5400 ft.								
CHLORANTHACEÆ.									
<i>Hedyosmum brasiliense</i> ?, Mart.	Upper slopes.	..	*						
PROTEACEÆ.									
<i>Roupala Schomburgkii</i> , Klotzsch ..	at 6000 ft.								
LORANTHACEÆ.									
<i>Loranthus</i> sp.	to 6000 ft.								
<i>Phoradendron Roraimæ</i> , Oliver	Ledge & Summit.								
EUPHORBIACEÆ.									
<i>Phyllanthus pycnophyllus</i> , Muell.-Arg.	at 5400 ft.								

† Apparently *C. auratum*, Miq.

		N. Brazil.	S. Brazil. Paraguay.	Colombia to Bolivia.	Venezuela and Trinidad.	W. Indies.	Central America.	Elsewhere in America.	Old World.
EUPHORBIACEÆ (cont.).									
<i>Phyllanthus</i> sp.									
<i>Croton lobatus</i> , Linn.	at 6000 ft.	*	*	*	*	*	*	..	Africa.
" sp.	Upper slopes.								
BURMANNIACEÆ.									
<i>Burmannia bicolor</i> , Mart.	to Upper slopes.	*	*	*	*	*	..	..	Africa.
<i>Dictyostegia orobanchoides</i> , Miers ..	to Upper slopes.	*	*	*	*				
ORCHIDACEÆ.									
<i>Pleurothallis stenopetala</i> , Lindl.	to Upper slopes.	..	*	..	*				
" <i>roraimensis</i> , Rolfe	Summit.								
<i>Stelis grandiflora</i> , Lindl.	Upper slopes.	*							
" <i>tristyla</i> , Lindl.	Upper slopes.	*							
<i>Lepanthes</i> sp.	at 5400 ft.								
<i>Masdevallia picturata</i> , Reichb. f. ..	to Upper slopes.	..	..	*	*	..	*		
<i>Scaphosepalum breve</i> , Rolfe	to Upper slopes.								
<i>Octomeria parviflora</i> , Rolfe	Summit.								
" <i>Connellii</i> , Rolfe	Summit.								
" sp.	Upper slope.								
<i>Bulbophyllum</i> sp. (<i>B. geraense</i> , Ridley, vix Reichb. f.) ..	to 5400 ft.								
" <i>roraimense</i> , Rolfe	Summit.								
<i>Elleanthus furfuraceus</i> , Reichb. f. ..	Upper slopes.	..	..	*	*				
<i>Epidendrum elongatum</i> , Jacq.	to Summit.	*	*	*	*	*			
" <i>alsum</i> , Ridley	Summit.								
" <i>Imthurii</i> , Ridley	Summit.								
" <i>tigrinum</i> , Lindl.	to Upper slopes.	..	..	*	*				
" <i>durum</i> , Lindl.	to 5400 ft.	..	*						
" <i>montigenum</i> , Ridley	Summit.								
" <i>violascens</i> , Ridley	Summit.								
<i>Cattleya Lawrenceana</i> , Reichb. f. ..	to 6000 ft.								
<i>Zygopetalum Burkei</i> , Reichb. f.	to Upper slopes.								
<i>Eriopsis Schomburgkii</i> , Reichb. f. ..	to Upper slopes.	*							
<i>Houlletia roraimensis</i> , Rolfe	to Upper slopes.								
<i>Catasetum discolor</i> , Lindl.	to Upper slopes.	*	*						
<i>Maxillaria Quelchii</i> , Rolfe	Summit.								
<i>Oncidium nigratum</i> , Lindl.	to 4500 ft.								
<i>Sobralia Liliastrum</i> , Lindl.	to 6000 ft.	*	*						
" sp.	Upper slope.								
<i>Epistephium lucidum</i> , Cogn.	to Upper slopes.	*	*						
<i>Stenoptera viscosa</i> , Reichb. f.	to Upper slopes.	..	..	*	*				
" <i>alnata</i> , Ridley	Upper slopes.								
<i>Spiranthes bifida</i> , Ridley	to 5400 ft.								
<i>Pogonia parviflora</i> , Lindl.	to Upper slopes.	..	*	..	..	..	..	S.	
" <i>rosea</i> , Hemsl.	to Upper slopes.	*	..	*	*				
" <i>tenuis</i> , Reichb. f.	Summit.	*	..	*	*				
<i>Habenaria roraimensis</i> , Rolfe	Summit.								
" <i>parviflora</i> , Lindl.	to 5400 ft.	*	*	*	*	..	..	S.	
" <i>Moritzii</i> , Ridley	to Upper slopes.	..	..	..	*				
<i>Selenipedium Lindleyanum</i> , Reichb. f.	to 5400 ft.								
BROMELIACEÆ.									
<i>Brocchinia cordylinoides</i> , Baker	to Ledge.								
<i>Puya floccosa</i> , Morren	at 6000 ft.	..	..	*	*				
<i>Connellia Augustæ</i> , N. E. Br.	to Summit.								
" <i>Quelchii</i> , N. E. Br.	Summit.								
<i>Tillandsia rhodocincta</i> , Baker	Summit.								

		N. Brazil.	S. Brazil, Paraguay.	Colombia to Bolivia.	Venezuela and Trinidad.	W. Indies.	Central America.	Elsewhere in America.	Old World.
LILIACEÆ.									
<i>Nietneria corymbosa</i> , Klotzsch & Rich. Schomb.	to Summit.								
<i>Tofieldia Schomburgkiana</i> , Oliver ..	to Summit.								
XYRIDACEÆ.									
<i>Xyris setigera</i> , Oliver	to Ledge.	..	*						
„ <i>Seubertii</i> , Nills.†	Upper slopes (? Ledge).								
„ <i>witserioides</i> , Oliver	to 7300 ft.								
„ <i>concinna</i> , N. E. Br.	Summit.								
<i>Abolboda Sceptrum</i> , Oliver	to Summit.								
RAPATEACEÆ.									
<i>Saxofredericia regalis</i> , Rob. Schomb.	to 5600 ft.								
<i>Stegolepis guianensis</i> , Klotzsch	to Summit.								
PALMÆ.									
<i>Euterpe</i> sp.	at 5400 ft.								
<i>Geonoma Appuniana</i> , Spruce	to Upper slopes.								
ARACEÆ.									
<i>Anthurium roraimense</i> , N. E. Br. ..	at 5400 ft.								
ERIOCAULACEÆ.									
<i>Peperanthus Schomburgkii</i> , Klotzsch	to 5400 ft.								
„ <i>flavescens</i> , Koern.	to Upper slopes.	*	*	*	*				
„ <i>Roraimæ</i> , Oliver	to Summit.								
„ <i>fraternus</i> , N. E. Br. ..	Summit.								
CYPERACEÆ.									
<i>Fimbristylis exilis</i> , Roem. & Schult...	to 5400 ft.	*	..	..	*	*	*	..	Africa.
<i>Rhynchospora stenophylla</i> , Britton ‡.	at 5400 ft.	..	..	..	..	*	..	N.	
„ <i>tenuis</i> , Britton	to 5400 ft.	*	*	..	.	*	*	S.	
<i>Lagenocarpus stellatus</i> , C. B. Clarke (<i>Cryptangium stellatum</i> , Boeckl.) .	Upper slopes.								
<i>Everardia montana</i> , Ridley	to Summit.								
„ <i>angusta</i> , N. E. Br.	Summit.								
<i>Scleria hirtella</i> , Sw.	to 5400 ft.	*	*	*	*	*	*	N.	Africa.
„ <i>bracteata</i> , Cav.	to 5400 ft.	*	*	*	*	..	*	..	*
GRAMINEÆ.									
<i>Saccharum cayennense</i> , Beauv.	to 5400 ft.	*	*	*	*	*	*	S.	
<i>Ischaemum latifolium</i> , Kunth	to 5400 ft.	*	*	*	*	*	*		
<i>Arundinella brasiliensis</i> , Raddi	to 5400 ft.	*	*	*	*	*	*	S.	*
<i>Paspalum stellatum</i> , Fluegge	to 5400 ft.	*	*	*	*				
<i>Panicum eligulatum</i> , N. E. Br.	Summit.								
„ <i>nervosum</i> ?, Lam.	to 5400 ft.	*	..	..	*				
<i>Echinolena hirta</i> , Desv.	to Upper slopes.	*	*						
<i>Arundo roraimensis</i> , N. E. Br.	Summit.								
<i>Arundinaria deflexa</i> , N. E. Br.	Summit.								
<i>Chusquea linearis</i> , N. E. Br.	Summit.								
<i>Guadua</i> sp.	to Upper slopes.								
Total of SPERMATOPHYTA		67	55	53	60	30	33	11	10

† *X. Fontanesiana*, Oliver in Trans. Linn. Soc., ser. II. Bot. ii. (1887), p. 285, appears to be *X. Seubertii*. It is not *X. Fontanesiana*, Kunth.

‡ *R. capillacea*, Oliver l. c. p. 287, not of Torrey (teste C. B. Clarke).

		N. Brazil.	S. Brazil, Paraguay.	Colombia to Bolivia.	Venezuela and Trinidad.	W. Indies.	Central America.	Elsewhere in America.	Old World.
FILICES (cont.).									
<i>Polypodium axanthotrichum</i> , Klotzsch	Upper slopes.								
„ <i>capillare</i> , Desv.	to Upper slopes.	*	..	*	*	*			
„ <i>Kalbreyeri</i> , Baker	Upper slopes.	..	..	*	*	*			
„ <i>rigescens</i> , Bory	to Summit.	*	*	*	*	*	..		*
„ <i>firmum</i> , Klotzsch	Upper slopes.	..	..	*	*	*			
„ <i>meridense</i> , Klotzsch ..	Summit.	..	*	*	*	*			
„ <i>subsessile</i> , Baker	Upper slopes.	..	..	*	*	*			
„ <i>melanotrichum</i> , Baker ..	at 5400 ft.							S.	
„ <i>loriceum</i> , Linn.	to 7200 ft.	*	*	*	*	*	*	N.	
„ <i>aweum</i> , Linn.	to 5400 ft.	*	*	*	*	*	*		Australia.
„ <i>angustifolium</i> , Sw.	to 5400 ft.	*	*	*	*	*	*		
<i>Gymnogramme daphnoglossoides</i> , Baker	to Summit.								
„ <i>cyclophylla</i> , Baker ..	Summit.								
„ <i>flexuosa</i> , Desv.	to Summit.	*	*	*	*	..	*		
„ <i>Schomburgkiana</i> , Kunze	Upper slopes.								
„ <i>hirta</i> , Desv.	Upper slopes.	..	..	*	*	..	*		
<i>Enteromorpha Campbellianum</i> , Baker ..	Summit.								
<i>Vittaria lineata</i> , Sw.	to Upper slopes.	*	*	*	*	*	*	N. & S.	*
„ <i>stipulata</i> , Kunze	Upper slopes.	..	..	*	*	*	*		
<i>Acrostichum latifolium</i> , Sw.	to Summit.	*	*	*	*	*	*	..	*
„ <i>squamosum</i> , Sw.	to Upper slopes.	*	*	*	*	*	*	..	*
„ <i>leptophlebium</i> , Baker ..	Upper slopes.								
„ <i>muscosum</i> , Sw.	to 5400 ft.	*	*	*	..	*	*		
„ <i>stenopteris</i> , Klotzsch ..	at 5400 ft.	..	..	*	*	*	*		
„ <i>decoratum</i> , Kunze	to 5400 ft.	..	*	*	..	*	*		
„ <i>peltatum</i> , Sw.	to Upper slopes.	*	..	*	*	*	*		
<i>Schizera dichotoma</i> , Sw.	to 5400 ft.	..	..	*	*	*	..	..	*
„ <i>elegans</i> , Sw.	to 5400 ft.	*	*	*	*	*	*		
<i>Aneimia tomentosa</i> , Sw.	to 5400 ft.	*	*	*	*	*	*	S.	
Total of PTERIDOPHYTES		55	50	67	59	55	45	17	25
BRYINEÆ.									
<i>Dicranum longisetum</i> , Hook.	Ledge.	..	..	*	*				
<i>Dicranodontium pulchroalare</i> , Broth.	Ledge.								
<i>Campylopus chionophyllus</i> , Mitt. ..	Ledge.	..	..	*					
„ <i>Roraimae</i> , Broth.	Ledge.								
„ <i>atratus</i> , Broth.	Summit.								
<i>Leucobryum megalophyllum</i> , Mitt. ..	Ledge.	..	*	*	*	*	..	*	
„ <i>levifolium</i> , Broth.	Summit.								
<i>Octoblepharum Mittenii</i> , Jaeg.	Summit.	*	*	..	*				
<i>Leptodontium cirrhifolium</i> , Mitt.	Summit.	..	..	*					
<i>Zygodon subdenticulatus</i> , Hampe	Summit.	..	..	*					
<i>Funaria calvescens</i> , Schwaeg.	Ledge.	*	*	*	*	*	*	N. & S.	*
<i>Breutelia scoparia</i> , Besch.	Summit.	..	..	..	..	*			
<i>Rhizogonium Lindigii</i> , Mitt.	Summit.	..	..	*	..	..	*		
<i>Mnium rostratum</i> , Schrad.	to Ledge.	*	*	*	*	*	*	*	*
<i>Polytrichum aristiflorum</i> , Mitt.	to 5400 ft.	..	*	*	*	*	*		
<i>Rhaecocarpus Humboldtii</i> , Lindb.	Summit.	..	*	*	..	*	..	S.	*
<i>Hookeria pilotrichelloides</i> , Broth. ..	Ledge.								
„ <i>crispa</i> , C. Muell.	to 5400 ft.	..	*	*	*	..	*		
<i>Ectropothecium amabile</i> , Mitt.	to Summit.	..	..	*					
<i>Ctenidium malacodes</i> , Mitt.	Ledge.	..	..	*					
<i>Thuidium pseudoprotensum</i> , Mitt.	Ledge.	..	..	..	*				
<i>Hypopterygium tamarisci</i> , Brid.	to 5400 ft.	*	*	*	*	*	*	S.	*
SPHAGNACEÆ.									
<i>Sphagnum sanguinale</i> , Warnst.	Ledge.								
„ <i>medium</i> , Limpr.	Ledge.	..	*	..	..	..	..	N. & S.	*
MARCHANTIACEÆ.									
<i>Dumortiera hirsuta</i> , Nees	Ledge.	..	*	..	..	*	*	..	*

		N. Brazil.	S. Brazil, Paraguay.	Colombia to Bolivia.	Venezuela and Trinidad.	W. Indies.	Central America.	Elsewhere in America.	Old World.
JUNGERMANNIACEÆ.									
<i>Aneura Schwaneckei</i> , Steph.	Summit.	..	*	..	..	*			
" <i>algoides</i> , Steph.	Summit.	..	..	*					
" <i>roraimensis</i> , Steph.	Summit.	..	..	..	..	*			
" <i>Breutelii</i> , Steph.	Summit.	..	..	..	..	*			
" <i>fucoides</i> , Steph. (<i>A. bipinnata</i> , Mitt.†)	Ledge.	..	*	..	..	*	*		
<i>Metzgeria inflata</i> , Steph.	Summit.	..	..	..	..	*			
" <i>hamata</i> , Lindb.	Ledge.	..	..	*	..	*	*	N. & S.	*
<i>Pallavicinius Wallisii</i> , Jack & Steph.	Summit.	..	..	*	..	..	*	S.	*
<i>Jamesoniella colorata</i> , Spruce	Summit.	..	..	*	..	..	*		
<i>Syzygiella Quelchii</i> , Steph.	Ledge.	..	..	..	..	*			
" <i>perfoliata</i> , Spruce	Summit.	..	..	..	*	*			
<i>Plagiocbila cerea</i> , Tayl.	Ledge.	..	..	..	*	..	*		
" <i>guyana</i> , Steph.	Ledge.	..	..	..	..	..			
" <i>remotifolia</i> , Hampe & Gottsche	Ledge.	..	..	..	..	*			
" <i>rutilans</i> , Lindb.	Ledge.	*	*	*	*	*	*		
" <i>adiantoides</i> , Lindb.	to Upper slopes.	..	..	*	..	*	*		
<i>Leioscyphus fragilis</i> , Jack & Steph. ..	Summit.	..	..	*	..	..			
<i>Lophocolea Breutelii</i> , Gottsche	Ledge.	..	..	..	..	*			
<i>Mastigobryum roraimense</i> , Steph. ..	Summit.	..	..	..	..	..			
" <i>dissodontum</i> , Spruce ..	Summit.	..	..	*	..	..			
" <i>vincentinum</i> , Lehm. & Lindenb.	Ledge.	..	..	..	..	*			
" <i>Krugianum</i> , Steph. ..	Ledge.	..	..	..	..	*			
" <i>gracile</i> , Hampe & Gottsche	Ledge.	..	..	..	..	*			
<i>Blepharozia Roraime</i> , Mitt.	Summit.	..	..	..	..	..			
<i>Micropterygium grandistipulum</i> , Steph.	Ledge.	..	..	..	..	..			
" <i>pterygophyllum</i> , Spruce	Summit.	*	*	*	..	*			
<i>Lepidozia commutata</i> , Steph.	Summit.	..	..	..	..	*			
<i>Schisma juniperinum</i> , Nees	Summit.	..	..	*	..	*	*	N. & S.	*
" <i>pensilis</i> , Steph.	Ledge.	..	..	*	..	..			
" <i>Durandii</i> , Steph.	Ledge.	..	..	..	..	..	*		
" <i>subdentatum</i> , Steph.	Ledge.	..	..	..	..	..			
<i>Trichocolea sphagnoides</i> , Steph.	Ledge.	..	..	*	..	..			
<i>Scapania portoricensis</i> , Hampe & Gottsche	Summit.	..	..	*	..	*			
<i>Pleurozia paradoxa</i> , Jack ..	Summit.	..	..	*	..	..			
<i>Harpaljeunea tenax</i> , Steph.	Ledge.	..	..	..	..	..			
<i>Frullanea mirabilis</i> , Jack & Steph. ..	Summit.	..	..	*	..	..			
" <i>atrata</i> , Nees	Summit.	*	*	*	*	*	*		
" <i>longicollis</i> , Lindenb. & Gottsche	Summit.	..	..	*	..	..	*		
Total of BRYOPHYTA		7	17	31	13	25	17	8	9
<i>Phoma Psammisia</i> , Massée	Summit.	..	..	..	..	..	..		*
<i>Echinobotryum roseum</i> , Massée	Summit.	..	..	..	..	..	..		*
<i>Stenophyllum ericoctonum</i> , R. Br. ..	Summit.	..	..	..	..	..	..		*
<i>Macrosporium ramulosum</i> , Sacc.	Summit.	..	..	*	..	..	..		*
<i>Capnodium fibrosum</i> , Berk.	Summit.	..	..	..	..	..	..		*
<i>Rhipidonema membranaceum</i> , Sacc.	Ledge.	*	..	..	..	..	..		Pacific.
<i>Sphaerophora compressum</i> , Achar. ..	Summit.	..	..	..	..	*	..	S.	*
<i>Cladonia rangiferina</i> , Hoffm.	Summit.	*	*	*	*	*	*	N. & S.	*
<i>Parmelia dictyosporia</i> , Massée	Summit.	..	..	..	..	..	..		
" <i>perforata</i> , Achar.	Summit.	..	..	..	..	..	*	N.	Pacific.
<i>Alectoria ochroleuca</i> , Nyl.	Summit.	..	..	..	..	..	*	N.	*
Total of THALLOPHYTA		2	1	2	1	2	3	4	8

CHIEF LITERATURE.

Schomburgk's journeys.

- Sir Robert H. Schomburgk, in *Proc. Roy. Geogr. Soc.* x. 1840, pp. 196-242.
 Sir Robert H. Schomburgk, 'Twelve Views in the Interior of Guiana' (London, 1841).
 Sir Richard Schomburgk, 'Reisen in Britisch-Guiana' (Leipzig, 1847-48), ii. pp. 152-300;
 iii. pp. 1041-1104.
 Sir Richard Schomburgk, 'Botanical Reminiscences of British Guiana' (Adelaide, 1876).
 Sir Robert H. Schomburgk, in *Verh. Gartenb. Ver.* xv., 1841.
 G. Benthams, in *Ann. Nat. Hist.* ii.-iii.; in *Hooker's Journ. Bot.* ii.-iv.; in *Hooker's Lond. Journ. Bot.* i.-vii.; in *Trans. Linn. Soc.* xviii. & xxii.; &c.
 J. F. Klotzsch, in *Verh. Gartenb. Ver.* xviii., 1847; in 'Linnæa,' xxiv. 1851, p. 1; &c.
 Various authors in Martius, 'Flora Brasiliensis' (Leipzig, 1840-1900).

Appun's journey.

- K. Appun, 'Unter den Tropen' (Jena, 1871), ii. pp. 105-384.

Brown and Sawkins's journey.

- C. B. Brown, 'Canoe and Camp-life, in Guiana,' 2nd edit. (London, 1877), pp. 118-124.
 J. G. Sawkins, in *Proc. Roy. Geogr. Soc.* xv. 1870-71, p. 131.
 C. B. Brown and J. G. Sawkins, 'Report on the Physical, Descriptive and Economic Geology of British Guiana' (London, 1875), pp. 61-63.

Boddam-Whetham and McTurk's journey.

- J. W. Boddam-Whetham, 'Roraima and British Guiana' (London, 1879), pp. 204-254.

im Thurn and Perkins's journey.

- E. F. im Thurn, in *Proc. Roy. Geogr. Soc.* vii. 1885, pp. 497-521, and in 'Timehri,' iv. 1885, pp. 1-48 and 256-267.
 H. I. Perkins, in *Proc. Roy. Geogr. Soc.* vii. 1885, pp. 529-530.
 E. F. im Thurn, D. Oliver, and others in *Trans. Linn. Soc. Ser. II. Bot.* in. pp. 249-300; reprinted in 'Timehri,' v. 1886, pp. 145-223.

Kromer's and Dressel's journeys.

- 'Timehri,' n. s., i. 1887, p. 330.

McConnell and Quelch's first journey.

- J. J. Quelch in 'Timehri,' n. s., viii. 1894, p. 381; ix. 1895, pp. 107-188.

ENUMERATION OF THE PLANTS COLLECTED.

I. **SPERMATOPHYTA.**

By N. E. BROWN, A.L.S. (the Orchids by R. A. ROLFE, A.L.S.).

SARRACENIACEÆ.

HELIAMPHORA NUTANS, Benth. in Proc. Linn. Soc. i. (1840), p. 53, and in Trans. Linn. Soc. xviii. (1840), p. 429, t. 29; R. Schomb. Reisen in Brit.-Guiana, ii. p. 263, and iii. p. 1090; R. H. Schomb., Views in the Interior of Guiana, p. 15; Bot. Mag. t. 7093; Fl. des Serres, Sér. II. xi. (1875), p. 149, tt. 2246-47; Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 271.

Summit of Mount Roraima, *McConnell & Quelch*, 78, 679; also collected on the lower slope in the "Eldorado Swamp" by all other collectors.—Endemic.

VIOLACEÆ.

IONIDIUM IPECACUANHA, Vent. Jard. Malm. sub t. 27.

Ireng Valley, *McConnell & Quelch*, 258.

ALSODEIA FLAVESCENS, Spreng. Syst. i. p. 806.

Mazaruni River, 300 ft., *McConnell & Quelch*, 715.

SAUVAGESIA ERECTA, Linn. Sp. Pl. ed. i. p. 203; Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 271.

Upper slopes and ledge of Mount Roraima, *McConnell & Quelch*, 7. Ireng Valley, *McConnell & Quelch*, 263.—Widely distributed in Tropical America.

SAUVAGESIA SPRENGELII, A. St. Hil. in Mém. Mus. Par. xi. (1824), p. 97.

Savannahs generally, *McConnell & Quelch*, 330.

LEITGEBIA IMTHURNIANA, Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 271, t. 37A.

Summit of Mount Roraima, *McConnell & Quelch*, 97, 98, 316, 654.—Endemic.

POLYGALACEÆ.

POLYGALA GLOCHIDIATA, H. B. & K. Nov. Gen. & Sp. v. p. 400.

Upper slopes of Mount Roraima, *McConnell & Quelch*, 38.—Widely distributed in Tropical America.

POLYGALA HYGROPHILA, H. B. & K. Nov. Gen. & Sp. v. p. 395; Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 271.

Upper slopes of Mount Roraima, *McConnell & Quelch*, 39.—Eastern Tropical America from Panama to Paraguay.

POLYGALA TIMOUTOU, Aubl. Pl. Guian. ii. p. 737, t. 295; R. Schomb. Reisen in Brit.-Guiana, iii. p. 1097.

Kotinga Valley, *McConnell & Quelch*, 196. Ireng Valley, *McConnell & Quelch*, 250.

POLYGALA CELOSIODES, Mart. *ex* A. W. Benn. in Mart. Fl. Bras. xiii. pt. III. p. 35.

Ireng Valley, *McConnell & Quelch*, 223.

SECURIDACA MARGINATA, Benth. in Hook. Journ. Bot. iv. (1842), p. 103.

Kotinga Valley, *McConnell & Quelch*, 157.

MONNINA CACUMINA, N. E. Brown, sp. n. Folia oblonga, acuta vel obtusa, apiculata, exstipulata, glabra. Racemi densiflori, floribus breviter pedicellatis. Bracteae lanceolatae, acutae, floribus breviores. Alae ellipticae, glabrae, basi ciliatae. Carina glabra, ciliata. Petala lateralibus utrinque pubescentia. Vagina staminea supra hirta. Fructus compresso-ellipsoideus, exalatus, unilocularis, glaber.

Stem subglabrous, or very thinly covered with very short hairs. Leaves about 2 in. long, $\frac{2}{3}$ -1 in. broad, oblong or oblong-lanceolate, acute or obtuse and apiculate, rather thin in the dried state, exstipulate. Racemes 3-4, terminal, 2-4 in. long, rather densely many-flowered. Bracts 2 lin. long, $\frac{2}{3}$ lin. broad, lanceolate, acute, caducous. Pedicels $\frac{1}{2}$ -1 lin. long. Dorsal sepal $1\frac{1}{3}$ lin. long, $\frac{2}{3}$ lin. broad, ovate, acute, concave, ciliolate; the two lower sepals similar but rather smaller, connate to their middle; lateral sepals or alae about $2\frac{1}{4}$ lin. long, 2 lin. broad, elliptic, obtuse, with incurved margins, glabrous, ciliate towards the base. Lower petal or keel $2\frac{1}{2}$ lin. long, 2 lin. broad, obscurely 3-crenulate at the apex, glabrous ciliate along the sides. Lateral petals 2- $2\frac{1}{2}$ lin. long, obliquely tapering from a broad base to an obtuse apex, adnate to the staminal tube, densely ciliate along the upper margin, pubescent along the middle of the outer side and more densely all over within. Staminal tube villose at the apex. Ovary elliptic, with a large unilateral gland at its base, glabrous; style thickened upwards, slightly incurved. Fruit 3- $3\frac{1}{2}$ lin. long, about $1\frac{3}{4}$ lin. broad, elliptic, subacute, compressed, not winged, 1-celled, glabrous.

Summit of Mount Roraima, 8600 ft. *McConnell & Quelch*, 645.

Allied to *M. cestrifolia*, H. B. & K., but differing in its much larger flowers.

KRAMERIA SPARTIOIDES, Klotzsch, *ex* O. Berg, in Bot. Zeit. xiv. (1856), p. 761.

Ireng Valley, *McConnell & Quelch*, 244, 268.

VOCHYSIACEÆ.

TRIGONIA SUBCYMOSA, Benth. in Hook. Lond. Journ. Bot. ii. (1843), p. 373.

Ireng Valley, *McConnell & Quelch*, 222.

PORTULACACEÆ.

PORTULACA SEDIFOLIA, N. E. Brown, sp. n. Caules humiles, diffusi. Folia parva, lineari-oblonga vel oblongo-lanceolata, breviter petiolata, axillis dense pilosa. Calycis lobi

late ovati, obtusi, apice inflexi, glabri. Petala calyce duplo longiora. Stamina 5. Stylus 3-fidus. Semina tuberculata.

A small succulent herb, branching at the base, with branches 2-3 in. long, glabrous, with the exception of the tufts of long white hairs in the axils of the leaves and around the flowers. Leaves $1\frac{1}{2}$ -3 lin. long, $\frac{1}{2}$ - $\frac{2}{3}$ lin. broad, flattened, linear-oblong or oblong-lanceolate, obtuse, tapering at the base into a short petiole, glabrous. Flowers small. Sepals $\frac{3}{4}$ lin. long, rather more than $\frac{1}{2}$ lin. broad, broadly ovate, obtuse, inflexed at the apex, concave, not keeled on the back, glabrous. Petals $1\frac{1}{2}$ lin. long, only seen in a withered state, apparently obtuse. Stamens 5, shorter than the petals. Style filiform, with 3 filiform stigmas. Upper half of the capsule broadly conical. Seeds $\frac{1}{4}$ lin. diam., tuberculate, the tubercles on the sides very much larger than those on the back, black.

Ireng Valley, *McConnell & Quelch*, 237.

Allied to *Portulaca parvula*, A. Gray, but differing in its shorter leaves, fewer stamens, and larger and more coarsely tuberculate seeds.

GUTTIFERÆ.

CLUSIA PLANCHONIANA, Engl. (?) in Mart. Fl. Bras. xii. pt. i. p. 431.

Kukenaam River, *McConnell & Quelch*, 130.

I am doubtful of my identification of this plant, and it may prove to be a new species, but the specimen only consists of three flowers and two leaves, all detached. They agree fairly well with the flowers and leaves of *C. Planchoniana*, with the exception that the leaves are rather more obtuse and have stouter petioles.

TERNSTRÆMIACEÆ.

TERNSTRÆMIA SCHOMBURGKIANA, Benth. in Hook. Lond. Journ. Bot. ii. (1843), p. 362;

Appun, Unter den Tropen, ii. p. 292.

Kotinga Valley, *McConnell & Quelch*, 138.

BONNETIA SESSILIS, Benth. in Hook. Lond. Journ. Bot. ii. (1843), p. 363; Appun, Unter den Tropen, ii. p. 292; R. Schomb. Reisen in Brit.-Guiana, ii. pp. 218, 251, and iii. p. 1093.

Irengwatong Creek in the Ireng Valley, *McConnell & Quelch*, 269, 322.—Only known from this western part of British Guiana.

BONNETIA RORAIMÆ, Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 272, t. 37. f. B.

Summit of Mount Roraima, *McConnell & Quelch*, 84, 664.—Endemic. According to Mr. Quelch, "this is the commonest and most widely spread species on the summit, rising to 35-40 ft. in height, where sheltered."

ARCHYTÆA MULTIFLORA, Benth. in Hook. Lond. Journ. Bot. ii. (1843), p. 363; R. Schomb.

Reisen in Brit.-Guiana, ii. p. 218, and iii. p. 1093.

Kotinga Valley, 3000 ft. *McConnell & Quelch*, 162; Ireng Valley, *McConnell & Quelch*, 311.

MAHURIA EXSTIPULATA, Benth. in Hook. Lond. Journ. Bot. ii. (1843), p. 365.
Silima Creek in the Ireng Valley, *McConnell & Quelch*, 270.

MALVACEÆ.

SIDA, sp. Specimen insufficient for determination, but apparently a new species.
Ireng Valley, *McConnell & Quelch*, 267.

PAVONIA SPECIOSA, H. B. & K. Nov. Gen. & Sp. v. p. 281.

Kotinga Valley, *McConnell & Quelch*, 176. Karona Falls on the Ireng River,
McConnell & Quelch, 228. Ireng Valley, *McConnell & Quelch*, 257.

PAVONIA CANCELLATA, Cav. Diss. iii. p. 135.

Ireng Valley, *McConnell & Quelch*, 248.

FUGOSIA CAMPESTRIS, Benth. in Hook. Journ. Bot. iv. (1842), p. 120.

Ireng Valley, *McConnell & Quelch*, 213, 239.

STERCULIACEÆ.

HELICTERES GUAZUMÆFOLIA, H. B. & K. Nov. Gen. & Sp. v. p. 304.

Ireng Valley, *McConnell & Quelch*, 212.

MELOCHIA MELISSÆFOLIA, Benth. in Hook. Journ. Bot. iv. (1842), p. 129.

Ireng Valley, *McConnell & Quelch*, 231.

MELOCHIA HIRSUTA, Cav. Diss. vi. p. 323.

Ireng Valley, *McConnell & Quelch*, 255.

WALTHERIA VISCOSISSIMA, A. St. Hil. Fl. Bras. Mer. i. p. 150.

Kotinga Valley, *McConnell & Quelch*, 168.

MALPIGHIACEÆ.

BYRSONIMA VERBASCIFOLIA, Rich. *ex* Juss. in Ann. Mus. Par. xviii. (1811), p. 481.

Ireng Valley, *McConnell & Quelch*, 226.

HETEROPTERIS OLEÆFOLIA, Griseb. in Linnæa, xxii. (1849), p. 19.

H. daphnoides, Griseb. in R. Schomb. Reisen in Brit.-Guiana, iii. p. 1096.

Upper slopes of Mount Roraima, *McConnell & Quelch*, 21, 319.—Endemic.

TETRAPTERIS SQUARROSA, Griseb. in Mart. Fl. Bras. xii. pt. i. p. 87.

Ireng Valley, *McConnell & Quelch*, 241.

GERANIACEÆ.

OXALIS DISTANS, A. St. Hil. Fl. Bras. Mer. i. p. 115.

O. Schomburgkiana, Prog. in Mart. Fl. Bras. xii. pt. ii. p. 500.

Ireng Valley, *McConnell & Quelch*, 202.

OCHNACEÆ.

PÆCILANDRA RETUSA, Tul. in Ann. Sc. Nat. Sér. III. viii. (1847), p. 342.

Kotinga Valley, *McConnell & Quelch*, 145, 167. Ireng Valley, *McConnell & Quelch*, 325.

ILICACEÆ.

ILEX APICIDENS, N. E. Brown, sp. n. (Plate 1, figs. 1-6). Frutex glaber. Folia parva, coriacea, elliptica, obtusa vel emarginata, basi obtusa, apice denticulata. Cymæ axillares vel ad apices ramulorum aggregatæ, 2-3-floræ, vel flores solitarii, glabræ. Calyx 4-lobus, lobis brevissimis rotundatis. Petala 4, elliptica vel suborbiculata, obtusissima. Stamina 4.

A glabrous shrub; branches tuberculated with the scars of the fallen leaves. Leaves moderately crowded, coriaceous; petiole $\frac{1}{2}$ -2 lin. long; blade 5-11 lin. long, 4-7 lin. broad, elliptic or somewhat obovate, obtusely rounded or emarginate and with 5-9 small teeth at the apex, obtusely rounded or somewhat cuneate at the base, entire along the sides, with slightly revolute margins, not punctate-dotted beneath; primary lateral veins about 4 on each side, uniting in broad loops, impressed above, prominent beneath; stipules minute, about $\frac{1}{3}$ lin. long, subulate. Flowers either solitary or in 2-3-flowered pedunculate cymes in the axils of the leaves, or crowded into a small, terminal, hemispherical corymb $\frac{1}{2}$ - $\frac{3}{4}$ in. diam. Peduncles $1\frac{1}{2}$ -3 lin. long, glabrous. Pedicels of the solitary flowers 2-3 $\frac{1}{2}$ lin. long, of those in cymes 1-2 $\frac{1}{2}$ lin. long, glabrous. Calyx about $\frac{1}{2}$ lin. long, cupular, shortly 4-lobed, lobes broadly rounded, obtuse, glabrous. Petals 4, $1\frac{1}{4}$ - $1\frac{1}{2}$ lin. long, $1\frac{1}{4}$ - $1\frac{1}{3}$ lin. broad, elliptic or suborbiculate, very obtuse, concave, glabrous. Stamens 4, a little shorter than the petals. Pistillode globose, glabrous. Female flowers not seen.

Summit of Mount Roraima, 8600 ft., *McConnell & Quelch*, 634.

Allied to *I. obcordata*, Sw., but differing in the apically toothed leaves, with the midrib and veins much more prominent beneath.

ILEX RETUSA, Klotzsch, in R. Schomb. Reisen in Brit.-Guiana, iii. p. 1097; Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 273.

Summit of Mount Roraima, *McConnell & Quelch*, 655. Collected by Schomburgk on the bank of a mountain-stream in the vicinity of Roraima.—Endemic.

CYRILLACEÆ.

CYRILLA ANTILLANA, Michx. Fl. Bor.-Am. i. p. 158.

Stachyanthemum Schomburgkii, Klotzsch, in R. Schomb. Reisen in Brit.-Guiana, iii. p. 1097, ex Loesener, in Engler & Prantl, Pflanzenfam. iii. pt. 5, p. 459.

Kotinga Valley, 3000 ft., *McConnell & Quelch*, 193. Ireng Valley, *McConnell & Quelch*, 203.

CYRILLA BREVIFOLIA, N. E. Brown, sp. n. (Pl. 1, figs. 7-16.) Folia lanceolato-elliptica vel obovato-elliptica vel lanceolata, obtusa vel acuta, glabra, rigide coriacea, costa infra vix prominente. Racemi 1-2 poll. longi, dense multiflori. Sepala lanceolata, acuminata, glabra. Petala lanceolata, involuto-acuminata, glabra.—*C. antillana*, Michx., var. *brevifolia*, Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887) p. 273.

Leaves crowded towards the ends of the branches, glabrous; petiole 2-2½ lin. long; blade $\frac{3}{4}$ -1¼ in. long, 4-8 lin. broad, lanceolate-elliptic or slightly obovate-elliptic and obtuse, or occasionally lanceolate and acute, sometimes minutely apiculate, rigidly coriaceous, entire, shining and smooth or coarsely pitted above, coarsely pitted between the veinlets beneath; midrib impressed above, prominent and keeled beneath at the basal part only, not raised above the general surface in the apical half. Racemes 1-2 in. long, 3-4 lin. diam., densely many-flowered, erect, glabrous. Bracts 1½-2 lin. long, subulate, very acute. Pedicels $\frac{3}{4}$ -1 lin. long, ascending curved, glabrous. Sepals 1-1¼ lin. long, acuminate from an ovate base, glabrous. Petals 1½-1¾ lin. long, $\frac{1}{2}$ lin. broad, lanceolate, acuminate from the margins being inrolled, subacute when expanded, glabrous. Stamens about $\frac{2}{3}$ as long as the petals. Ovary oblong, irregularly wrinkled in the dried state, 2-3-celled, glabrous; style very short and stout; stigmas 2-3. Ovules 2, pendulous from the apex of each cell.

Summit of Mount Roraima, 8600 ft., *in Thurn*, 334; *McConnell & Quelch*, 88, 318, 638.

This species differs from *Cyrilla antillana*, Michx., in having smaller and more rigidly coriaceous leaves, with the midrib not prominent beyond the middle on the underside, shorter and denser racemes, and longer and more acuminate sepals and petals.

LEGUMINOSÆ.

CROTALARIA ANAGYROIDES, H. B. & K. Nov. Gen. & Sp. vi. p. 404.

Kotinga Valley, *McConnell & Quelch*, 169.

TEPHROSIA ADUNCA, Benth. in Ann. Nat. Hist. Ser. I. iii. (1839), p. 432.

Kotinga Valley, *McConnell & Quelch*, 131, 199.

CLITORIA GUYANENSIS, Benth. in Journ. Linn. Soc., Bot. ii. (1858), p. 40.

Kotinga Valley, *McConnell & Quelch*, 155; Ireng Valley, *McConnell & Quelch*, 209.

DIOCLEA GUIANENSIS, Benth. in Ann. Wien. Mus. ii. (1838), p. 134.

Kotinga Valley, *McConnell & Quelch*, 158.

PHASEOLUS SEMIERECTUS, Linn. Mant. i. p. 100, var.

Kotinga Valley, *McConnell & Quelch*, 171.

PHASEOLUS LASIOCARPUS, Mart. (?) *ex* Benth. in Ann. Wien. Mus. ii. (1838), p. 140,

Kotinga Valley, *McConnell & Quelch*, 177.

ERIOSEMA CRINITUM, Benth. in Hook. Journ. Bot. ii. (1840), p. 62.

Kotinga Valley, *McConnell & Quelch*, 186.

DIPTERYX OPPOSITIFOLIA, Willd. Sp. Pl. iii. p. 910.

Arabapu River, *McConnell & Quelch*, 143, 303. Roraima Range, 3500 ft., *McConnell & Quelch*, 717.—Guiana and Brazil.

CASSIA INSIGNIS, N. E. Brown, sp. n. Foliola bijuga, ampla, oblique oblonga vel elliptico-oblonga, breviter obtuse acuminata, venosa, glandula magna inter inferiora. Stipulæ foliaceæ, falcato-lanceolatae. Racemi laxi subcorymbosi. Pedicelli elongati. Stamina perfecta 7; staminodia 3. Antheræ perfectæ subæquales, 3 inferiores breviter rostratae.

Leaves large, bijugate; petiole below the basal pair of leaflets, $\frac{3}{4}$ –1 in. long, minutely puberulous, becoming nearly glabrous; leaflets obliquely oblong or elliptic-oblong, shortly and obtusely acuminate, obliquely rounded at the base, glabrous to the eye, but with a minute scattered pubescence as seen under a lens, with the veins and veinlets very conspicuous on both sides; the basal pair $1\frac{1}{2}$ – $3\frac{1}{4}$ in. long, $\frac{3}{4}$ – $1\frac{3}{4}$ in. broad; the terminal pair 4–9 lin. distant, $2\frac{3}{4}$ – $4\frac{1}{2}$ in. long, 1–2 in. broad; gland $1\frac{1}{2}$ lin. long, 1 lin. thick, erect, ovoid, obtuse, placed between the basal pair of leaflets. Stipules leafy, $\frac{1}{2}$ – $\frac{3}{4}$ in. long, 2–4 lin. broad, falcate-lanceolate, obtuse or acute, tapering at the base into a curved petiole. Racemes loosely subcorymbose at the ends of the branches, solitary or in pairs from the axils of the uppermost reduced leaves, $1\frac{1}{2}$ –3 in. long, 7–14-flowered, softly and minutely pubescent. Pedicels 1– $1\frac{3}{4}$ in. long, pubescent. Sepals about $5\frac{1}{2}$ lin. long, $\frac{1}{4}$ in. broad, obovate, obtuse, puberulous, minutely ciliate towards the apex. Petals 8–11 lin. long, 4–8 lin. broad, elliptic or elliptic-obovate, very distinctly clawed, minutely puberulous on both sides. Stamens with 7 subequal perfect and 3 reduced abortive anthers, 3 of the perfect anthers shortly beaked. Ovary linear, curved, densely pubescent with yellowish adpressed hairs. Legume not seen.

On the upper slopes of Mount Roraima, *McConnell & Quelch*, 23.

Allied to *C. latifolia*, G. F. W. Mey., but may be at once distinguished by its narrow-based, petiolate stipules, and much longer pedicels.

CASSIA APOUCOUITA, Aubl. Pl. Guian. i. p. 379, t. 146.

Mazaruni River, 300 ft., *McConnell & Quelch*, 716.

This appears to be the true plant of Aublet, which is easily distinguished from the Brazilian plants described by Benth. as *C. Apoucouita*, by its narrowly winged petiole and differently shaped leaflets. The *C. Apoucouita* of Benth. appears to me to include 2 or 3 perfectly distinct species.

CASSIA HISPIDULA, Vahl, Eclog. Am. iii. p. 10.

Kotinga Valley, *McConnell & Quelch*, 170. Ireng Valley, *McConnell & Quelch*, 246.

CASSIA POLYSTACHYA, Benth. in Hook. Journ. Bot. ii. (1840), p. 77; R. Schomb. Reisen in Brit.-Guiana, ii. pp. 187, 207, and iii. p. 1207.

Ireng Valley, *McConnell & Quelch*, 229. Upper slopes of Mount Roraima, *McConnell & Quelch*, 317.—Only known from this western corner of British Guiana.

CASSIA UNIFLORA, Spreng. Neue Entdeck. i. p. 291.

Kotinga Valley, near Roraima, *McConnell & Quelch*, 133. Kotinga Valley, *McConnell & Quelch*, 151. Savannahs generally, *McConnell & Quelch*, 320.

CASSIA RORAIMÆ, Benth. in Trans. Linn. Soc. xxvii. (1871), p. 574; Oliver in Trans. Linn. Soc. Ser. II. ii. (1887) p. 273.

Kotinga Valley, *McConnell & Quelch*, 140. Roraima Range, 3500 ft., *McConnell & Quelch*, 719.—Only known from this region.

MIMOSA MICROCEPHALA, Humb. & Bonpl. *ex* Willd. Sp. Pl. iv. p. 1039.

Kotinga Valley, *McConnell & Quelch*, 173.

PITHECOLOBIUM FERRUGINEUM, Benth. in Hook. Lond. Journ. Bot. iii. (1844), p. 208; R. Schomb. Reisen in Brit.-Guiana, iii. p. 1038.

Upper and lower slopes of Mount Roraima, *McConnell & Quelch*, 19. Roraima Range, 3500 ft., *McConnell & Quelch*, 712.—Endemic.

The fruit described by Bentham under *P. ferrugineum* in Trans. Linn. Soc. xxx. (1875) p. 584, does not belong to that species, but to some totally different plant.

ROSACEÆ.

MOQUILEA TURIUVA, Hook. f. in Mart. Fl. Bras. xiv. pt. II. p. 25.

Savannahs generally, *McConnell & Quelch*, 308.

HIRTELLA AMERICANA, Linn. Sp. Pl. ed. 1, p. 34.

Savannahs generally, *McConnell & Quelch*, 309.

RUBUS URTICÆFOLIUS, Poir. Encycl. vi. p. 246.

Upper slopes of Mount Roraima, *McConnell & Quelch*, 11.—Widely dispersed throughout Tropical America.

It is uncertain whether *R. Schomburgkii*, Klotzsch, in R. Schomb. Reisen in Brit.-Guiana, iii. p. 1102, is a synonym of this species or identical with *R. guyanensis*, Focke, in Brem. Abh. iv. (1874), p. 160, as no description is given by Klotzsch, and Schomburgk collected both species.

SAXIFRAGACEÆ.

WEINMANNIA GUYANENSIS, Klotzsch, in R. Schomb. Reisen in Brit.-Guiana, iii. p. 1089;

Engler, in Linnæa, xxxvi. (1870), p. 605; Oliver, in Trans. Linn. Soc. Ser. II.

Bot. ii. (1887), p. 273.

Summit of Mount Roraima, *McConnell & Quelch*, 92, 636.—Endemic.

The foliage of this plant is somewhat variable. In some examples the leaves are all trifoliolate, this being the typical form; in others they all consist of one leaflet only, whilst others have trifoliolate and unifoliolate leaves on the same branch.

WEINMANNIA FAGARIOIDES, H. B. & K. Nov. Gen. & Sp. vi. p. 54.

W. glabra, Linn. f. *var.*?, Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 273.

Summit of Mount Roraima, *McConnell & Quelch*, 639.—Also on the Andes of Peru, Bolivia, and Ecuador, from 6000–10,000 ft. elevation.

The specimens collected by im Thurn upon Roraima have longer leaves, with more

numerous and narrower leaflets and longer and more lax racemes than those of the present collections; but I believe both forms belong to *Weinmannia fagaroides*, H. B. & K., which seems to be somewhat variable in the characters mentioned.

DROSERACEÆ.

DROSERA COMMUNIS, A. St. Hil. Pl. Rem. Bres. p. 267; Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 273.

D. Roraimæ, Klotzsch, in R. Schomb. Reisen in Brit.-Guiana, iii. p. 1090.

Upper slopes of Mount Roraima, *McConnell & Quelch*, 42. Summit of Mount Roraima, *McConnell & Quelch*, 85, 681. Arabapu River, *McConnell & Quelch*, 154.—Widely distributed in eastern South America, westward to New Granada, and southward to Paraguay.

MYRTACEÆ.

CAMPOMANESIA COÆTANEA, Berg, in Mart. Fl. Bras. xiv. pt. i. p. 44.

Kanuku Mountains, *McConnell & Quelch*, 275.

MYRTUS MYRICOIDES, H. B. & K. Nov. Gen. & Sp. vi. p. 131, t. 539.

M. sp. nov. aff. myricoides, Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 273.

Summit of Mount Roraima, *McConnell & Quelch*, 495.—Also in the Andes of Colombia up to 11,000 ft. elevation.

This was considered to be a new species by Prof. Oliver, but I cannot distinguish it by any specific character from *M. myricoides*, H. B. & K.

MYRTUS RORAIMENSIS, N. E. Brown, sp. n. Omnino glabra. Folia parva, anguste oblonga vel oblongo-lanceolata, obtusa, coriacea, utrinque glanduloso-punctata, supra nitida, rugulosa. Flores axillares. Petala orbicularia, glanduloso-punctata. Stamina quam petala breviora. Ovarium 3-loculare.

A branching shrub, quite glabrous in all parts. Leaves small, 4–6 lin. long, including the $\frac{1}{2}$ –1 lin. long petiole, $1\frac{1}{2}$ – $3\frac{1}{2}$ lin. broad, usually narrowly oblong or oblong-lanceolate, occasionally ovate, obtuse, rounded at the base, rigidly coriaceous, shining and rugulose above in the dried state, pale beneath, gland-dotted on both sides. Flowers axillary, solitary. Pedicels $1\frac{1}{2}$ –3 lin. long, bibracteolate at the apex. Bracteoles $1\frac{1}{2}$ – $1\frac{3}{4}$ lin. long, $\frac{1}{3}$ – $\frac{1}{2}$ lin. broad, linear or linear-lanceolate, obtuse, coriaceous, gland-dotted. Calyx-tube $\frac{3}{4}$ lin. long, hemispherical; lobes about 1 lin. long, $\frac{2}{3}$ – $\frac{3}{4}$ lin. broad, oblong or ovate-oblong, obtuse, gland-dotted. Petals about 2 lin. long and broad, suborbicular, very obtuse, gland-dotted. Stamens shorter than the petals. Ovary 3-celled; ovules several in each cell. Berry drooping, globose, about 3 lin. diam., crowned with the calyx-lobes.

Summit of Mount Roraima, 8600 ft., *McConnell & Quelch*, 635, 641.

This species somewhat resembles *M. myricoides*, H. B. & K., but differs in having perfectly glabrous stems, smaller, narrower, and less acute leaves, shorter pedicels, and smaller calyx-lobes.

LECYTHIS VENUSTA, Miers, in Trans. Linn. Soc. xxx. (1874), p. 214.

Mazaruni River, 200 ft., *McConnell & Quelch*, 713.

MELASTOMACEÆ.

MICROLICIA BRYANTHOIDES, Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 274 t. 39. f. B; Cogn. in DC. Monog. Phan. vii. p. 49.

Summit of Mount Roraima, *McConnell & Quelch*, 630.—Endemic.

RHYNCHANThERA GRANDIFLORA, DC. Prod. iii. p. 107.

Ireng Valley, *McConnell & Quelch*, 230.

SIPHANTHERA MICROLICIOIDES, Cogn. in Mart. Fl. Bras. xiv. pt. III. p. 199; and in DC. Monog. Phan. vii. p. 113.

Meisneria cordifolia, Benth. in Hook. Journ. Bot. ii. (1840), p. 299; R. Schomb. Reisen in Brit.-Guiana, iii. p. 1101.

M. microlicioides, Naud. in Ann. Sc. Nat. Sér. III. xii. (1849), p. 204; Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 274.

Upper slopes of Mount Roraima, *McConnell & Quelch*, 8. Kotinga Valley, *McConnell & Quelch*, 182.—Apparently endemic in the Roraima region.

DESMOSCELIS VILLOSA, Naud. in Ann. Sc. Nat. Sér. III. xiii. (1850), p. 30.

Kotinga Valley, *McConnell & Quelch*, 195.

MACAIREA ASPERA, N. E. Brown, sp. n. (Plate 2.) Folia oblonga, obtusa, supra scaberrima, subtus appresse rufo-tomentosa. Calyx appresse pubescens et glandulosus. Petala obovato-oblonga, obtusissima, glanduloso-ciliata. Stamina filamenta superne glandulosa. Ovarium superne dense glandulosum, stylo parce glanduloso.

A branching shrub; clothed on the young shoots, branches of the panicle, pedicels and calyx with rather long, rust-coloured, adpressed hairs. Leaves opposite; petiole 4–6 lin. long, pubescent like the stems; blade $1\frac{1}{4}$ –3 in. long, 8–17 lin. broad, oblong, obtuse, or shortly and stoutly apiculate, cuneately rounded into the petiole at the base, coriaceous, 3-nerved, with the lateral nerves close to the revolute margin, scabrid with minute, conical, hair-tipped tubercles on the upper surface, tomentose beneath with a dense adpressed pubescence of rust-coloured hairs. Panicles 4–6 in. long, 2–3 $\frac{1}{2}$ in. broad, many-flowered. Pedicels 1–2 $\frac{1}{2}$ lin. long, minutely bibracteolate in the middle. Calyx with sessile glands mingled with the hairs outside; tube campanulate, 1 $\frac{1}{2}$ lin. long; lobes 1 $\frac{1}{2}$ lin. long, deltoid, acuminate. Petals 3–4 lin. long, 2 $\frac{1}{2}$ –3 lin. broad, oblong-ovate, very obtuse, minutely gland-ciliate, glabrous. Stamens glandular on the apical part of the filaments. Anthers with the connective dilated at the base into a cordate-orbicular shield. Ovary free, densely covered with glands on the upper part; style filiform, sparsely glandular.

Upper slopes of Mount Roraima, *McConnell & Quelch*, 31.

Another specimen (no. 24, *McConnell & Quelch*), collected at the same place as no. 31, is identical with it in all particulars excepting the anthers, in which the portion of the connective between the pollen-cells and the dilated base is very much longer than in 31,

as shown in figs. 6-7, but I cannot consider this variation of sufficient importance to separate the two specifically.

Allied to *Macairea thyrsiflora*, DC., and *M. albiflora*, Cogn.; differing in the very much rougher upper surface and different indumentum of the lower surface of the leaves and in the stouter anthers. *M. albiflora*, Cogn., according to Spruce's label, has purple flowers, not white as described by Cogniaux.

MACAIREA PACHYPHYLLA, Benth. in Hook. Journ. Bot. ii. (1840), p. 292; R. Schomb. Reisen in Brit.-Guiana, iii. pp. 1100, 1192; Cogn. in DC. Monog. Phan. vii. p. 178.

Kotinga Valley, *McConnell & Quelch*, 181.—Apparently endemic to the vicinity of Roraima.

TIBOUCHINA FRATERNA, N. E. Brown, sp. n. Folia anguste elliptica, utrinque æqualiter subacuta, 3-nervia, subtus appresse squamosa, supra striolata, et marginibus et secundum lineas duas inter nervos squamosa. Bracteæ liberæ, oblongæ, obtusæ. Calyx appresse squamosus, 5-lobus, tubo cylindrico quam lobis longiore. Petala elliptico-obovata, unguiculata, glanduloso-ciliata.

Branchlets densely clothed with imbricating, lanceolate, acuminate, brownish scales about $\frac{1}{2}$ lin. long, those at the nodes about twice as long and more acuminate. Leaves spreading; petiole 1-1 $\frac{1}{2}$ lin. long, scaly; blade $\frac{3}{4}$ -1 in. long, 4-6 lin. broad, narrowly elliptic, about equally acute or subacute at both ends, apiculate, 3-nerved, the underside clothed with adpressed acuminate scales, those on the nerves much larger than the rest, the upper surface finely striate, with a longitudinal row of minute scales near each of the lateral nerves and along the margins. Cymules 2-3-flowered, subsessile or with peduncles up to 3 $\frac{1}{2}$ lin. long, axillary and terminal, forming a terminal inflorescence 1-2 in. long, 1 in. broad, clothed to the tips of the calyx-lobes with pale, rosy-tinted acuminate scales. Bracts free, 2-4 lin. long, about 1 $\frac{1}{2}$ lin. broad, oblong, obtuse, resembling reduced leaves; bracteoles smaller, thinner, more glabrous, free, ciliate with long hair-like scales. Pedicels very short, scarcely 1 lin. long. Calyx-tube 3 lin. long and about 1 $\frac{1}{2}$ lin. diam., cylindric-campanulate; lobes 2 lin. long, 1 $\frac{1}{4}$ lin. broad at the base, whence they gradually taper in a nearly straight line to an acute point, erect or very slightly spreading, glabrous inside, ciliate. Petals 7 lin. long, 3 $\frac{1}{2}$ lin. broad, elliptic-obovate, very obtuse, clawed, minutely ciliate with gland-tipped hairs, and with a few long simple hairs at the apex, otherwise glabrous. Stamens 10, unequal, glabrous; the longer anthers 5 lin. long, the shorter 4 lin. long, all attenuate to a subulate point, opening by one terminal pore; connective bifid at its insertion on the filament. Ovary free, oblong, densely covered with hair-like scales; style slender, glabrous.

Upper slopes of Mount Roraima, *McConnell & Quelch*, 17.

A very distinct species, which, according to the arrangement of Cogniaux, I think should be placed in the section *Lepidotæ*, although it bears very little resemblance to the only two species (*T. lepidota*, Baill., and *T. paleacea*, Cogn.) at present placed under that section. It has much more resemblance to *T. Spruceana*, Cogn., and *T. aspera*, Aubl.,

but the free (not connate) bracts and bracteoles at once separate it from the section to which they belong.

TIBOUCHINA LASIOPHYLLA, Cogn. in Mart. Fl. Bras. xiv. pt. III. p. 297; and in DC. Monog. Phan. vii. p. 270.

Pterolepis lasiophylla, Triana, in Trans. Linn. Soc. xxviii. (1871), p. 40; Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 274.

Kotinga Valley, *McConnell & Quelch*, 183, 197. Ireng Valley, *McConnell & Quelch*, 205. Slope of Mount Roraima, at 5400 ft. *in Thurn*, 59.—Only known from the above localities.

TIBOUCHINA ASPERA, Aubl. Pl. Guian. i. p. 446, t. 177; R. Schomb. Reisen in Brit.-Guiana, iii. pp. 1100, 1191.

Pteroma tibouchinum, Triana, in Trans. Linn. Soc. xxviii. (1871), p. 45, t. 3; Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 274.

Ireng Valley, *McConnell & Quelch*, 217.—Widely spread in Tropical South America, extending westward to Colombia and Peru.

MAR CETIA TAXIFOLIA, DC. Prod. iii. p. 124; R. Schomb. Reisen in Brit.-Guiana, ii. p. 216, and iii. p. 1100; Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 273.

Upper slopes of Mount Roraima, *McConnell & Quelch*, 13, 20; *in Thurn*, 68; *Appun*, 1114, 1153. Kotinga Valley near Roraima, *McConnell & Quelch*, 149.

With the exception of the above localities, this species is only known from Brazil, where it is a common and widely spread plant.

MAR CETIA JUNIPERINA, DC. Prod. iii. p. 125; Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 274.

Summit of Mount Roraima, *McConnell & Quelch*, 86, 644.—Also in Venezuela at Cumana and Merida.

CHÆTOLEPIS ANISANDRA, Naud. in Ann. Sc. Nat. Sér. III. xiv. (1850), p. 140; Cogn. in DC. Monog. Phan. vii. p. 171.

On the face of the cliff of Mount Roraima, at 7000–8000 ft., *McConnell & Quelch*, 22.—Endemic.

MONOCHÆTUM BONPLANDII, Naud. in Ann. Sci. Nat. Sér. III. xiv. (1850), p. 51; Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 274; Cogn. in DC. Monog. Phan. vii. p. 393.

Kotinga Valley, *McConnell & Quelch*, 189.—Also in Venezuela, Colombia, Ecuador, and Peru.

MICONIA HOLOSERICEA, DC. Prod. iii. p. 181.

M. albicans, Steud. Nom. ed. 2, ii. p. 139.

Upper slopes of Mount Roraima, *McConnell & Quelch*, 28.—Throughout Tropical America from Mexico to Paraguay.

MICONIA TINIFOLIA, Naud. in Ann. Sc. Nat. Sér. III. xvi. (1851), p. 225.

Kotinga Valley, *McConnell & Quelch*, 191. Summit of Mount Roraima, *McConnell & Quelch*, 633.—Also in Venezuela.

LYTHRACEÆ.

CUPHEA GRACILIS, H. B. & K. Nov. Gen. & Sp. vi. p. 199, var. MEDIA, Koehne, in Mart. Fl. Bras. xiii. pt. II. p. 284.

Ireng Valley, *McConnell & Quelch*, 225.

ONAGRACEÆ.

JUSSIEA LONGIFOLIA, DC. in Mém. Soc. Phys. Genève. ii. pt. II. (1824), p. 141.

Ireng Valley, *McConnell & Quelch*, 208, 214, 227.

TURNERACEÆ.

TURNERA ULMIFOLIA, Linn. Sp. Pl. ed. 1, p. 271.

Kotinga Valley, *McConnell & Quelch*, 174. Ireng Valley, *McConnell & Quelch*, 256, 260.

TURNERA VELUTINA, Benth. in Hook. Journ. Bot. iv. (1842), p. 116.

Ireng Valley, *McConnell & Quelch*, 240.

PIRIQUETA GUIANENSIS, N. E. Brown, sp. n. Perennis, caulibus simplicibus, pilis stellatis cum setulis intermixtis tomentosa. Folia petiolata, anguste vel oblongo-lanceolata, obtusa, obtuse dentata, dense tomentosa. Flores axillares, solitarii. Pedicelli 4–10 lin. longi, supra medium articulati. Petala purpurea?, glabra. Fructus globosus, dense appresse pubescens. Semina oblonga, reticulato-foveolata.

A perennial herb growing to about 1 ft. high. Stems several from the same root-stock, rather slender, usually simple, tomentose with a short down of stellate hairs intermingled with longer spreading hairs. Leaves petiolate; petiole 1–2 lin. long, blade $\frac{1}{2}$ –1 in. long, $1\frac{1}{2}$ –4 lin. broad, narrowly oblong or oblong-lanceolate, obtuse, obtusely 3–6-toothed on each side, densely stellate tomentose on both sides. Flowers axillary, solitary. Pedicels 4–10 lin. long, jointed above the middle, stellate-tomentose. Calyx 3–4 $\frac{1}{2}$ lin. long, stellate-tomentose outside; tube about 1 lin. long, shortly obconical, lobes 2–3 $\frac{1}{2}$ lin. long, 1–1 $\frac{1}{2}$ lin. broad, ovate-lanceolate or lanceolate, acute. Petals much longer than the calyx, obovate, glabrous, with a fringed scale at their base, apparently purple. Stamens about 2 $\frac{1}{4}$ lin. long, with glabrous filaments. Ovary ovoid, densely adpressed-pubescent, not tubercled; styles 2 lin. long, with a few scattered hairs or almost glabrous; stigmas dilated. Capsule globose, about 2 lin. diam., 3-valved, densely covered with adpressed, rather silky hairs. Seeds 1 lin. long, oblong, rather coarsely pitted, pale reddish brown.

British Guiana; without precise locality, *Appun*, 1858. Ireng Valley, *McConnell & Quelch*, 264.

Allied to *P. Duarteana*, Urb.

PIRIQUETA VILLOSA, Aubl. Pl. Guian. i. p. 298, t. 117.

P. cistoides, Mey. ex Steud. Nom. ed. 2, ii. p. 344.

Kotinga Valley, *McConnell & Quelch*, 178.

PASSIFLORACEÆ.

PASSIFLORA QUELCHII, N. E. Brown, sp. n. (Plate 3.) Frutex vel arbor, glaber. Folia petiolata, cuneato-oblonga, acuta vel obtusissima, undulata, coriacea; petiolus apice biglandulosus. Stipulæ minutæ, subulatæ, deciduæ. Cirrhi nulli. Flores in racemum abbreviatum dispositi, glabri. Calycis tubus elongatus, cylindricus. Sepala petalis subconformia, oblongo-lanceolata, obtusa. Corona faucialis uniserialis, multipartita, segmentis complanatis prope apicem dilatatis oblique acuminatis; corona interior e tubo versus basin emergens, multipartita, segmentis complanatis lineari-falcatis.

Shrubby or arborescent, not climbing; branches woody, glabrous. Leaves simple, entire, very coriaceous, 2-4 lin. distant, spreading, glabrous; petiole 4-6 lin. long, moderately stout, with two large sessile glands at the apex; blade 3-5 in. long, $\frac{3}{4}$ -1 $\frac{1}{2}$ in. broad, cuneately oblong, gradually tapering from $\frac{3}{4}$ the way up in a slightly curved line to an acute base, shortly acute or very obtuse and slightly emarginate with an apiculus in the notch, with strongly undulated margins; primary lateral veins 12-15 on each side of the midrib, very spreading, uniting close to the margin in a series of broad loops; secondary veins much reticulated, all prominent on both sides. Stipules minute, subulate, very deciduous. Tendrils none. Flowers in short racemes or clusters, probably on the older parts of the branches; axis of the inflorescence $\frac{1}{3}$ - $\frac{3}{4}$ in. long, several-flowered, glabrous. Pedicels 1-2 lin. long, glabrous. Calyx-tube about 1 in. long, 1 $\frac{1}{2}$ -2 lin. diam. in the dried state, cylindric, slightly enlarged at the mouth, glabrous; the lobes 10-12 lin. long, 2 $\frac{1}{3}$ -3 $\frac{1}{2}$ lin. broad, oblong-lanceolate, obtuse or subacute. Petals similar to the calyx-lobes and of about the same size. Corona in two series; one placed towards the bottom of the calyx-tube, consisting of numerous linear-falcate, acute, erect filaments about 1 $\frac{1}{2}$ lin. long; the other at the mouth of the tube, consisting of numerous flattened, somewhat hatchet-shaped segments rather more than $\frac{1}{2}$ in. long, with the dilated part slightly toothed and dorsally produced into a short point. Gynophore nearly 1 $\frac{1}{2}$ in. long, terete or perhaps sulcate at the base, glabrous. Filaments of the stamens 2 lin. long. Ovary oblong, 9-striate; styles 2 lin. long, stigmas capitate.

Ireng Valley, *McConnell & Quelch*, 207.

A very distinct species, easily distinguished from all the others of shrubby habit by its cuneately oblong, undulated leaves, shortly racemose flowers, and long calyx-tube, besides other characters.

ARALIACEÆ.

DIDYMOPANAX RUGOSUM, N. E. Brown, sp. n. Folia 3-foliolata, foliolis ellipticis vel elliptico-oblongis utrinque obtusis, rigide coriaceis, supra glabris rugosis, subtus dense tomentosis. Umbellæ compositæ, longe pedunculatæ, fulvo-tomentosæ, multifloræ. Petala ovata, acuta. Drupa immatura valde compressa, suborbiculata.

Leaves digitately trifoliolate; petiole stout, $\frac{1}{4}$ in. thick, tomentose, broken in the only specimen seen and about $3\frac{1}{2}$ in. long; leaflets rigidly coriaceous, above glabrous and rugose, with a narrow prominent midrib and impressed veins and veinlets, beneath tomentose on the very stout prominent midrib and the primary veins, and covered with a dense blanket-like tawny tomentum between them; the lateral pair $3\frac{1}{2}$ –4 in. long, $2\frac{1}{2}$ – $2\frac{3}{4}$ in. broad, elliptic, obtuse at both ends, mucronulate, with stout petiolules about 2 lin. long; the terminal leaflet $4\frac{3}{4}$ in. long, $2\frac{3}{4}$ in. broad, elliptic-oblong, slightly emarginate at the obtuse apex, mucronulate, with a stout petiole 4 lin. long. Umbel very compound, 6–8-rayed, many-flowered, $2\frac{3}{4}$ – $3\frac{1}{2}$ in. diam., with a stout, more or less flattened peduncle 4–7 in. long, 2–3 lin. thick, everywhere, to the outside of the petals, densely covered with a tawny tomentum, becoming greyish with age. Secondary umbels 6–7-rayed, some of the rays bearing a single flower, others dividing into tertiary umbels, some of which again divide in a similar manner, forming 2–4-flowered umbels of a fourth order. Bracts and bracteoles suborbicular, about $\frac{1}{2}$ –1 lin. diam. Pedicels $1\frac{1}{2}$ –2 lin. long, stout, not jointed. Calyx-limb shortly cupular, 5-toothed; teeth $\frac{1}{4}$ – $\frac{1}{3}$ lin. long, deltoid, acute. Bud globose. Petals 5, about $\frac{3}{4}$ lin. long, and nearly as broad, deltoid-ovate, acute, rather thick, glabrous within. Stamens 5, not longer than the petals. Ovary 2-celled; styles 2, somewhat flattened, divergent in young fruit, which is suborbicular in outline and compressed.

Summit of Mount Roraima, 8600 ft., *McConnell & Quelch*, 663.

A most distinct species, in general habit more like *Didymopanax Spruceanum*, Seem., than any other; but the very thick rigid leaves, rugose above and tomentose beneath, at once distinguish it from that and all others.

SCIADOPHYLLUM UMBELLATUM, N. E. Brown, sp. n. Folia digitata, foliolis 6–8 petiolulatis oblongo-lanceolatis vel oblanceolato-oblongis obtuse acutis basi cuneatis supra glabris subtus tomento appresso pallido vestitis. Umbellæ compositæ, longissime pedunculatæ, multifloræ. Petala ovata, acuta. Fructus immaturus 5-angularis, compressus.

Leaves digitately 6–8-foliolate; petiole 4–6 in. long, $1\frac{1}{2}$ –2 lin. thick, glabrous; leaflets coriaceous, unequal, $2\frac{1}{2}$ –6 in. long, excluding the 2–7 lin. long petiolules, 1 – $2\frac{1}{4}$ in. broad, oblong-lanceolate or oblanceolate-oblong, obtusely pointed at the apex, cuneate at the base, entire or very slightly repand, slightly revolute along the margins, glabrous above, with a thin layer of pale close tomentum beneath; midrib prominent on both sides, stouter beneath; primary lateral veins 9–16 on each side of the midrib, rather slender and not very prominent. Umbel compound, 2–3 in. diam., on a peduncle 9–12 in. long, $2\frac{1}{2}$ lin. thick; rays numerous, unequal, $\frac{1}{2}$ – $1\frac{1}{4}$ in. long, $\frac{3}{4}$ –1 lin. thick. Bracts minute, rounded. Secondary umbels 10–16-flowered. Pedicels 1 – $1\frac{3}{4}$ lin. long. Calyx-limb annular, entire or very minutely 5-toothed. Petals 5, acute. Stamens 5, not exceeding the petals. Ovary 3–5-celled; style or column of united styles $1\frac{1}{3}$ lin. long, minutely 3–5-fid at the apex. Young fruit deeply 3–5-angled, angles compressed.

Summit of Mount Roraima, 8600 ft. *McConnell & Quelch*, 666.

Allied to *S. coriaceum*, Marchal, from which it differs in its more pointed leaves, very

much shorter and stouter pedicels, and different indumentum. The indumentum which covers the whole of the inflorescence is rather peculiar, and is neither scurfy nor hairy, and when highly magnified seems to consist of a very thin dense covering of very minute, flattened, and very closely adpressed hairs, producing a greyish hue.

This species and *Sciadophyllum coriaceum* differ from all the others in the genus in having the flowers collected into a compound umbel terminating a long peduncle, and not paniculate; for although *S. coriaceum* has a second whorl of umbels below the terminal one, they all spring from one level, and the inflorescence cannot be called a panicle. The styles also of these two species are united in a column, and in this character, and indeed in their entire floral structure, they quite agree with *Heptapleurum*, a genus that is only artificially separated from *Sciadophyllum* by this one point, as has already been pointed out in Benth. & Hook. f. *Genera Plantarum*, i. pp. 940, 942. As there are some species of *Heptapleurum* in which the styles are at first united in a column and during the growth of the young fruit become free, it appears to me that *Sciadophyllum* should include *Heptapleurum*, when it would form a genus fairly uniform in character and of world-wide distribution within the tropics.

RUBIACEÆ.

HENRIQUEZIA JENMANI, K. Schum. in Mart. Fl. Bras. vi. pt. VI. p. 135. (Plate 4.)

Mazaruni River, *McConnell & Quelch*, 711; *Jenman*, 629.

This handsome tree appears only to have been collected in this single locality.

CHALEPOPHYLLUM SPECIOSUM, N. E. Brown, sp. n. (Plate 5, figs. 10-17.) Frutex glaber, ramis tetragonis. Folia opposita ad apicem ramorum subconferta, coriacea, sessilia, obovata, obtusa, subapiculata. Flores in axillis supremis (foliis delapsis) solitarii, breviter pedicellati, bibracteati. Calycis lobi æquales vel inæquales, lineares vel oblongi, acuti. Corolla hypocrateriformis, 5-lobata, tubo longissimo, lobis anguste lanceolatis vel oblongis acutis. Stamina inclusa. Stylus sæpius inclusus.

A stoutly-branched shrub, glabrous in all parts excepting the inside of the corolla. Branches 4-angled, 2 lin. thick; internodes very short, $1\frac{1}{2}$ -5 lin. long. Leaves opposite, in a cluster of 3-5 pairs at the tips of the branches, subsessile, $1\frac{1}{4}$ -2 in. long, $\frac{1}{2}$ -1 in. long, more or less obovate, obtuse, bluntly apiculate, cuneate at the base, rigidly coriaceous in the dried state, shining above, pale beneath, slightly revolute along the margins; midrib stout and conspicuous beneath; veins inconspicuous or not at all visible. Stipules broad-based, abruptly contracted into a linear point about $1\frac{1}{2}$ lin. long, persistent long after the leaves have fallen. Flowers usually 2 to each shoot, or sometimes 1 only, arising immediately below the terminal tuft of leaves in the axils of fallen leaves or in those of the lowest pair, solitary in each axil. Pedicels 1-4 lin. long, stout, flattened, bibracteate at the base of the ovary. Bracts 4-8 lin. long, $1\frac{1}{2}$ -5 lin. broad, of the form and substance of reduced leaves. Calyx-lobes 5-10 lin. long, $1-2\frac{1}{2}$ lin. broad, equal or unequal, varying from linear to oblong, acute, erect, coriaceous. Corolla very variable in size, hypocrateriform, regular, 5-lobed; tube $1\frac{3}{4}$ - $3\frac{1}{2}$ in. long, about $1\frac{1}{2}$ lin.

diam., cylindric, 5-grooved, slightly enlarged and densely hairy inside at the throat around the anthers; lobes spreading or recurved, 1–2 in. long, 2–6 lin. broad, linear-lanceolate or oblong, acute, glabrous, contorted in the bud. Stamens 5, included, inserted in the throat of the corolla-tube; filaments about $2\frac{1}{2}$ lin. long; anthers 3– $3\frac{1}{2}$ lin. long, dorsifixed, with a rather long and broad attachment to the filaments, linear, acute, bifid at the base, glabrous. Ovary $2\frac{1}{2}$ –3 lin. long, stoutly obconical, 2-celled; style about as long as the corolla-tube, or shorter than the anthers or rarely exserted, glabrous; stigma of two oblong obtuse lobes about 1 lin. long. Ovules numerous, flattened, on broad placentas that are peltately attached along their middle by a very short narrow plate to the septum. Fruit not seen.

Summit of Mount Roraima, 8600 ft., *McConnell & Quelch*, 100, 305, 653.

The variation in the length of the style would seem to imply that the flowers are heterostyled, but it may be due to a lengthening at various periods of development. The stems in the figure on Plate 5 are not represented of sufficient stoutness.

SIPANEA PRATENSIS, Aubl. Pl. Guian. i. p. 147, t. 56; in Thurn & Oliver, in Trans. Linn. Soc. ser. II. ii. (1887), pp. 261, 276.

Kotinga Valley, *McConnell & Quelch*, 175, 185, 198.—Widely distributed in eastern Tropical America from Trinidad to Minas Geraes and westward to Colombia.

DIDYMOCHLAMYS CONNELLII, N. E. Brown, sp. n. (Plate 5, figs. 1–9.) Folia alterna, oblique oblanceolata, subabrupte acuta, glabra. Bracteae exteriores suborbiculares, brevissime cuspidatae. Corollae tubus supra medium ampliatus, ore contractus; lobi infra medium bilobulati lobulis lineari-oblongis undulatis subtortis.

A small herb, with a simple stem $1\frac{1}{4}$ –6 in. long. Leaves alternate, distichous, 1– $1\frac{1}{3}$ in. long, including the very short petiole, $2\frac{1}{2}$ – $4\frac{1}{2}$ lin. broad, obliquely or subfalcately oblanceolate, acute or rounded in at the apex to a very short cuspidate point, tapering from above the middle to an unequal base, the upper side of which is very narrow and acute, and the lower side broadly rounded or subtruncate, glabrous on both sides; under surface pale, densely covered with large stomata, which appear as whitish dots under a lens; stipule solitary, 3–4 lin. long, linear, acuminate, glabrous. Peduncle terminal, about $\frac{1}{4}$ in. long, bearing two suborbicular, apiculate, glabrous bracts, about $\frac{1}{2}$ in. in length and breadth, enclosing 2 (or more?) shortly pedicellate flowers and 1 (or more?) linear acute bracteole. Calyx-lobes slightly unequal, $\frac{3}{4}$ –1 lin. long, lanceolate, acute, erect, glabrous. Corolla-tube about $\frac{1}{2}$ in. long, probably somewhat compressed, dilated a little above the middle, then narrowed to the mouth, glabrous, with the exception of a ring of hairs inside near the base; lobes divided to slightly below the middle into two diverging, linear-oblong, obtuse, undulated and somewhat twisted lobules $1\frac{1}{2}$ lin. long and rather more than $\frac{1}{2}$ lin. broad, the basal entire part being about 1 lin. long and broad, deltoid-ovate, slightly concave, with a minute, obtuse, incurved point at the apex between the lobules. Stamens 5, included, inserted below the middle of the corolla-tube, unequal, 2 longer than the rest and inserted higher up, having on one side a pair and on the other side one of the shorter stamens alternating with them; filaments glabrous,

anthers linear. Glands of the disk 4, in 2 opposite pairs, the larger pair partly enclosing the smaller pair. Ovary inferior, laterally compressed, 2-celled; style included, filiform; stigma of 2 short oblong lobes; placenta ascending from near the base of the septum, thin; ovules numerous, flat.

On the Roraima Range at 3500 ft., *McConnell & Quelch*, 714; Kaieteur Falls on the Potaro River, *Jenman*, 7462.

This is one of the most remarkable plants in the collection; its alternate leaves and general habit give it the appearance of a Gesneraceous rather than of a Rubiaceous plant. The two large bracts enclosing the flowers and the bilobed corolla-lobes are very remarkable characteristics. It is very closely allied to *Didymochlamys Whitei*, Hook. f., from New Granada, the only other known species, but differs in its much smaller leaves and very much longer corolla-lobules. In *D. Whitei* the leaves are $1\frac{3}{4}$ – $3\frac{1}{2}$ in. long, 4–11 lin. broad, and taper at the apex into a long acuminate point, and the subquadrate lobules of the corolla are only about $\frac{1}{2}$ a line long and the same in breadth.

To the generic characters of *Didymochlamys* may be added the following description of the fruit, taken from a specimen of *D. Whitei* sent to Kew by Mr. R. B. White in 1882:—Fruit a laterally-compressed 2-celled capsule, dehiscing at the apex between the glands; placentas ascending from near the base of the septum, thin; seeds numerous, minute, ovate, flattened, thinly covered with rather long hairs, with a dense tuft at the narrower (micropyle?) end; testa rather coriaceous, moderately thin; albumen copious, oily; embryo minute.

The two smaller glands of the disk in the fruiting stage of *D. Whitei* are a little longer than the larger glands, they are tubular in the upper part and may assist in the dispersion of the seeds.

TOCOYENA NEGLECTA, N. E. Brown, sp. n. Folia breviter petiolata, lanceolata vel elliptico-lanceolata, acuta, basi cuneata, supra pubescentia, subtus tomentosa. Cymæ terminales, subsessiles, 9–20-floræ. Calyx acute 5-dentatus, plus minusve tomentosus. Corollæ tubus 3 – $3\frac{1}{2}$ poll. longus, extra tomentosus, intra glaber, fauce hirtus; lobi oblique oblongo-lanceolati, subacuti, glabri, ciliati.

Branches subterete or obscurely 4-angled, with a light brown bark, tomentose when young. Leaves opposite, 3–7 in. long, $1\frac{3}{4}$ – $3\frac{1}{2}$ in. broad, lanceolate, elliptic-lanceolate or slightly obovate, acute, cuneately tapering at the base into a petiole $1\frac{1}{2}$ –3 lin. long, somewhat harshly pubescent above, densely tomentose beneath, with the hairs along the veins distinctly diverging on either side; primary veins 9–13 on each side of the midrib, and together with the reticulated veinlets impressed above, prominent beneath. Stipules about 2 lin. long, triangular, acute, at first tomentose, becoming glabrous with age. Cymes terminal, subsessile, about 9–20-flowered. Flowers sessile. Calyx more or less tomentose; the tube $1\frac{1}{2}$ – $2\frac{1}{2}$ lin. long, oblong, slightly tapering downwards; the limb cup-shaped, with 5 deltoid acuminate teeth $\frac{3}{4}$ –2 lin. long. Corolla hypocrateriform, 5-lobed, white; the tube 3 – $3\frac{1}{2}$ in. long, about $1\frac{1}{2}$ lin. diam., slightly enlarged at the mouth, tomentose outside, glabrous within, except in the throat, which is densely hairy; the

lobes 10–12 lin. long, 5–6 lin. broad, obliquely oblong-lanceolate, subacute, glabrous on both sides, ciliate along the outer margin, but in the younger stage, when contorted into the lanceolate acuminate bud, the outside towards the base is more or less pubescent or tomentose. Anthers sessile at the mouth of the corolla-tube, partly exerted, 4 lin. long, oblong-lanceolate, obtuse. Ovary 2-celled; style as long as the corolla-tube, slender, glabrous; stigma of 2 oblong obtuse lobes, about 2 lin. long, 1 lin. broad. Fruit globose, about $\frac{3}{4}$ in. diam., thinly pubescent or subglabrous.

British Guiana: Maimatta on the Rupununi River, *Jenman*, 5525, 5742; Kotinga Valley, *McConnell & Quelch*, 163; Roraima, *Schomburgk*, 478; Suruma River, *Schomburgk*, 772 B.

Allied to *Tocoyena foetida*, Poepp. & Endl., but differs in its much shorter petioles and corolla-tube. From *T. velutina*, Spruce (which is a very distinct species, wrongly referred to *T. foetida* by K. Schumann), it differs in the less elliptic form of the leaves, in the indumentum, and very different calyx-limb. Schomburgk's specimens are quoted by K. Schumann in the 'Flora Brasiliensis,' vi. pt. VI. p. 348, under *T. formosa*, K. Schum., which is a quite different species, with orbicular leaves and short, ellipsoidal, obtuse buds.

RETINIPHYLLUM LAXIFLORUM, N. E. Brown. (Plate 6.) Folia petiolata, elliptica vel elliptico-oblonga, obtusa basi subcuneata, glabra. Racemus terminalis, 3–9-florus, floribus longe pedicellatis. Bracteoli sub ovario in cupulam disciformem minute denticulatam connati. Calyx non costatus, truncatus, minutissime 5-dentatus. Corolla utrinque sericeo-puberula, lobis reflexis tubo brevioribus.—*Patima laxiflora*, Benth. in Hook. Journ. Bot. iii. (1841) p. 220; R. Schomb. Reisen in Brit.-Guian. iii. p. 1142. *Synisoon Schomburgkianum*, Baill. in Bull. Soc. Linn. Paris (1879), p. 208; Baill. Hist. Pl. vii. p. 433; K. Schum. in Mart. Fl. Bras. vi. pt. VI. p. 398.

A shrub or tree, glabrous in all parts excepting the flowers, and exuding a varnish-like resin on various parts of the young branches and inflorescence. Branches moderately slender, with a brown bark. Leaves opposite, petiolate, thinly coriaceous; petiole 2–8 lin. long; blade $1\frac{1}{2}$ – $4\frac{3}{4}$ in. long, 1 – $2\frac{1}{4}$ in. broad, shining above, paler beneath with distinctly reticulated veins. Stipules subtruncate, minutely apiculate. Raceme erect, terminal, very lax, 2 – $2\frac{3}{4}$ in. long, consisting of 1–4 pairs of opposite flowers and one terminal one. Bracts reduced to a minutely denticulate raised line. Pedicels 6–8 lin. long, ascending, slender. Bracteoles beneath the ovary connate into a very small, disk-like, minutely-denticulate involucre. Calyx $3\frac{1}{2}$ –4 lin. long, with the limb produced for 2 – $2\frac{1}{3}$ lin. above the ovary, tubular, truncate, with 5 very minute distant teeth, glabrous, but more or less coated with the varnish-like secretion. Corolla erect, 5-lobed, minutely adpressed-pubescent outside and inside, with a dense ring of white hairs near the base of the tube inside, below which it is glabrous; tube 8–9 lin. long, $1\frac{1}{4}$ – $1\frac{1}{3}$ lin. diam., cylindric; lobes pendulous, reflexed from their base, 7 lin. long, $1\frac{1}{2}$ lin. broad, linear-oblong, obtuse, nearly straight along one margin curved along the other, strictly contorted in the bud. Stamens inserted at the mouth of the tube, reflexed with the corolla lobes; filaments 5 lin. long, pubescent; anthers 2 lin. long, dorsifixed, lanceolate, produced beyond the cells into a subulate point at the apex and into an oblong,

emarginate, membranous appendage at the base, glabrous; pollen apparently red. Ovary 5-celled, surmounted by a rather deep ring-like disk; style much exserted, $1\frac{1}{3}$ in. long, rather slender, pubescent; stigma slightly thickened, ovoid-conical, tipped with 5 minute points. Ovules 2 in each cell, collateral, axile, pendulous from near the top of the cell, and partly covered at the top by a cap-like outgrowth of the funicle-like placenta. Fruit a globose, ribbed berry about $\frac{1}{4}$ in. diam., crowned with the calyx-tube; endocarp hardened into 5 separable nutlets, 3-keeled on the back, triangular in transverse section, with a false cell under each lateral wing. Seeds not seen.

Aroie Creek, *im Thurn*, 6. Upper slopes of Mount Roraima, *McConnell & Quelch*, 18; *Appun*, 1175; *Schomburgk*, 724 (815 B), 158.

This very distinct species of *Retiniphyllum* appears to have been misunderstood by all authors. Originally it was doubtfully referred by Benth. to the genus *Patima*, and erroneously described as having very numerous seeds. Next, in Benth. and Hooker's 'Genera Plantarum,' ii. p. 98, it is confused with *Kutchubaea insignis*, Fisch., and the part of the generic description in that work, relating to the fruit, is drawn up from the present plant and not from the true *Kutchubaea*, which indeed it greatly resembles in habit, but differs in having a 5-lobed corolla, 5 exserted, reflexed stamens, with long filaments, a long exserted style, and in the presence of a minute disk-like involucl at the apex of the pedicel, which is jointed to the calyx; whilst in *Kutchubaea* the corolla is 8-lobed, the stamens are 8, with the anthers almost sessile and included in the throat of the corolla-tube, the style is included, and the pedicel is continuous with the calyx, without any trace of an involucl; the foliage and stipules of the two plants are also very different, besides disagreeing in several minor details. Finally Baillon made a distinct genus of it, which he placed next to *Knoxia*, where it has no affinity, for it is undoubtedly a true *Retiniphyllum*, allied to *R. scabrum*, Benth., but differing from that and all others of the genus by the long slender pedicels of its lax, few-flowered inflorescence. The involucl is characteristic of the genus *Retiniphyllum*.

PSYCHOTRIA CONCINNA, Oliver, in Trans. Linn. Soc. Ser. II. ii. (1887) p. 276, t. 42. f. B. Summit of Mount Roraima, *McConnell & Quelch*, 80, 89, 667.—Endemic.

PSYCHOTRIA CRASSA, Benth. in Hook. Journ. Bot. iii. (1841), p. 227; *im Thurn & Oliver*, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), pp. 265, 276; R. Schomb. Reisen in Brit.-Guiana, iii. p. 945.

Summit of Mount Roraima, *McConnell & Quelch*, 632. Also collected on Roraima by *Appun*, 1112.—Kanuku Mountains.

DIODIA HYSSOPIFOLIA, Cham. & Schlecht. in Linnæa, iii. (1828), p. 350.

Kotinga Valley, *McConnell & Quelch*, 190.

COMPOSITÆ.

VERNONIA ACUTA, N. E. Brown, sp. n. Rami tomentosi. Folia alterna, brevissime petiolata, lanceolata, acutissima vel subpungentia, basi acuta, supra glabra, subtus tenuiter et appresse pubescentia, glanduloso-punctata; petiolus tomentosus. Capitula in axillis aggregata, sessilia, sub-10-flora. Involuceri squamæ 5-6 seriatae, imbricatae, lanceolatae, acuminatae, pungentes, subtomentosae, exteriores gradatim minores. Achænia appresse pubescentia. Pappus biserialis, squamis exterioribus linearibus acutis quam setis interioribus multo brevioribus.

Branchlets densely tomentose. Leaves alternate, spreading; petiole 1-2 lin. long, tomentose; blade $1\frac{1}{4}$ -2 in. long, $\frac{1}{4}$ - $\frac{1}{2}$ in. broad, lanceolate, very acute and subpungent at the apex, acute at the base, entire, with revolute margins, rather rigid, glabrous above, except the midrib, which is pubescent, thinly covered with short adpressed hairs and gland-dotted beneath, with the veins and reticulated veinlets impressed above, prominent beneath. Heads in clusters of 3 or 4, sessile in the axils of the leaves at the ends of the branchlets, about $\frac{1}{4}$ in. diam., and about 10-flowered. Involucre-scales in 5-6 series, adpressed, lanceolate, acuminate, pungent-pointed, adpressed-pubescent or subtomentose, the innermost about 3 lin. long, $\frac{1}{2}$ - $\frac{2}{3}$ lin. broad, the others gradually smaller. Corolla $\frac{1}{4}$ in. long, 5-lobed, with the linear acute lobes of equal length with the tube, glabrous. Style-branches only about $\frac{1}{2}$ lin. longer than the stamens. Achenes rather more than 1 lin. long, adpressed-pubescent. Pappus in two series, white; the outer of numerous linear acute scales, the inner of numerous bristles, slightly scabrous, much longer than the outer series.

Kotinga Valley, *McConnell & Quelch*, 192.

EUPATORIUM IVÆFOLIUM, Linn. Syst. ed. 10, p. 1205, var. **GRACILLIMUM**, Baker, in Mart. Fl. Bras. vi. pt. II. p. 290.

Ireng Valley, *McConnell & Quelch*, 242.

EUPATORIUM SUBOBTUSUM, DC. Prod. v. p. 161.

Ireng Valley, *McConnell & Quelch*, 211, 304.

EUPATORIUM RORAIMENSE, N. E. Brown, sp. n. Herbacea, ramis breviter pubescentibus.

Folia opposita, petiolata, ovato-lanceolata, acuminata, glabra, crenato-dentata, petiolo pubescente. Capitula corymbosa, circa 20-flora. Involuceri squamæ lineares, acutæ, glabræ. Corollæ 5-dentata, glabra, tubo supra medium ampliato, campanulato, dentibus ovatis, acutis. Achænia 5-angulata, angulis scaberulis. Pappi setæ scaberulæ, roseo-purpureæ.

Stems herbaceous, with a very fine spreading pubescence. Leaves opposite; petiole 4-6 lin. long, pubescent; blade $1\frac{1}{4}$ - $2\frac{1}{2}$ in. long, $\frac{1}{2}$ -1 in. broad, ovate-lanceolate, acuminate, crenately toothed, with teeth 1-2 lin. broad, glabrous on both sides, rather thin and submembranous in texture in the dried state. Corymbs terminal, compact, 2- $2\frac{3}{4}$ in. diam., its branches $\frac{1}{4}$ -1 in. long, and, together with the slender $1\frac{1}{2}$ -3 lin. long pedicels, pubescent like the stem. Bracteoles almost filiform. Heads about $2\frac{1}{2}$ lin. diam., about 20-flowered. Involucre-scales 2-seriate, 2- $2\frac{1}{4}$ lin. long, linear, acute, nearly

or quite glabrous. Corolla glabrous, the tube 1 lin. long, slender below, dilated into a campanulate cup above the middle, with 5 ovate acute teeth $\frac{1}{2}$ lin. long. Achenes 1 lin. long, 5-angled, slightly scabrous along the angles, blackish. Pappus-bristles slightly scabrous, rosy-purple.

Summit of Mount Roraima, *McConnell & Quelch*, 640.

This somewhat resembles *Eupatorium Pentlandianum*, DC. (*Mandon*, 259), but the corolla, pappus, and achenes are all much shorter, the corolla is very different in form, the involucre-scales are longer and narrower, and the leaves are less sharply toothed.

EUPATORIUM FUSCUM, N. E. Brown, sp. n. Rami validi, cum petiolis et corymbis nigrescente-tomentosi. Folia breviter petiolata, late ovata, obtusa, basi rotundata vel subcordata, irregulariter et obtuse dentata, subcoriacea, supra glabra, glanduloso-punctata, subtus fusco-pubescentia. Corymbi compacti. Capitula 15-16-flora. Involucris squamæ biseriatæ, interiores lineari-oblongæ acutæ, exteriores minores. Corolla subcylindrica, breviter 5-dentata, tubo glabro, dentibus dorso minute papillatis. Pappi setæ scabridæ, plus minusve flexuosæ, subfulvescentes.

Flowering branches stout, 2-2 $\frac{1}{2}$ lin. thick, densely covered with a short blackish tomentum. Leaves opposite, rather crowded, 4-9 lin. distant; petiole 2-3 lin. long, stout, with a blackish tomentum, which spreads along the veins of the leaf; blade 1-1 $\frac{3}{4}$ in. long, 10 lin. to 1 $\frac{3}{4}$ in. broad, broadly ovate, obtuse, rounded or subcordate, very shortly and irregularly dentate, with subobtuse teeth, or nearly entire, somewhat coriaceous, glabrous (except along the veins,) and densely gland-dotted above, with a brownish or fuscous pubescence beneath; primary lateral veins 4-5 on each side of the midrib, 2-3 of them arising close to the base, the others at or above the middle, prominent beneath. Corymbs terminal, 1 $\frac{1}{2}$ -2 $\frac{1}{2}$ in. diam., compact, with short branches, clothed with the blackish tomentum as are also the pedicels and involucre. Heads about $\frac{1}{3}$ in. diam., 15-16-flowered. Involucral-scales in two series; the inner about 2 lin. long, $\frac{1}{2}$ lin. broad, linear-oblong acute; the outer a little shorter, narrower and tapering from the base to an acute point. Corolla 2 lin. long, tubular, subcylindric, 5-toothed, glabrous, with some minute papillæ on the back of the $\frac{1}{3}$ lin. long, acute teeth. Ovary pubescent in the upper part with short fuscous subglandular hairs. Pappus-bristles numerous, scabrid, more or less flexuose, dull tawny. Ripe achenes not seen.

Summit of Mount Roraima, 8600 ft., *McConnell & Quelch*, 648.

A very distinct species, which would appear to be best placed near *E. nummularia*, Hook. & Arn.

BACCHARIS VITIS-IDÆA, Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 277, t. 43. f. A.

Upper slopes and summit of Mount Roraima, *McConnell & Quelch*, 27, 91, 650.—Endemic.

HETEROTHALAMUS DENSUS, N. E. Brown, sp. n. (Plate 7, figs. 1-7.) Planta nana, perennis, dioica, dense foliosa. Folia patentia, lanceolata, acuta vel obtusa, basi in petiolum attenuata, integra, coriacea, nitida. Capitula sessilia, 2-4-na ad apices ramorum conferta vel solitaria, discoidea. Flores hermaphroditi steriles, corollæ tubo anguste

infundifuliformi, lobis 5 linearibus acutis patentibus. Florum foeminarum corolla angustissime tubulosa, truncata, vel oblique truncata, minutissime denticulata, stylo multo brevior.

A dwarf dioecious perennial with erect woody stems, apparently about 3-5 in. high, rooting at the base, sparingly branched in the upper part, densely leafy to the top, quite glabrous in all parts. Leaves densely crowded, spreading, $\frac{1}{3}$ - $\frac{2}{3}$ in. long (including the petiole), 1-2 lin. broad, lanceolate, acute or obtuse, tapering at the base into a linear petiole, entire, coriaceous, shining above, wrinkled from shrinkage in drying, veinless. Heads subunisexual, solitary or 2-4 together, sessile at the apex of the branches, and surrounded, but not overtopped by the leaves, about $2\frac{1}{2}$ lin. diam., discoid, many-flowered. Involucre campanulate, a little shorter than the florets, its scales in 2-3 series, subequal, about $1\frac{1}{2}$ lin. long, $\frac{2}{3}$ lin. broad, oblong or lanceolate, obtuse or acute, ciliate. Receptacle naked in the sterile heads, but with several linear or linear-lanceolate acute scales in the female heads, scattered among, and about as long as, the florets. Florets of the sterile heads structurally hermaphrodite, but with barren ovaries; corolla with a tube 1 lin. long, gradually widening at the mouth, and 5 linear, acute, spreading lobes $\frac{1}{2}$ lin. long, glabrous. Florets of the female heads with a very slender tubular corolla, truncate or oblique and very minutely 5-toothed at the apex, much shorter than the shortly bifid style, glabrous. Ovary glabrous. Pappus-bristles numerous and alike in both kinds of flowers, equalling the style in the female florets, slightly scabrid, whitish.

Summit of Mount Roraima, 8600 ft., *McConnell & Quelch*, 631.

PTEROCAULON VIRGATUM, DC. Prod. v. p. 454, var.

Kotinga Valley, *McConnell & Quelch*, 159. Savannahs generally, *McConnell & Quelch*, 306.

STIFFTIA CONDENSATA, Baker, in Mart. Fl. Bras. vi. pt. III. p. 351; in Thurn & Oliver, in Trans. Linn. Soc. Ser. II. ii. (1887), pp. 254, 278.

Summit of Mount Roraima, *McConnell & Quelch*, 87, 651.—Endemic.

STIFFTIA CONNELLII, N. E. Brown, sp. n. (Plate 8.) Rami 2-3 lin. crassi, tomento compacto cinereo dense obtecti. Folia petiolata, coriacea, oblonga vel oblongo-obovata, obtusa, basi cuneato-rotundata, supra glabra, subtus tomento deciduo vestita. Capitula solitaria, magna. Involucris subinfundibuliformis bractearum multiseriatarum; interiores ligulatarum obtusarum; exteriores gradatim minores, ovatarum, obtusarum, subtomentosarum. Receptaculum bracteis linearibus minute dentato-ciliatis paucis onustum. Corolla infra medium 5-loba, glabra, lobis linearibus acutis spiraliter revolutis. Stylus longe exsertus, apice minute bifidus.

Branches woody, 2-3 lin. thick, covered with a dense, felt-like, greyish tomentum. Leaves alternate, petiolate, coriaceous; petiole $\frac{1}{4}$ - $\frac{3}{4}$ in. long, tomentose; blade $3\frac{1}{2}$ - $5\frac{1}{2}$ in. long, $1\frac{1}{2}$ - $2\frac{1}{2}$ in. broad, oblong or obovate-oblong, obtuse, somewhat cuneately rounded at the base, entire, glabrous above, clothed with a deciduous tomentum beneath, which falls away in patches; midrib and veins about equally prominent on both sides. Heads solitary, terminal, about $2\frac{1}{2}$ in. long and 2 in. diam. Involucre somewhat funnel-shaped,

shorter than the florets, with imbricating bracts in 10 or 12 series, the innermost about $1\frac{1}{3}$ in. long, $1\frac{3}{4}$ – $3\frac{1}{2}$ lin. broad, strap-shaped to lanceolate-oblong, obtuse, the outer gradually smaller, ovate, obtuse, tomentose on the back. Receptacle with a few deciduous bracts, scattered among the florets, about $1\frac{1}{4}$ in. long, linear, obtuse, sparsely and minutely denticulate-ciliate. Corolla glabrous, with a tube about $\frac{1}{2}$ in. long, slightly enlarging upwards, and spirally revolute linear-acute lobes, about 8 lin. long, $\frac{1}{2}$ – $\frac{2}{3}$ lin. broad. Anthers about 7 lin. long, with the tails connate in pairs. Ovary $\frac{1}{4}$ in. long, compressed, glabrous. Style much exserted, minutely bifid at the apex. Pappus-bristles copious, unequal, the longest about 10 lin. long, minutely adpressed pubescent, dull straw-coloured, or pale tawny.

Summit of Mount Roraima, 8600 ft., *McConnell & Quelch*, 661.

A very distinct species, readily distinguished by its large heads, and the very long exserted styles. It appears to be a connecting link between *Stiffia* and *Wunderlichia*. The presence of bracts upon the receptacle and the minutely bifid stigma, technically place it in the genus *Wunderlichia*, Riedel, but the plant has a different appearance, and the receptacular bracts are not persistent and not of the same character or scarious texture as those of *Wunderlichia*. In all probability when more species are discovered it will be found that the two genera will have to be united.

QUELCHIA, N. E. Brown, gen. nov.

Capitula 1-flora in glomerulum vel cymam densam aggregata. Involucrum conico-tubulosum, bracteis imbricatis interioribus elongatis, exterioribus gradatim brevioribus ovatis. Receptaculum parvum, nudum. Corolla regularis, profunde 5-loba, tubo brevissimo, lobis linearibus. Anthæræ basi caudato-sagittatæ, caudis retrorsim subbarbatis. Styli rami breves, erecti, truncati. Achænia subteretia, leviter costata. Pappi setæ copiosæ.—Frutex, foliis alternis integris coriaceis. Capitulorum glomeruli pedunculati.

QUELCHIA CONFERTA, N. E. Brown. (Plate 7, figs. 8–14.) Rami apice tomentosi. Folia petiolata, obovata vel oblongo-oblanceolata, obtusa vel apiculata basi cuneato-acuta, primum subtomentosa, demum glabrata. Glomeruli vel corymbi $\frac{3}{4}$ –1 poll. diam. Pedunculi tomentosi. Capitula sessilia, 1-flora, conferta. Involucri bracteæ interiores anguste lanceolatæ, obtusæ, glabræ; exteriores ovatæ, dorso tomentosæ. Corolla glabra. Achænia $2\frac{1}{2}$ lin. longa, glabra. Pappi setæ 3–4 lin. longæ vix vel leviter scabridæ.

Apparently a shrub, densely tomentose on the youngest parts of the shoots. Leaves alternate, coriaceous, petiolate; petiole 2–6 lin. long, tomentose in the young state becoming glabrous; blade 1– $2\frac{1}{2}$ in. long, 5–10 lin. broad, obovate or oblong-oblanceolate, obtuse or more or less apiculate, cuneate-acute at the base, entire, at first with a thin tomentum on both sides, which soon disappears and the leaf becomes glabrous; midrib impressed and pubescent above, very prominent beneath, veins reticulated, impressed above, scarcely prominent beneath. Corymbs pedunculate, small, hemispherical or subglobose, very densely many-headed, $\frac{3}{4}$ –1 in. diam. Peduncles $\frac{1}{2}$ – $1\frac{1}{2}$ in. long,

tomentose, bearing a few small leaf-like bracts at the base of the very short branches of the corymb. Heads sessile, densely crowded, about 4 lin. long, 1-flowered. Involucre cylindric-conical, with 12-16 spirally imbricating scales, the innermost about $3\frac{1}{2}$ lin. long, $\frac{1}{2}$ -1 lin. broad, narrowly lanceolate obtuse, glabrous, the outer gradually shorter and more and more ovate, and more or less tomentose on the back. Floret not exceeding the involucre (?), hermaphrodite. Corolla regular, very deeply 5-lobed, glabrous, the lobes about $2\frac{1}{2}$ lin. long, linear, acute, incurved at the apex. Stamens with free filaments and syngenesious anthers, which are produced at the base into long, slightly barbed tails. Achenes $2\frac{1}{2}$ lin. long, subterete, slightly grooved, glabrous. Pappus-bristles copious, 3-4 lin. long, nearly smooth or very slightly scabrid, pale tawny-white.

Summit of Mount Roraima, 8600 ft., *McConnell & Quelch*, 652.

This plant appears to be nearly allied to the genus *Moquinia*, DC., but differs in habit and in its 1-flowered, densely-crowded heads. I am somewhat doubtful as to the exertion of the floret from the involucre during flowering, as I have only seen the corollas in the bud state; when in fruit the achene is as long as the involucre and the pappus is entirely exerted.

CAMPANULACEÆ.

CENTROPOGON SURINAMENSIS, Presl, Prod. Monog. Lobel. p. 48; in Thurn & Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), pp. 261, 278.

Kotinga Valley, *McConnell & Quelch*, 156.—Widely dispersed throughout Tropical America except on the west of the Andes.

VACCINIACEÆ.

PSAMMISIA CORIACEA, N. E. Brown, sp. n. Frutex glaber. Folia breviter petiolata, obovata, oblonga vel orbiculata, crasse coriacea, integra, infra reticulato-venosa. Flores in axillis aggregati, penduli. Pedicelli recurvi, bibracteolati. Calyx cum pedicello articulatus, limbo breviter 4-5-dentato. Corolla $\frac{1}{2}$ poll. longa, tubulosa, ore angustata, breviter 4-5-loba, lobis erectis lineari-oblongis obtusis. Stamina inclusa, biseriata, libera, glabra, antheris longe bitubulosis. Ovarium subglobosum. Stylus corolla subæquilongus, nec exsertus.

A shrub with a greyish-brown bark. Leaves shortly petiolate, thick and rigidly coriaceous, glabrous; petiole 1-2 lin. long, stout; blade $\frac{3}{4}$ - $1\frac{1}{2}$ in. long, $\frac{1}{2}$ - $1\frac{1}{2}$ in. broad, varying from obovate to orbicular, obtuse, entire, reticulately veined beneath. Flowers in clusters of 3-9 in the axils of the leaves, pedicellate, drooping. Pedicels about 4 lin. long, curved, bibracteolate at the middle or below, glabrous. Bracteoles opposite or subopposite, $\frac{1}{4}$ - $\frac{1}{3}$ lin. long, broadly ovate, obtuse or subapiculate, very minutely ciliate. Calyx about $2\frac{1}{4}$ lin. long, glabrous, jointed to the pedicel, truncate at the base, with a subglobose tube and a cup-shaped shortly 4-5-toothed limb, the teeth very much broader than long, acute or apiculate. Corolla $\frac{1}{2}$ in. long, about 2 lin. diam., tubular, narrowed at the mouth, 4-5-lobed, glabrous inside and outside; lobes 1 lin. long, erect, linear-oblong, obtuse. Stamens 8-10, included, biseriate, glabrous; the shorter filaments $1\frac{1}{3}$ lin. long, the longer $1\frac{3}{4}$ lin. long; anthers $\frac{1}{4}$ in. long, with the cells passing into two

long tubes about 3 times as long as the united part, opening by long oblique slits. Ovary subglobose, 4-5-celled; style $5\frac{1}{2}$ -6 lin. long, not exserted, filiform, glabrous; stigma small, discoid. Ovules numerous in each cell. Berry globose, many-seeded; seeds about $\frac{2}{3}$ lin. long, variable in form, with a pale brown reticulated testa.

Summit of Mount Roraima, 8600 ft., *McConnell & Quelch*, 662.

VACCINIUM RORAIMENSE, N. E. Brown, sp. n. Fruticulus, ramis patente pubescentibus foliosis. Folia parva, breviter petiolata, rigide coriacea, lanceolata obovata vel elliptica, obtusa vel rare acuta, crenato-dentata vel integra, glabra, marginibus subtus incrassata. Racemi breves, terminales, bracteati. Pedicelli bracteolati, pubescentes. Bracteolæ lanceolatae, acutæ, tenuiter glanduloso-ciliatæ. Calyx glaber, lobis deltoideo-ovatis acutis. Corolla $2\frac{3}{4}$ lin. longa, urceolata, breviter 5-loba. Stamina 10, inclusa, filamentis a basi attenuatis ciliatis, antheris acutis nec aristatis. Ovarium obconicum, 5-loculare, glabrum, loculis 2-3-ovulatis.—*Vaccinium*, an *V. floribundum*, H. B. & K.?, Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887) p. 278.

A small branching shrub, probably of dwarf habit. Branches rather slender, very leafy, pubescent with short spreading hairs. Leaves small, petiolate, thick, rigidly coriaceous, glabrous; petiole $\frac{1}{2}$ - $1\frac{1}{4}$ lin. long; blade $3\frac{1}{2}$ -7 lin. long, 2- $5\frac{1}{2}$ lin. broad, varying from lanceolate to elliptic, obtuse or bluntly pointed, rarely acute, crenately toothed or entire, with the margins much thickened beneath, and the veins usually inconspicuous above, more or less evident beneath. Flowers in short terminal bracteate racemes, or aggregated at the apex of the branches. Pedicels $1\frac{1}{2}$ -3, rarely 4-5 lin. long, pubescent, bracteolate. Bracts and bracteoles similar, 1- $1\frac{1}{2}$ lin. long, lanceolate, acute, or occasionally the bracts are larger and resemble very small leaves, glabrous, more or less gland-ciliate. Calyx jointed to the pedicel, with a 5-lobed limb, glabrous; lobes 1 lin. long, deltoid-ovate, acute. Corolla $2\frac{3}{4}$ lin. long, urceolate, 5-lobed, glabrous; lobes about $\frac{3}{4}$ lin. long, broadly ovate, obtuse. Stamens 10, included; filaments 1 lin. long, tapering upwards from a broad base, ciliate; anthers acute but not aristate. Ovary obconical, surmounted by a thick ring-like disk, 5-celled, with 2-3 ovules in each cell, glabrous; style about $1\frac{1}{2}$ lin. long; stigma dilated cushion-like.

Summit of Mount Roraima, 8600 ft., *im Thurn*, 329, 333; *McConnell & Quelch*, 79, 494, 642.

Allied to *V. polystachyum*, Benth., but very different in foliage.

Besides the above, there is a specimen belonging to this family which is too imperfect for identification; it was collected in the Kotinga Valley, *McConnell & Quelch*, 187.

ERICACEÆ.

PERNETTYA MARGINATA, N. E. Brown, sp. n. Frutex foliosus, ramis gracilibus plus minus hirtis. Folia breviter petiolata, coriacea, ovata vel elliptica, acuta vel subacuta, basi rotundata, supra glabra, nitida, infra sæpe parce setulosa, marginibus infra incrassatis setuloso-denticulatis. Flores axillares, solitaires. Pedicelli bracteolati, pubescentes.

Bracteolæ ovatæ, glanduloso-denticulatæ. Calyx glaber, lobis ovatis acutis minute ciliatis. Corolla $3\frac{1}{2}$ lin. longa, urceolata, breviter 5-loba, lobis recurvis. Stamina inclusa, filamentis basi dilatis ciliatis, antheris apicibus 4-aristatis. Bacca globosa, $\frac{1}{4}$ poll. diam.—*P. sp. aff. parvifoliæ*, Benth., Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. p. 278.

A shrub with very leafy slender branches, which are more or less covered with scattered spreading hairs, but are scarcely hispid, and occasionally nearly glabrous. Leaves small, rather crowded, petiolate, rigidly coriaceous; petiole about 1 lin. long; blade 5–8 lin. long, $3\frac{1}{2}$ – $4\frac{1}{2}$ lin. broad, ovate or elliptic, acute or subacute, rounded at the base, thickened and slightly recurved along the denticulate margins beneath, the teeth tipped with a minute bristle, usually with a few scattered bristle-like hairs beneath, otherwise glabrous, shining and rugose from depressions (caused by shrinkage in drying?) between the veins above. Flowers axillary, solitary. Pedicels 2–5 lin. long, more or less pubescent, bracteolate. Bracteoles 5 or 6, about 1 lin. long, ovate, acute, gland-toothed, the lowermost crowded. Calyx 5-lobed nearly to the base, not enlarging in the fruiting stage; lobes $1\frac{1}{2}$ lin. long, $\frac{3}{4}$ –1 lin. broad, ovate, acute, minutely ciliate, otherwise glabrous. Corolla $3\frac{1}{2}$ lin. long, urceolate, glabrous, with 5 ovate obtuse recurved lobes $\frac{3}{4}$ –1 lin. long. Stamens shorter than the corolla, 2 lin. long; filaments ovate-lanceolate, acuminate, ciliate with rather long hairs; anthers tipped with 4 bristles about $\frac{1}{3}$ as long as the cells. Ovary subglobose, glabrous; style $1\frac{1}{2}$ lin. long, glabrous; stigma dilated, discoid. Berry globose, about $\frac{1}{4}$ in. in diam.

Summit of Mount Roraima, 8600 ft., *McConnell & Quelch*, 637; *in Thurn*, 333 A.

Allied to *Pernettya rigida*, DC., but among other characters the more thickened margin of the leaves and more numerous bracteoles serve readily to distinguish it.

GAULTHERIA SETULOSA, N. E. Brown, sp. n. Frutex setulis glanduliferis ubique vestitus.

Folia brevissime petiolata, late cordato-ovata vel suborbicularia, coriacea, integra, utrinque setuloso-scabrida, ciliata. Racemi solitarii vel 2–3-ni ad apices ramorum, bracteati. Calycis lobi deltoideo-ovati, acuminati. Corolla urceolata, breviter 5-loba, intra glabra. Stamina inclusa, filamentis linearibus ciliatis, antheris obtusis muticis. Ovarium depressum, 5-lobum, pubescens.—*Gaultheria aff. G. vestitæ*, Benth., Oliver in Trans. Linn. Soc. Ser. II. Bot. ii. (1887) p. 278.

A shrub, more or less clothed with long setæ or gland-tipped hairs on the branches, leaves, and inflorescence. Leaves very shortly petiolate, $\frac{3}{4}$ – $1\frac{3}{4}$ in. long, $\frac{2}{3}$ – $1\frac{1}{2}$ in. broad, ovate or suborbicular, obtuse or subacute, cordate at the base, entire, rigidly coriaceous, more or less setose or scabrid on both sides, ciliate, with the veins much reticulated and prominent beneath. Racemes solitary or 2–3 together at the apex of the branches, erect, bracteate, 1–4 in. long, 7–15-flowered, rather lax, puberulous and more or less densely covered with gland-tipped bristles on the axis, bracts, pedicels, calyx, and corolla. Bracts $\frac{1}{4}$ – $\frac{1}{2}$ in. long, 2– $3\frac{1}{2}$ lin. broad, elliptic subacute, concave, spreading, persistent. Pedicels 3–8 lin. long, ascending or spreading. Calyx 2– $2\frac{1}{2}$ lin. long, 5-lobed to $\frac{3}{4}$ the way down, the lobes $1\frac{1}{2}$ –2 lin. long, deltoid-ovate, acuminate. Corolla $3\frac{1}{2}$ –4 lin. long, $2\frac{1}{2}$ lin. diam., urceolate, with 5 recurved, ovate, obtuse lobes $\frac{3}{4}$ lin. long, glabrous inside. Stamens 10, included, $\frac{2}{3}$ as long as the corolla; filaments linear, ciliate; anthers $\frac{3}{4}$ lin.

long, oblong, obtuse, shortly bifid at the apex, but not in the least produced into awns or tubes, opening by longitudinal slits nearly to the base. Ovary depressed, 5-lobed, pubescent; style included, rather stout, glabrous; stigma simple, not enlarged. Capsule depressed, 5-angled, pubescent.

Summit of Mount Roraima, 8600 ft., *in Thurn*, 332; *McConnell & Quelch*, 104, 647.

Allied to *Gaultheria cordifolia*, H. B. & K., and *G. vestita*, Benth., but differing from both in the scabrid upper surface of the leaves, the stouter gland-tipped setæ which clothe it, and by the obtuse, not aristate anthers.

LEDOTHAMNUS SESSILIFLORUS, N. E. Brown, sp. n. Fruticulus nanus, ericoideus. Folia verticillata, 3-na vel 4-na ad nodos posita, imbricata, brevissime petiolata, linearia vel lineari-lanceolata, acuta, minute puberula. Flores solitarii, terminales, sessiles. Sepala 7 vel 6, anguste lanceolata, acuta, puberula vel subglabra, ciliata. Petala 7 vel 6, oblonga obtusa, glabra. Stamina 7 vel 6, filamentis quam antheris multo longioribus. Discus nullus. Ovarium subglobosum vel ovoideum, granulosum, 5-8-loculare; stylus crassus; stigma subinfundibuliforme, minute 5-8-lobulatum.

A dwarf shrublet 4-8 in. high. Stems erect, simple or branched, densely leafy to the apex, at first minutely puberulous, becoming glabrous, and rough from the persistent scars of the fallen leaves. Leaves resembling those of an *Erica*, in densely crowded whorls of 3-4, ascending, imbricating, $1\frac{1}{2}$ - $2\frac{1}{2}$ lin. long, $\frac{1}{2}$ - $\frac{2}{3}$ lin. broad, linear or narrowly linear-lanceolate, acute, narrowed at the base into a very short petiole, thick, flat on the face, grooved down the back, minutely puberulous, ciliate or entire. Flowers terminal, solitary, $\frac{3}{4}$ - $1\frac{1}{4}$ in. diam., sessile, surrounded by a few bracts that are intermediate in character between the sepals and leaves. Sepals 7 or occasionally 6, spreading, 3 - $4\frac{1}{2}$ lin. long, $\frac{2}{3}$ - $\frac{3}{4}$ lin. broad, narrowly lanceolate, acute, minutely puberulous or subglabrous, ciliate. Petals usually 7, very spreading, bright crimson, 5-7 lin. long, 2-3 lin. broad, oblong, obtuse, cuneately narrowed at the base, entire, glabrous. Stamens 7 or 6, shorter than the petals, glabrous; filaments $3\frac{1}{2}$ - $4\frac{1}{2}$ lin. long, gradually dilated towards the base; anthers $1\frac{1}{2}$ lin. long, oblong, obtuse. Ovary subglobose or ovoid, granulate-rugose, 5-8-celled, with a stout style $1\frac{1}{4}$ - $1\frac{1}{2}$ lin. long, slightly dilated at the apex into a funnel-shaped minutely 5-8-lobed stigma. Capsule sessile, 2 - $2\frac{1}{2}$ lin. diam., 5-8-valved, granulate.

Summit of Mount Roraima, 8600 ft., *McConnell & Quelch*, 643.

Var. GLABER, N. E. Brown, var. n. Folia et sepala glabra, ciliata.

Leaves and sepals glabrous, ciliate.

L. guyanensis, Meissn., var. *minor*, Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 278, t. 44. f. A.

Mount Roraima, on the upper part of the Ledge and on the summit, *Im Thurn*, 308; *McConnell & Quelch*, 99.

This plant is very distinct from *L. guyanensis*, Meissn., with which it has been associated by Oliver, differing not only by its smaller size and smaller leaves, but more especially by its sessile flowers and the long filaments of the stamens. In *L. guyanensis* the flowers have pedicels $\frac{1}{3}$ - $\frac{2}{3}$ in. long covered with gland-tipped hairs, and the filaments of the stamens are only about 1 lin. long and shorter than the anthers.

BEFARIA GUIANENSIS, Klotzsch, in R. Schomb. Reisen in Brit.-Guiana, iii. p. 1088; Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 278; Appun, Unter den Tropen, ii. pp. 232, 287, 292.

Upper slopes and Ledge of Mount Roraima, *McConnell & Quelch*, 40.—Endemic.

BEFARIA IMTHURNII, N. E. Brown, sp. n. Fruticulus nanus, ramis pilis glanduliferis vel setulis vestitus. Folia conferta, breviter petiolata, coriacea, elliptico-oblonga vel elliptico-lanceolata, utrinque obtusa vel acuta, supra glabra, subtus glauca et glabra vel costa plus minus setulosa, marginibus integris vel glanduloso-scabridis vel ciliatis. Racemi subumbelliformes, densi. Pedicelli subhispidi. Calyx campanulatus, 7-8-lobus, lobis biseriatis late ovatis obtusis. Petala 6-8, oblonga, obtusa. Stamina 12-16, petalis subæquilonga, basi pilosa.—*Befaria* aff. *B. resinosa*, Mutis; Oliver in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 278.

A dwarf shrublet, with the branchlets densely covered with spreading simple or gland-tipped hairs, very leafy. Leaves crowded, very shortly petiolate, coriaceous; petiole $\frac{1}{2}$ -1 lin. long; blade $\frac{1}{2}$ -1 in. long, $\frac{1}{4}$ - $\frac{1}{2}$ in. broad, elliptic-oblong or elliptic-lanceolate, about equally acute or obtuse at each end, with slightly revolute margins, which are either quite entire, minutely gland-scabrid, or ciliate with short gland-tipped bristles or longer simple hairs, glabrous and smooth above, glaucous and either quite glabrous or with some simple or gland-tipped bristles scattered along the midrib beneath. Flowers several, crowded into a short terminal umbel-like raceme or cluster, each flower being solitary in the axil of one of the uppermost leaves. Pedicels $\frac{3}{4}$ -7 lin. long, more or less densely covered with stiff spreading hairs. Calyx about 2 lin. long, campanulate, 7-8-lobed to about the middle, with the lobes in two series, broadly ovate, obtuse, glabrous, minutely ciliate. Corolla glabrous, pink; petals 6-8, slightly spreading, 8-9 lin. long, 3 lin. broad, oblong, obtuse, cuneate at the base. Stamens 12-16, about as long as the petals; filaments filiform, slightly thickened and hairy at the base; anthers 1 lin. long, cuneately oblong, obtuse. Ovary depressed, lobulate, glabrous, with a glabrous style 6-8 lin. long, elongating to 1-1 $\frac{1}{4}$ in. long in fruit, and a capitate slightly lobed stigma. Capsule about $\frac{1}{4}$ in. diam., woody, 6 (or more ?)-valved.

Summit of Mount Roraima, 8600 ft., *im Thurn*, 310; *McConnell & Quelch*, 94, 646.

Allied to *B. guianensis*, Klotzsch (*Schomburgk*, 1041), from which it differs in its less elongated, glabrous leaves. In *B. guianensis* the leaves are 1-1 $\frac{1}{2}$ in. long, and their upper surface is thinly covered with short gland-tipped bristles, tubercular at the base, becoming more or less scabrid after the bristles have fallen or have been rubbed off; on the under side they are more or less densely covered all over with fine gland-tipped hairs, with longer hairs on the midrib, and they appear to be less glaucous than in *B. Imthurnii*.

MYRSINACEÆ.

AEDISIA QUELCHII, N. E. Brown, sp. n. Frutex dioicus, glaber, ramis crassis. Folia alterna, petiolata, coriacea, obovata vel subelliptica, obtusa vel rare subacuta, integra, subtus minutissime rubro-glandulosa. Paniculæ axillares, anguste oblongæ vel

pyramidales, minutissime rubro-glandulosæ, floribus parvis ad apices ramorum confertis. Calyx profunde 4-lobus, glanduloso-punctatus, lobis oblongis obtusis. Corolla profunde 4-loba, glanduloso-punctata, lobis oblongis obtusis patentibus. Stamina 4, corollæ lobis breviora et prope basin inserta, filamentis quam antheris paulo brevioribus.

A dicecious shrub, probably of dwarf habit, glabrous in all parts, with branches $\frac{1}{4}$ in. thick, terete. Leaves chiefly near the apex of the stem, alternate, petiolate, coriaceous; petiole 2-4 lin. long, rather stout; blade $1\frac{1}{2}$ - $2\frac{1}{2}$ in. long, 10-16 lin. broad, obovate or occasionally subelliptic, obtuse, rarely subacute, with revolute margins, entire, more or less densely covered with very minute red glands beneath. Panicles axillary, near the summits of the branches, narrowly oblong or pyramidal, $1-3\frac{1}{4}$ in. long, $\frac{1}{3}$ - $1\frac{1}{4}$ in. broad, covered with minute red glands, with the flowers crowded at the ends of the short, rather distant, simple branches. Bracts $\frac{1}{2}$ -1 lin. long, oblong, obtuse, dotted with rather large dark-coloured glands. Calyx 4-lobed nearly to the base; lobes $\frac{1}{2}$ - $\frac{2}{3}$ lin. long, oblong, obtuse, conspicuously gland-dotted. Corolla about 2 lin. diam., 4-lobed nearly to the base; lobes $\frac{3}{4}$ -1 lin. long, rather more than $\frac{1}{2}$ lin. broad, conspicuously gland-dotted. Stamens 4, shorter than and inserted towards the base of the corolla-lobes, with stout filaments rather shorter than the oblong, $\frac{1}{3}$ lin. long anthers. Ovary rudimentary, minute, subulate, or none. Female flowers not seen.

Summit of Mount Roraima, 8600 ft., *McConnell & Quelch*, 665.

A very distinct species, unlike any other.

APOCYNACEÆ.

MANDEVILLA GLABRA, N. E. Brown, sp. n. Caules volubiles, glabrescentes. Folia parva, distantia, petiolata, coriacea, lanceolata, acuminata, apice obtusa, glabra. Racemi brevissimi, pauciflori, axillares, glabri. Pedunculi et pedicelli brevissimi. Sepala parva, ovata, acuta, glabra. Corolla infundibularis, $2\frac{1}{4}$ poll. longa, extra glabra, intra ad insertionem staminum pilis deflexis dense barbata, lobis dolabri-formibus. Stamina inclusa, filamentis oblongis apice barbatis, antheris acutis, basi cordatis obtusis.

Stem twining, about 1 lin. thick, glabrous or with a very few short scattered hairs, brown. Leaves opposite, distant, petiolate, coriaceous, glabrous; petiole 2-3 lin. long; blade $\frac{3}{4}$ -1 in. long, $3\frac{1}{2}$ -4 lin. broad, lanceolate, acuminate, obtuse at the point, entire, with the midrib impressed above, prominent beneath, and having 5 or 6 small tubercles or glands scattered along it on the upper side, the lateral veins spreading, slightly impressed on both sides in the dried state, not very conspicuous. Racemes axillary, from one axil, very short, about 3-flowered in the specimen seen; glabrous. Peduncle about 1 lin. long, stout. Pedicels alternate, $1-1\frac{1}{2}$ lin. long, stout. Sepals 1 lin. long, $\frac{2}{3}$ - $\frac{3}{4}$ lin. broad, ovate, acute, glabrous, each with a very short, transversely oblong, denticulate scale at its base inside. Corolla about $2\frac{1}{4}$ in. long, funnel-shaped; tube curved, cylindrical and about $1\frac{1}{2}$ lin. diam. in the lower half, much enlarged and sub-campanulate in the upper half, glabrous outside and inside, except 5 broad lines that are

densely bearded with deflexed white hairs at the insertion of the stamens within; lobes about $\frac{3}{4}$ in. long, hatchet-shaped, one side being dilated into a large square membranous obtuse lobe. Stamens inserted at the top of the narrow part of the tube included; filaments about 1 lin. long and nearly as broad, flat, oblong, slightly bearded near the apex on the inner face; anthers $3\frac{1}{3}$ lin. long, linear-oblong, acute, shortly cordate at the base, connate, adnate to the stigma. Disk shortly lobed. Ovary glabrous; style about 14 lin. long, slender, glabrous; stigma conical, 5-winged.

Upper slopes of Mount Roraima, *McConnell & Quelch*, 16.

MANDEVILLA LINEARIS, N. E. Brown, sp. n. Caules volubiles, graciles, puberuli. Folia breviter petiolata, linearia, acuta, supra glabra, subtus tomentosa, marginibus fere vel usque ad costam revolutis. Racemi breves, pauciflori, minute puberuli. Bracteae parvae, acuminatae. Pedicelli brevissimi. Sepala parva, deltoidea, acuminata. Corolla infundibularis, $2\frac{1}{4}$ poll. longa, recta, extra glabra, intra ad mediam dense barbata. Stamina inclusa, filamentis ellipticis glabris, antheris subacutis basi breviter cordatis vel emarginatis.

Stem twining, slender, puberulous. Leaves opposite, $\frac{3}{4}$ – $2\frac{1}{2}$ in. distant, petiolate, thinly coriaceous; petiole 1– $1\frac{1}{2}$ lin. long, rather slender, puberulous; blade $1\frac{1}{2}$ – $2\frac{1}{4}$ in. long, linear, acute, obtuse at the base, with the margins revolute nearly or quite to the midrib, glabrous and shining, with an impressed midrib above, tomentose, with a prominent midrib beneath, the tomentum usually hidden by the revolute margins. Racemes short, $\frac{1}{2}$ – $\frac{3}{4}$ in. long, more than half of which is peduncle, 3–4-flowered, very minutely puberulous. Bracts about $\frac{3}{4}$ lin. long, ovate, acuminate. Pedicels alternate, about 1 lin. long, rather stout. Calyx-lobes about $1\frac{1}{4}$ lin. long, deltoid, acuminate, subglabrous or very minutely puberulous, each with a small ovate scale at the base inside. Corolla about $2\frac{1}{4}$ in. long, funnel-shaped, straight, 5-lobed, glabrous outside, the middle part of the tube inside densely covered with deflexed white hairs; tube cylindric and about $1\frac{1}{2}$ lin. diam. for $\frac{2}{3}$ of its length, upper part much enlarged, subcampanulate; lobes about 5 lin. long, hatchet-shaped, with an incurved apiculus, one side being dilated into a subrectangular, obtuse, membranous lobe. Stamens 5, included, inserted at the top of the narrow part of the tube; filaments about $\frac{3}{4}$ lin. long, elliptic, glabrous; anthers 2 lin. long, linear-oblong, subacute, slightly narrowed at the emarginate or shortly cordate base. Disk of 5 oblong lobes. Ovary glabrous; style about $1\frac{1}{4}$ in. long, filiform, glabrous; stigma subquadrate when viewed sideways, obtuse, 5-winged.

Kotinga Valley, 3000 ft., *McConnell & Quelch*, 132, 194.

MANDEVILLA SCABERULA, N. E. Brown, sp. n. Caules volubiles, puberuli. Folia distantia, petiolata, oblonga, obtusa, apiculata, basi cordata, supra scaberula, subtus tomentosa, reticulato-venosa. Racemi axillares, multiflori, puberuli. Bracteae oblongae vel oblongo-lanceolatae. Pedicelli brevi. Calycis lobi ovati, acuti, intra squama ovata instructi. Corolla 2 poll. longa, infundibularis, extra pubescens, intra ad insertionem staminum pilis deflexis dense barbata, tubo curvato inferne angusto basi leviter inflato, superne ampliato subcampanulato, lobis dolabriformibus. Stamina inclusa, filamentis linearibus dense barbatis, antheris acutis basi cordatis obtusis.

Stem twining, $1\frac{1}{4}$ – $1\frac{1}{3}$ lin. thick, puberulous. Leaves opposite, very distant, $2\frac{1}{2}$ – $3\frac{1}{2}$ in. long, $1\frac{1}{4}$ –2 in. broad, oblong, obtuse, apiculate, cordate at the base, scaberulous above, finely greyish-tomentose beneath, reticulated with dark veins; petiole $1\frac{1}{2}$ –2 lin. long, puberulous. Racemes axillary, from one axil, 2–3 in. long, including the $\frac{3}{4}$ in. long peduncle, many-flowered, densely puberulous in all parts. Bracts about 2 lin. long. $1\frac{2}{3}$ – $1\frac{1}{2}$ lin. broad, oblong or oblong-lanceolate, acute, thin, spreading, deciduous. Pedicels alternate, 2–3 lin. long, moderately stout. Calyx-lobes $1\frac{1}{3}$ lin. long and nearly as broad, ovate, acute, each with an ovate acute slightly toothed scale at its base inside. Corolla about 2 in. long, funnel-shaped, curved, 5-lobed, pubescent outside, densely bearded with deflexed white hairs at the insertion of the stamens within, elsewhere glabrous; the lower half of the tube is slightly inflated at its base and there about $2\frac{1}{2}$ lin. diam., narrowed above to $1\frac{1}{2}$ lin. diam.; the upper half much enlarged, funnel-shaped or subcampanulate; lobes about $\frac{3}{4}$ in. long, hatchet-shaped, one side being dilated into a large subquadrate, obtusely rounded, membranous lobe. Stamens 5, included, inserted at the top of the narrow part of the tube; filaments short, flat, linear, densely bearded on the inner face; anthers $2\frac{1}{2}$ lin. long, linear-oblong, acute, shortly cordate at the base, connate, adnate to the stigma. Disk tubular, $\frac{3}{4}$ lin. deep, 5-crenate. Ovary pubescent; style about $1\frac{1}{4}$ in. long, very slender, glabrous; stigma conical, 5-winged.

Tolimbaru Creek near Mount Roraima, *McConnell & Quelch*, 146.

ASCLEPIADACEÆ.

DITASSA TAXIFOLIA, Decne. in DC. Prod. viii. p. 578.

Kotinga Valley, *McConnell & Quelch*, 165, 166. Ireng Valley, *McConnell & Quelch*, 323.

LOGANIACEÆ.

BONYUNIA MINOR, N. E. Brown, sp. n. (Plate 9, figs. 1–5.) Frutex glaber. Folia opposita, approximata, breviter petiolata, coriacea, rotundato-ovata, subacuta, basi rotundata vel leviter cordata. Cymæ subdensæ, ramulis puberulis, floribus sessilibus bracteolatis. Calyx campanulatus, minute puberulus, dentibus deltoideis subacutis. Corollæ tubus subcylindricus, extra pubescens, intra parte inferiore puberulus, superiore glaber; lobi patentes, recurvi, superne carinato-incrassati, superne etiam carina et marginibus puberuli, area basali triangulare glabri. Ovarium breviter pilosum, stylo pubescente, stigmatе bilobo.

A shrub. Branches terete, glabrous, dark brown. Leaves opposite, subimbricate, ascending, shortly petiolate, coriaceous, glabrous; petiole 1 lin. long; blade $\frac{2}{3}$ – $1\frac{1}{4}$ in. long, $\frac{1}{2}$ –1 in. broad, roundish-ovate, subacute, broadly rounded or slightly cordate at the base, with prominent veins beneath. Cymes terminal, rather dense, 1 – $1\frac{1}{4}$ in. diam., with puberulous branches 2–4 lin. long. Flowers sessile, bracteolate. Bracteoles $\frac{2}{3}$ –1 lin. long, linear or oblong-lanceolate, obtuse, minutely puberulous. Calyx about $1\frac{1}{2}$ lin. long, campanulate, puberulous, with 5 deltoid subacute teeth. Corolla tubular, with 5 spreading recurved lobes; tube $3\frac{1}{2}$ lin. long, subcylindric, pubescent outside, puberulous within in the lower $\frac{2}{3}$, glabrous above; lobes about $2\frac{1}{2}$ lin. long, spreading,

recurved, thickened and keeled in the upper half and along the margins below and there densely puberulous, enclosing a narrowly triangular glabrous basal area. Stamens 5, inserted at the mouth of the tube, glabrous; filaments $\frac{1}{4}$ lin. long, slender; anthers about 1 lin. long, partly exserted, linear, acute. Ovary shortly hairy; style $1\frac{2}{3}$ – $1\frac{3}{4}$ lin. long, pubescent; stigma 2-lobed. Capsule 4–6 lin. long, $2\frac{1}{2}$ –3 lin. diam., ellipsoidal, sparingly pubescent or nearly glabrous, 2-valved.

Kotinga Valley, *McConnell & Quelch*, 161. Ireng Valley, *McConnell & Quelch*, 331.

ANTONIA OVATA, Pohl, Pl. Bras. ii. p. 13, t. 109.

Kotinga Valley, *McConnell & Quelch*, 164. Upper slopes of Mount Roraima, *McConnell & Quelch*, 324.—Only known from Brazil and Guiana.

GENTIANACEÆ.

SCHULTESIA BRACHYPTERA, Cham. in Linnæa, viii. (1833), p. 8.

Kotinga Valley, *McConnell & Quelch*, 148.

SCHULTESIA HETEROPHYLLA, Miq. in Linnæa, xix. (1847), p. 136.

Ireng Valley, *McConnell & Quelch*, 249.

SCHULTESIA BENTHAMIANA, Klotzsch, ex Griseb. in Linnæa, xxii. (1849), p. 34.

Ireng Valley, *McConnell & Quelch*, 247 (and 224, apparently a starved state).

COUTOUBEA REFLEXA, Benth. in Ann. Nat. Hist. ii. (1839), p. 442.

Kotinga Valley, *McConnell & Quelch*, 141.

COUTOUBEA SPICATA, Aubl. Pl. Guian. i. p. 72.

Ireng Valley, *McConnell & Quelch*, 221, 259.

LISIANTHUS IMTHURNIANUS, Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 279.

Summit of Mount Roraima, *McConnell & Quelch*, 101, 682.—Endemic.

LISIANTHUS ULIGINOSUS, Griseb. Gen. et Sp. Gent. p. 181.

Kotinga Valley, *McConnell & Quelch*, 172. Ireng Valley, 204, 252. Upper slopes of Mount Roraima (a small-flowered form), *McConnell & Quelch*, 15.—Widely dispersed in Guiana and Brazil.

LISIANTHUS QUELCHII, N. E. Brown, sp. n. (Plate 9, figs. 6–9.) Fruticulus glaber, ramis tetragonis internodiis $1\frac{1}{2}$ –6 lin. longis. Folia opposita, petiolata, oblonga vel elliptico-oblonga vel obovata, subacuta vel obtusa et apiculata, coriacea. Flores 1–3-ni ad apices ramorum, pedicellati. Calyx campanulatus, profunde 5-lobus, lobis oblongis obtusis. Corolla 1– $1\frac{1}{2}$ poll. longa, subinfundibularis, curvata, lobis rotundatis obtusis. Stamina inclusa, filamentis apice recurvis, antheris erectis. Ovarium ovoideum, glabrum, in stylum elongatum angustatum, stigmate bilobo.

A dwarf shrub, glabrous in all parts. Branches 4-angled, 1– $1\frac{1}{2}$ lin. thick, with internodes $1\frac{1}{2}$ –6 lin. long, leafy at the summit only, minutely tuberculate-rugulose.

Leaves opposite, petiolate, coriaceous; petiole 1-2½ lin. long; blade ¾-1¾ in. long, ⅓-1 in. broad, oblong, elliptic or obovate, subacute or obtuse and apiculate, more or less acute at the base, slightly revolute along the margins, with the midrib impressed above, prominent beneath, and the veins invisible. Flowers terminal, solitary or in a sessile 3- (or more?) flowered cyme. Pedicels 2-4 lin. long, rather stout, slightly rough from minute tubercles. Bracts and bracteoles ⅓-1 lin. long, acuminate. Calyx 3½-4 lin. long, campanulate, deeply 5-lobed; lobes 2½-3 lin. long, oblong, obtuse. Corolla 1-1½ in. long, somewhat funnel-shaped, 5-lobed, curved or oblique; lobes 3½-4½ lin. long and broad, suborbicular, contracted at the base. Stamens included; filaments filiform, recurved near the apex and then shortly curved upwards at the very apex; anthers erect, 2-2½ lin. long, oblong subobtuse, cordate at the base. Ovary ovoid, narrowed into the 5 lin. long style, with a shortly 2-lobed stigma.

Summit of Mount Roraima, 8600 ft., *McConnell & Quelch*, 106, 649.

A very distinct species, perhaps nearer to *Lisianthus ovalis*, Ruiz & Pav., than to any other.

LISIANTHUS ELISABETHÆ, Griseb. in *Linnæa*, xxii. (1849), p. 40.

Lisianthus aff. *L. macrantho*, Oliver, in *Trans. Linn. Soc. Ser. II. Bot. ii.* (1887), p. 279.

Leiothamnus Elisabethæ, R. H. Schomb. in *Verh. des Ver. Beförd. Gartenb. in Preuss.* xviii. (1847), p. 155, t. 1; R. Schomb. *Bot. Rem. Brit. Guiana*, p. 77.

Upper slopes and ledge of Mount Roraima, *im Thurn*, 188; *McConnell & Quelch*, 25.—Endemic.

HYDROPHYLLACEÆ.

HYDROLEA SPINOSA, Linn. *Sp. Pl.* ed. 1, p. 328.

Ireng Valley, *McConnell & Quelch*, 262. Kotinga Valley, *McConnell & Quelch*, 180. Savannahs generally, *McConnell & Quelch*, 313.

BORAGINACEÆ.

HELIOTROPIUM STRICTISSIMUM, Moric.?, *Pl. Nouv. Am.* p. 146, t. 87.

Ireng Valley, *McConnell & Quelch*, 220, 302.

CONVOLVULACEÆ.

IPOMÆA IRENGANA, N. E. Brown, sp. n. Caulis gracilis, volubilis, subpubescens, internodiis ⅓-1 poll. longis. Folia parva, petiolata, late cordata, obtusa, minute apiculata, supra velutina, subtus albo-tomentosa. Flores axillares, solitarii, pedicellati. Calycis lobi oblongi, obtusi, puberuli. Corolla 1¼-1½ poll. longa, extra pubescens, purpurea.

Stems very slender, twining, woody below, slightly pubescent, with internodes ⅓-1 in. long. Leaves small; petiole 2-5 lin. long, slender, terete, pubescent; blade ⅓-1 in. long, ⅓-¾ in. broad, cordate or roundish-cordate, obtuse, minutely apiculate, velvety-tomentose above, whitish-tomentose beneath. Flowers axillary, solitary. Pedicels 2-2½ lin. long, very minutely bracteolate at the base, pubescent. Sepals subequal, 3½-4 lin. long, 1½ lin. broad, oblong, obtuse, thinly pubescent or puberulous. Corolla 1¼-1½ in. long,

funnel-shaped, pubescent outside, purple. Stamens included; filaments pubescent at their base. Ovary and style glabrous; stigma of 2 globose lobes.

Ireng Valley, on ant-hills, *McConnell & Quelch*, 251, 265.

JACQUEMONTIA EVOLVULOIDES, Meissn. in Mart. Fl. Bras. vii. p. 307.

Ireng Valley, *McConnell & Quelch*, 233.

EVOLVULUS STRICTUS, Benth. in Hook. Lond. Journ. Bot. v. (1846), p. 354.

Ireng Valley, *McConnell & Quelch*, 218, 219, 261.

EVOLVULUS SERICEUS, Sw. Prod. Veg. Ind. Occ. p. 55.

Ireng Valley, *McConnell & Quelch*, 215.

SOLANACEÆ.

SOLANUM CRINITUM, Lam. Illust. ii. p. 20.

Ireng Valley, near Mataruka Mountain, *McConnell & Quelch*, 134.

SCROPHULARIACEÆ.

BEYRICHIA SCUTELLARIOIDES, Benth. Scroph. Ind. p. 9, in nota.

B. ocymoides, Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1889), p. 280, not of Cham.

Upper slopes of Mount Roraima, *McConnell & Quelch*, 5.—Trinidad and Brazil.

HERPESTIS GRATIOLOIDES, Benth. in Hook. Comp. Bot. Mag. ii. (1836), p. 57.

Ireng Valley, *McConnell & Quelch*, 266.

GERARDIA HISPIDULA, Mart. Nov. Gen. & Sp. iii. p. 13, t. 207.

Ireng Valley, *McConnell & Quelch*, 232.

BUCHNERA PALUSTRIS, Spreng. Syst. ii. p. 805.

Ireng Valley, *McConnell & Quelch*, 234.

BUCHNERA, sp. Too imperfect for determination.

Ireng Valley, *McConnell & Quelch*, 236, 238.

LENTIBULARIACEÆ.

UTRICULARIA HUMBOLDTII, R. H. Schomb. in Verh. des Ver. Beförd. Gartenb. in Preuss. xv. (1841), p. 139, t. 3; R. Schomb. Reisen in Brit.-Guiana, ii. p. 263, & iii. p. 1086; im Thurn & Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1889), pp. 256, 262, 265, 267, 269, & 280.

Upper slopes of Mount Roraima, *McConnell & Quelch*, 3.—Also on the Kaieteur Savannah.

UTRICULARIA ALPINA, Jacq. Enum. Pl. Carib. p. 11.

U. montana, Jacq. Select. Stirp. Amer. p. 7, t. 6.

Upper slopes of Mount Roraima, growing on trees, *McConnell & Quelch*, 43. Roraima

Range, 3500 ft., *McConnell & Quelch*, 718.—Also in the West Indies, Venezuela, Colombia, Peru, and Ecuador.

UTRICULARIA CAMPBELLIANUM, Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 280, t. 44. f. B, & pp. 266, 267, & 270.

Upper slopes of Mount Roraima, on trees, *McConnell & Quelch*, 35.—Endemic.

UTRICULARIA QUELCHII, N. E. Brown, sp. n. (Plate 10, figs. 12–16.) Perennis. Folia spathulata, longe petiolata, coriacea, glabra. Caulis 2–5 poll. altus, 1–2-squamosus, 1–2-florus, glaber. Bracteæ et bracteolæ 2–3½ lin. longæ, oblongæ vel elliptico-oblongæ, obtusæ, glabræ. Sepala magna, elliptica, obtusa. Corolla magna, labio superiore ovato et obtuso et apice recurvo, labio inferiore multo majore orbiculare reflexo, calcare ad medium abrupte procurvo subobtusum.—*Utricularia* aff. *U. montana*, Jacq.; Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 280.

Perennial. Roots fibrous and tuberous intermixed. Leaves all radical, erect, spathulate, coriaceous, glabrous; petiole $\frac{3}{4}$ –2 in. long; blade $\frac{1}{2}$ –1 in. long, 3–5 lin. broad, obovate or narrowly elliptic, obtuse, cuneately narrowed into the petiole. Stem 2–5 in. long, $\frac{1}{2}$ – $\frac{2}{3}$ lin. thick, 1–2-flowered, glabrous, bearing 1–2 lanceolate, acute, glabrous scales 1–2½ lin. long. Bracts and bracteoles subequal, basifixed, 2–3½ lin. long, 1–1½ lin. broad, oblong or elliptic-oblong, obtuse, glabrous, not ciliate. Pedicels 4–6 lin. long, glabrous. Sepals 4½–6 lin. long, 3½–4 lin. broad, elliptic, obtuse, concave, glabrous. Corolla large, glabrous, purple or deep red; upper lip 3–4 lin. long, included in the dorsal sepal, erect, recurved at the apex, ovate, obtuse, with the sides abruptly reflexed around the mouth, then again curved forwards; lower lip 7–9 lin. long, 8–15 lin. broad, abruptly reflexed from the mouth of the spur, suborbicular, very obtuse, entire; spur $\frac{3}{4}$ –1 in. long, gradually tapering from the 2½–3 lin. broad mouth to the obtuse apex, with the basal half hanging straight down and partly embraced by the lower sepal, and the apical half abruptly curved forwards or upwards. Stamens with thick clavate curved filaments, glabrous. Ovary globose, narrowed into a stout style, glabrous; upper lip of the stigma narrow, lower lip very broad, very obtusely rounded or subtruncate.

Summit of Mount Roraima, 8600 ft., *McConnell & Quelch*, 105, 683; *in Thurn*, 293.

Allied to *U. montana*, Jacq., but smaller, with larger petioles to the leaves, broader bracts, and differently coloured flowers, which, to judge from the dried specimens, appear to have been either purple or deep red. Probably it is the plant mentioned by Mr. Quelch in 'Timehri,' vol. ix. p. 178, as "the beautiful crimson *Utricularia montana*," since the flowers of the true *U. montana* (= *U. alpina*), Jacq. are white.

UTRICULARIA CONNELLII, N. E. Brown, sp. n. (Plate 10, figs. 1–6.) Aphylla, 1½–4 poll. alta. Caulis filiformis, 1–2-squamosus, glaber, 1–3-florus. Bracteæ et bracteolæ minutæ, ovatæ vel oblongæ, ciliatæ. Sepala elliptica, multi-nervosa, nervis pubescentibus; sepalum inferius minutum, bilobum. Corolla ringens, glabra, labio superiore erecto, anguste subspathulato-oblongo, labio inferiore suborbiculare obtuso reflexo, calcare leviter curvato subacuto.

Leafless at the time of flowering. Stems $1\frac{1}{2}$ –4 in. high, filiform, glabrous, 1–2-flowered and bearing 1–2 minute, peltately attached, very minutely ciliate scales. Bracts $\frac{1}{2}$ lin. long, ovate, acute, peltately attached, minutely ciliate. Bracteoles slightly shorter than the bracts, basifixed, oblong, acute, minutely ciliate. Pedicels 1–2 lin. long, glabrous. Sepals about $1\frac{1}{2}$ lin. long, the upper nearly 1 lin. broad, ovate, subacute, the lower rather broader, emarginate at the apex, both deeply concave, strongly nerved, with a few short hairs scattered along the nerves. Corolla ringent; upper lip erect, $3-3\frac{1}{2}$ lin. long, $\frac{3}{4}$ –1 lin. broad, subspathulate-oblong, obtuse, with recurved margins, minutely puberulous inside at the base; lower lip 4–5 lin. long, about 4 lin. broad, suborbicular, obtuse, abruptly deflexed from the mouth of the spur, where it is minutely puberulous. Spur $4\frac{1}{2}$ – $5\frac{1}{2}$ lin. long, nearly in a line with the upper lip, slightly curved forwards, gradually tapering from the $1\frac{1}{2}$ lin. broad base to the subacute apex, compressed. Filaments of the stamens incurved, clavate, glabrous. Ovary ovoid, narrowed into a short style, glabrous; stigma with a narrow acute upper lip and a broadly ovate obtuse lower lip.

Arabapu River, *McConnell & Quelch*, 127.

Allied to *U. cornuta*, Michx., but smaller, more slender, and with a more reflexed lower lip to the corolla.

UTRICULARIA RORAIMENSIS, N. E. Brown, sp. n. (Plate 11, figs. 1–4.) Perennis. Folia radicalia, orbiculato-spathulata, 3–7 lin. longa. Caules simplices vel divisi, $1\frac{1}{4}$ –3 poll. longi, subfiliformi, 1–2-flori, 1–3-squamosi, glabri. Bracteae minutae, trifidae. Flores parvi, pedicellati. Sepala inaequalia, glabra; sepalum superius ellipticum, obtusum, profunde concavum; inferius brevius, subquadratum, truncatum. Corollae labium posticum ovatum, obtusum, intra minute glandulosum; labium anticum late subreniforme breviter 3-lobum, abrupte reflexum, minute glandulosum; calcar late conicum, obtusum, minute glandulosum.

Perennial. Leaves all radical, spathulate, glabrous; petiole 2–5 lin. long, slender; blade 1–2 lin. long, $\frac{1}{2}$ – $1\frac{3}{4}$ lin. broad, obovate or orbicular, obtuse, cuneate at the base. Stems 1–4 to a plant, simple or once branched, $1\frac{1}{4}$ –3 in. high, subfiliform, 1–2-flowered, glabrous, bearing 1–3 minute, oblong, obtuse, glabrous scales. Bracts minute, trifid, with the lateral teeth narrow and acute, and the middle tooth broadly rounded, glabrous, not ciliate. Pedicels $1-2\frac{1}{2}$ lin. long, filiform, erect. Sepals unequal, thin, glabrous; the upper $\frac{3}{4}$ –1 lin. long, elliptic or elliptic-ovate, obtuse, deeply concave, with about 4 rather obscure veins; the lower $\frac{1}{2}$ lin. long, and about the same in breadth, subquadrate, truncate or very slightly emarginate, 4-veined. Corolla small; upper lip $1-1\frac{1}{4}$ lin. long, $\frac{2}{3}$ – $\frac{3}{4}$ lin. broad, ovate, obtuse, concave, erect or ascending, minutely glandular within; lower lip $1\frac{1}{3}$ – $1\frac{1}{2}$ lin. long, 2 lin. broad, subreniform, shortly 3-lobed, and more or less undulated, abruptly reflexed from the mouth of the spur on which it rests, minutely glandular on the lower part; spur $1\frac{1}{2}$ lin. long, $1\frac{1}{4}$ lin. broad at the mouth, broadly conical, obtuse, compressed, minutely glandular, directed at a right angle to the pedicel. Stamens with curved clavate filaments, glabrous. Ovary globose, abruptly contracted into a stout style, glabrous; stigma with a minute tooth-like upper lip and a large truncate lower lip.

Summit of Mount Roraima, 8600 ft., *McConnell & Quelch*, 685.

UTRICULARIA CONCINNA, N. E. Brown, sp. n. (Plate 10, figs. 7-11.) Pusilla, glabra.

Pedunculus uniflorus, $\frac{3}{4}$ -1 poll. longus. Bracteæ et bracteolæ lanceolatæ, obtusæ. Sepala magna, cordato-ovata. Corolla erecta, labiis calyce paulo longioribus, glabris, labio superiore elliptico vel subrectangulare et apice obscure 3-crenato, labio inferiore elliptico vel subrectangulare undulato-crenato; calcar labiis multo longius, erectum, acutum, prope apicem parcissime pilosulum.

Leaves 2-4, all radical, 3-5 lin. long, lanceolate, elliptic or oblong-lanceolate, obtuse or subacute, tapering into a short petiole at the base, glabrous, erect. Peduncle $\frac{3}{4}$ -1 in. high, 1-flowered, slender, with two distant minute bracts, one near the middle, the other near the base, glabrous. Flowering-bract $\frac{2}{3}$ -1 $\frac{1}{2}$ lin. long, lanceolate, subobtuse, glabrous, not ciliate. Bracteoles resembling the flowering-bract, but shorter and narrower. Pedicel 2-3 lin. long, glabrous. Sepals 2, erect, large in proportion to the corolla, 2 $\frac{1}{2}$ -3 $\frac{1}{2}$ lin. long, 2-2 $\frac{1}{2}$ lin. broad, cordate-ovate, obtuse, glabrous, not ciliate. Corolla erect, with the lips only slightly exceeding the sepals, and the spur about twice as long; upper lip 1 $\frac{3}{4}$ -2 lin. long, and about the same in breadth, elliptic or subrectangular, obscurely 3-crenate at the apex, with recurved margins, glabrous; lower lip about 2 $\frac{1}{2}$ -2 $\frac{3}{4}$ lin. long, elliptic or subrectangular, crenate and wavy at the apex, with the margins recurved, glabrous; spur about 3 $\frac{1}{2}$ -3 $\frac{3}{4}$ lin. long, erect, acute, with a few scattered hairs on the terminal half. Stamens glabrous. Ovary globose-ovoid, narrowed into a short stout style, glabrous; stigma with a narrow acute upper lip and a broad emarginate lower lip. Capsule about half as long as the sepals, ellipsoidal, obtuse, crowned with the short style and stigma.

Kaieeteur Savannah, *Jenman*, 1272; Mazaruni River, 300 ft., *McConnell & Quelch*, 710; and without precise locality, *Schomburgk*, 131.

The flowers of this diminutive species are stated by Messrs. McConnell and Quelch to be "white with purple points." I have only seen one corolla, and am uncertain if my description of it is quite correct in all details, as it was very much flattened; but my drawings of it (figs. 8 & 9) represent its general appearance as nearly as the material would allow, although they may not be accurate. The affinity of *U. concinna* is with *U. Campbelliana*, Oliver, and *U. montana*, Jacq., although it is very much smaller than either of those species.

UTRICULARIA NERVOSA, Weber?, ex Benj. in Mart. Fl. Bras. x. p. 247.

Upper slopes of Mount Roraima, *McConnell & Quelch*, 3 a.

Possibly a distinct species, but the material consists of one flowering stem only, which scarcely admits of proper dissection; externally, however, it much resembles *U. nervosa*, which is a Brazilian species.

UTRICULARIA sp.

Arabapu River, *McConnell & Quelch*, 150 a.

The specimen consists of leafy rhizomes without flowers, which were found entangled amongst the pitchers and false roots of *Gentlisea guianensis*, N. E. Brown.

GENLISEA GUIANENSIS, N. E. Brown, in Hooker, Icones Plant. t. 2629.

Arabapu River, *McConnell & Quelch*, 150.

This is the largest species of this genus at present discovered, and more nearly resembles *G. africana*, Oliver, (a native of Angola), than any other with which I am acquainted. The members of the genus *Genlisea* appear to be quite destitute of true roots, their place being supplied by modified leaves, which descend into the soil or water. Some of these modified leaves terminate in the curious tubular two-lobed utricles characteristic of the genus, others being quite simple and root-like; probably all fulfil the functions of roots, and the utricles supply an additional amount of nitrogen to the plants by the absorption of the decomposed remains of the minute animalculæ they capture. The utricles have a very remarkable structure, which will be found well described and illustrated in Darwin's 'Insectivorous Plants,' p. 446, and Goebel's 'Pflanzenbiologische Schilderungen,' ii. p. 121, t. 15-16.

GENLISEA RORAIMENSIS, N. E. Brown, sp. n. (Plate 11, figs. 5-12.) Folia obovato-vel rotundato-spathulata, crassiuscula vel subcoriacea, glabra. Utriculi bifformes, alter apice bilobus, alter apice acutus, minute 1-porosi. Caulis $1\frac{1}{2}$ -3 poll. longus, 3-6-squamosus, minute glanduloso-hirtellus, 1-4-florus. Bracteæ et bracteolæ minutæ, lanceolatae, acutæ. Pedicelli minute glanduloso-hirtelli. Calyx 5-lobus, minute pubescens. Corolla parva, lutea, labio superiore late ovato vel elliptico-ovato obtuso concavo glabro, labio inferiore reflexo obtuse trilobo glabro, calcari late conico obtusissimo minute glanduloso-hirtello.

A perennial with a very short branching rhizome densely covered with leaves. Leaves rosulate, 3-5 lin. long, $\frac{1}{2}$ - $\frac{3}{4}$ lin. broad, obovate- or orbicular-spathulate, rather thick or subcoriaceous, glabrous. Utricles $2\frac{1}{2}$ -5 lin. long, of two forms, both tubular and ovoid-inflated at the base and descending into the soil among the roots, one (the perfect form) dividing at the mouth into two long twisted lobes, the other (a transition form) acute, with a very minute orifice at the apex and entirely without lobes; a third and imperfect form is sometimes present standing erect among the leaves in which the terminal lobes are very short and scarcely or not at all twisted. Stem simple, $1\frac{1}{2}$ -3 in. high, $\frac{1}{4}$ - $\frac{1}{3}$ lin. thick, 1-4-flowered, minutely and rather sparsely glandular-hairy, bearing 3-6 lanceolate, acute, basifixed, glabrous scales about $\frac{2}{3}$ lin. long. Bracts and bracteoles subequal, basifixed, about $\frac{3}{4}$ lin. long, lanceolate, acute, sparsely pubescent and ciliate. Pedicels 1-2 lin. long, glandular-pubescent. Calyx 5-lobed almost to the base; lobes $\frac{3}{4}$ -1 lin. long, oblong-lanceolate, acute, thinly pubescent and slightly ciliate. Corolla small, yellow; upper lip $1\frac{1}{2}$ lin. long, 1 lin. broad, broadly ovate or elliptic-ovate, obtuse, concave, glabrous; lower lip 2 lin. long and as much in breadth, reflexed, obtusely 3-lobed, glabrous; spur $1\frac{1}{4}$ lin. long, $1\frac{1}{4}$ lin. broad at the mouth, stout, conical, very obtuse, compressed, slightly and minutely glandular-hairy, and darker in colour than the rest of the flower. Stamens glabrous; filaments curved, clavate; anthers deep blue. Ovary globose, thinly covered with very minute hairs; style short; stigma oblique. Capsule globose, about 1 lin. diam., minutely and sparsely hairy.

Summit of Mount Roraima, 8600 feet, growing in a somewhat sandy boggy soil, *McConnell & Quelch*, 684.

BIGNONIACEÆ.

TABEBUIA RORAIMÆ, Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 280, t. 45.

Upper slopes of Mount Roraima, *McConnell & Quelch*, 33.—Endemic.

ACANTHACEÆ.

RUELLIA VINDEK, Mart. ex Nees in Mart. Fl. Bras. ix. p. 42.

Ireng Valley, *McConnell & Quelch*, 253.

THYRSACANTHUS SCHOMBURGKIANUS, Nees, in Hook. Lond. Journ. Bot. iv. (1845), p. 636.

Upper Essequibo River, *McConnell & Quelch*, 277.

DIANTHERA GUIANENSIS, N. E. Brown, sp. n. Caulis bifariam pubescens. Folia subsessilia, lanceolata vel ovato-lanceolata, acuta, basi angustata rotundata, utrinque glabra vel sparsim pubescentia. Thyrsus terminalis, subspiciformis, pedunculatus. Bracteæ et bracteolæ lineari-lanceolatæ, acutissimæ, subglabræ, ciliatæ. Sepala 5, libera, lineari-lanceolata, acutissima, pubescentia, ciliata. Corolla superne pubescens, labio superiore oblongo marginibus reflexo apice minute bilobo, labio inferiore 3-lobo disco plicato-elevato.—*Justicia* sp., Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 280.

Shrubby, with terete, bifariously pubescent branches. Leaves spreading, subsessile, 2–4 in. long, 7–18 lin. broad, lanceolate or ovate-lanceolate, acute, narrowed below the middle to a rounded base, glabrous or sparsely pubescent on both sides. Peduncles terminal, $\frac{3}{4}$ – $3\frac{1}{2}$ in. long, naked or bibracteate. Thyrsus spike-like, moderately dense, 1– $2\frac{1}{2}$ in. long, 1– $1\frac{1}{2}$ in. diam. Bracts and bracteoles $2\frac{1}{2}$ – $3\frac{1}{2}$ lin. long, linear-lanceolate, very acute, glabrous or slightly pubescent and slightly ciliate. Sepals 5, free, 3– $3\frac{1}{2}$ lin. long, $\frac{2}{3}$ lin. broad, linear-lanceolate, very acuminate, pubescent, ciliate. Corolla two-lipped, pubescent in the upper part outside, apparently purple or red, spotted with darker; tube $\frac{1}{4}$ in. long; upper lip $4\frac{1}{2}$ in. long, oblong, with reflexed margins and minutely 2-lobed at the apex; lower lip $5\frac{1}{2}$ lin. long, 3-lobed, with an elevated plicate disk. Stamens 2, inserted at the middle of the tube; anthers 2-celled, with unequal cells, obtuse at each end, separated by a broad connective.

Kotinga Valley, *McConnell & Quelch*, 188; Kukenaam Valley, *im Thurn*, 81; Mount Roraima, *Appun*, 1387.

VERBENACEÆ.

LIPPIA SCHOMBURGKIANA, Schauer, in DC. Prod. xi. p. 577; Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 280.

Upper slopes of Mount Roraima, *McConnell & Quelch*, 2; Roraima Range, 3500 ft., *McConnell & Quelch*, 720; Ireng Valley, *McConnell & Quelch*, 235.

I doubt if this Guiana plant is really distinct from the Brazilian *L. glandulosa*, Schauer, described at the same place.

STACHYTARPHETA MUTABILIS, Vahl, Enum. i. p. 208.

Ireng Valley, *McConnell & Quelch*, 245.

LABIATÆ.

HYPTIS ARBOREA, Benth. in DC. Prod. xii. p. 132.

Kotinga Valley, *McConnell & Quelch*, 142, 184.

AMARANTACEÆ.

GOMPHRENA GLOBOSA, Linn. Sp. Pl. ed. 1, p. 224.

At Pelepowta Village in the Ireng Valley, *McConnell & Quelch*, 210.

PHYTOLACCACEÆ.

PHYTOLACCA THYRSIFLORA, Fenzl, ex J. A. Schmidt in Mart. Fl. Bras. ii. p. 343.

Kotinga Valley, *McConnell & Quelch*, 152.

PROTEACEÆ.

ROUPALA MONTANA, Aubl. Pl. Guian. i. p. 83.

Savannahs generally, *McConnell & Quelch*, 310.

LORANTHACEÆ.

PHORADENDRON RORAIMÆ, Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 281, and pp. 267, 269.

Summit of Mount Roraima, *McConnell & Quelch*, 82, 680.—Endemic.

BURMANNIACEÆ.

BURMANNIA BICOLOR, Mart. Nov. Gen. & Sp. i. p. 10, t. 5; R. Schomb. Reisen in Brit.-Guiana, iii. p. 1066.

Upper slopes of Mount Roraima, *McConnell & Quelch*, 14.—Widely dispersed in Tropical America from Cuba to Brazil, westward to Peru.

ORCHIDACEÆ. By R. A. ROLFE, A.L.S.

PLEUROTHALLIS RORAIMENSIS, Rolfe, sp. n. Lepanthiformis, caulibus gracilibus, vaginis apice ciliatis. Folia elliptica vel obovato-elliptica, minute apiculata, parva. Scapi subelongati, gracillimi, pauciflori. Bracteæ spathaceæ. Sepalum posticum basi ovatum, apice caudato-acuminatum; sepala lateralibus subsimilia, basi angustiora. Petala oblonga, obtusa. Labellum trilobum, lobis lateralibus rotundatis, intermedio oblongo obtuso.

Stems slender, $\frac{1}{2}$ in. long, clothed with three or four angular or striate sheaths, broader and ciliate at the apex. Leaves elliptical or obovate-elliptical, minutely apiculate, $2\frac{1}{2}$ –4 lin. long, $1\frac{1}{2}$ –2 lin. broad. Scapes very slender, $\frac{3}{4}$ –1 in. long, about 2- or

3-flowered. Bracts spathaceous, minute. Pedicels $1\frac{1}{2}$ lin. long. Dorsal sepal ovate at the base, caudate-acuminate above, 2 lin. long; lateral similar, but narrower at the base. Petals oblong, obtuse, 1 lin. long, thinly membranaceous. Lip trilobed, 1 lin. long; side lobes spreading, broadly rounded; front lobe oblong, obtuse; disc bearing a pair of thickened nerves. Column clavate, $\frac{3}{4}$ lin. long.

Roraima, summit, 8600 ft., *McConnell & Quelch*, 687.—Endemic.

Allied to the Colombian *Pleurothallis intricata*, Lindl., but smaller in all its parts, besides being different in structure. The species of the natural group *Lepanthiformes*, to which it belongs, were distributed by Lindley into three sections, owing to the exigencies of an admittedly artificial arrangement, which would place the present one in the section *Acuminatae*.

STELIS GUIANENSIS, Rolfe, sp. n. Cæspitosa. Caules brevissimi. Folia oblanceolato-oblonga, subobtusa, basi subattenuata. Scapi graciles, multiflori. Bracteæ triangulari-ovatae, acutæ. Pedicelli breves. Sepala orbiculari-ovata, obtusa, basi connata. Petala suborbicularia, concava. Labellum integrum, suborbiculare, petalis paullo angustius.

A dwarf, densely-tufted plant, with very short stems. Leaves oblanceolate-oblong, subobtuse, somewhat attenuate at the base, 4-9 lin. long, $1\frac{1}{2}$ -2 lin. broad. Scapes slender, $2\frac{1}{2}$ in. long, many-flowered. Bracts triangular-ovate, acute, $\frac{1}{2}$ lin. long. Pedicels $\frac{3}{4}$ lin. long. Sepals orbicular-ovate, obtuse, $\frac{3}{4}$ lin. long, slightly pubescent, the basal fourth connate. Petals broadly orbicular, somewhat concave, not half as long as the sepals. Lip entire, rather narrower than the petals, but otherwise very similar.

Roraima range, 3500 ft., *McConnell & Quelch*, 703.—Endemic.

A species belonging to Lindley's small group *Monastachyæ apodæ*, and allied to the Peruvian *S. pusilla*, H. B. & K. A specimen collected at the Kaieteur Savannah, Potaro River, British Guiana, by Jenman (1055), is either a form of the same or a closely allied species.

BRACHIONIDIUM BREVICAUDATUM, Rolfe, sp. n. Caulis primarius repens, brevis; secundarius brevissimus. Folia breviter petiolata, elliptico-lanceolata, tridenticulata. Pedunculi filiformes, uniflori. Bracteæ apice acuminatae. Sepalum posticum ovatum, apice abrupte acuminatum vel brevissime caudatum; sepala lateralina omnino connata et postico subsimilia. Petala ovata, acuminata vel brevissime caudata, margine plus minusve ciliata. Labellum sessile, trilobum, lobis lateralibus falcato-oblongis subobtusis, intermedio reflexo late ovato-rotundato. Columna lata.

Primary stem creeping, with short internodes, clothed with membranaceous sheaths; secondary very short, clothed with about two tubular membranaceous sheaths having a short acuminate apex. Leaves shortly petioled, elliptical-lanceolate, shortly tricuspidate, 8-10 lin. long, $2\frac{1}{2}$ - $3\frac{1}{2}$ lin. broad; petioles $1\frac{1}{2}$ -2 lin. long. Peduncles slender, $\frac{3}{4}$ - $1\frac{1}{4}$ in. long, bearing a short tubular acuminate sheath above the middle. Bract similar to the sheath but rather larger. Dorsal sepal ovate, abruptly acuminate or shortly caudate, 4 lin. long by nearly 3 lin. broad, trinerved; lateral pair united into a body closely resembling the dorsal sepal, but rather broader and 4-nerved. Petals ovate, shortly

acuminate, 3 lin. long, more or less ciliate, trinerved. Lip sessile, $1\frac{1}{2}$ lin. long, trilobed; side lobes falcate-oblong, subobtuse, suberect; front lobe reflexed, broadly rounded-oblong; disc with a slightly elevated crest between the side lobes. Column broad, $\frac{1}{2}$ lin. long.

Roraima range, 3500 ft., *McConnell & Quelch*, 705.—Endemic.

An interesting addition to this small genus, readily distinguished from its allies by the short secondary stem, and short tails to the sepals and petals.

MASDEVALLIA PICTURATA, Reichb. f. *Otia Bot. Hamb.* 16; *Ridl. in Trans. Linn. Soc. Ser. II. Bot. ii.* (1887), p. 281; *Woolw. Masd.* p. 87, t. 34.

Roraima, at 3500 ft., *McConnell & Quelch*, 704; Upper slope, *in Thurn*, 279.

In the mountains of Colombia, between 2500 and 6500 ft. elevation; extending along the Coast Andes of Venezuela to Caracas, at 6000 to 6500 ft.; also in Costa Rica. Roraima seems to be an outlying station for the species.

OCTOMERIA CONNELLII, Rolfe, sp. n. Caules secundarii elongati. Folia subsessilia, oblongo-lanceolata, subobtusa, crasso-coriacea. Flores subfasciculati, mediocres. Sepala et petala ovato-lanceolata, subobtusa. Labellum oblongum, subconcavum, obtusum vel obscure tridentatum, margine minutissime crenulatum.

Secondary stems stout, elongate, 4–12 in. long, clothed with four or five tubular sheaths $1-1\frac{1}{2}$ in. long. Leaf subsessile, thickly coriaceous or somewhat fleshy, oblong-lanceolate, subobtuse, 2–4 in. long, 4–10 lin. broad. Flowers few, or produced mostly in succession, medium-sized, shortly pedicelled. Sepals and petals ovate-lanceolate, subobtuse, 4–6 lin. long. Lip oblong, somewhat concave, obtuse or obscurely tridentate, margin minutely crenulate, 3–4 lin. long. Column clavate, $1\frac{1}{2}$ lin. long.

Roraima, summit at 8600 ft., *McConnell & Quelch*, 700.—Endemic.

A very distinct species, having stems much longer than the leaves, and flowers comparatively large for the genus. It is comparable with *O. grandiflora*, Lindl., but has longer stems, much shorter leaves, and the lip very different in structure.

OCTOMERIA PARVIFOLIA, Rolfe, sp. n. Cæspitosa. Caules secundarii graciles, vaginis tubulosis striatis obtecti. Folia lineari-oblonga, subacuta, parva. Flores fasciculati. Sepala ovato-oblonga, subobtusa. Petala lineari-oblonga, subobtusa. Labellum late deltoideo-trilobum, lobis lateralibus divergentibus anguste triangularibus, intermedio breviter et late triangulari obtuso.

Secondary stems densely tufted, slender, $2\frac{1}{2}$ –3 in. long, clothed with about five or six tubular striate sheaths. Leaves linear-oblong, subacute, $\frac{3}{4}$ – $1\frac{1}{4}$ in. long, $1-1\frac{1}{2}$ lin. broad, base somewhat attenuate. Flowers fasciculate. Pedicels very short. Sepals ovate-oblong, subobtuse, 1 lin. long. Petals linear-oblong, subobtuse, 1 lin. long. Lip broadly deltoid-triangular, over $\frac{1}{2}$ lin. broad; side lobes narrowly triangular, subobtuse; front lobe broadly triangular, obtuse; disc with a pair of thickened slightly arcuate keels in the centre, and a shorter one in front. Column short.

Roraima, summit, 8600 ft., *McConnell & Quelch*, 696.—Endemic.

In the relatively small proportion which the leaf bears to the rest of the plant, this

species approaches the Brazilian *Octomeria breviflora*, Cogn., but the structure of the lip is very different.

BULBOPHYLLUM RORAIMENSE, Rolfe, sp. n. Racemus multiflorus. Bracteæ oblongæ vel ovato-oblongæ, acutæ. Sepala ovato-lanceolata, acuminata, lateralalia carinata. Petala ovato-oblonga, subacuta. Labellum velutinum, pandurato-oblongum, obtusum, medio ad basin lateraliter compressum, basi utrinque auriculatum, disco elevato bicarinato. Columna brevis, utrinque bidentata.

Scape moderately slender; raceme 3 in. long, many-flowered. Bracts oblong or ovate-oblong, acute, $2\frac{1}{2}$ –3 lin. long. Pedicels 1 lin. long. Sepals ovate-lanceolate, acuminate, 4– $5\frac{1}{2}$ lin. long, lateral carinate. Petals ovate-oblong, subacute, $1\frac{1}{2}$ lin. long. Lip $2\frac{3}{4}$ lin. long, velvety, pandurate-oblong, obtuse; front lobe expanded into an ovate-oblong blade, middle and base laterally compressed, with a rounded erect basal auricle on either side; disc much elevated and channelled between the pair of obtuse keels. Column stout, 1 lin. long, with two pairs of teeth; upper pair subulate-filiform, incurved, with a broad base, as long as the column, lower pair subulate-oblong, a third as long as the upper.

Roraima, summit, 8600 ft., *McConnell & Quelch*, 103.—Endemic.

Allied to *B. geraense*, Reichb. f., but the scape and raceme more slender, the sepals proportionately broader, and the lip and teeth of the column different in detail. It is described from a single scape.

ELLEANTHUS LINIFOLIUS, Presl, Rel. Haenk. p. 97.

Ireng Valley, *McConnell & Quelch*, 301. Roraima, *Schomburgk*.—Also found in Cuba, Porto Rico, Nicaragua, Panama, Guiana, Brazil, and Bolivia.

DIACRIUM BICORNUTUM, Benth. in Journ. Linn. Soc., Bot. xviii. (1881), p. 312.

Kwaimatta Savannah, *McConnell & Quelch*, 280.—Also found in Guiana, Trinidad, and Tobago.

EPIDENDRUM GRANITICUM, Lindl. in Hook. Journ. Bot. iii. (1841), p. 83.

Kwaimatta Savannah, Rupununi River, *McConnell & Quelch*, 279. Ireng River, *im Thurn*, 13.—Found in British and Dutch Guiana, and Trinidad.

Nearly allied to *E. oncidoides*, Lindl., with which it is made synonymous by Lindley.

EPIDENDRUM SCHOMBURGKII, Lindl. Bot. Reg. (1838), Misc. p. 15; Ridl. in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 281.

Kotinga Valley, *McConnell & Quelch*, 135. “Flowers red, of all shades.”—Found in Trinidad, Guiana, Brazil, and Bolivia.

EPIDENDRUM FULGENS, Brongn. in Duperry, Voy. Coquille, Phan. p. 196, t. 43; Rolfe, in Orch. Rev. v. (1897), p. 264.

E. Schomburgkii, var. *confluens*, Lindl. Fol. Orch., Epidendr. p. 70.

E. Schomburgkii, Appun, Unter den Tropen, ii. p. 199 (non Lindl.).

Roraima, upper slopes, *McConnell & Quelch*, 1. North of Roraima, *Appun*, 1202. "Abundant."—Found in British Guiana, and the Brazilian provinces of Rio de Janeiro and Santa Catherina.

EPIDENDRUM ELONGATUM, Jacq. Coll. iii. p. 260 ; and 1c. Pl. Rar. iii. p. 17, t. 604 ; Ridl. in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 281.

Roraima, summit, at 8600 ft., *McConnell & Quelch*, 686. South side, at 5500 ft., *im Thurn*, 42.—Found in the Windward Islands, W. Indies, from Antigua to Trinidad, also in Guiana, Brazil, and Bolivia.

EPIDENDRUM ALSUM, Ridl. in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 281.

Roraima, summit, 8600 ft., *McConnell & Quelch*, 90, 699 ; *im Thurn*, 296.—Endemic.

EPIDENDRUM IMTHURNII, Ridl. in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 282, t. 46. fig. A.

Roraima, summit, 8600 ft., *McConnell & Quelch*, 697 ; *im Thurn*, 299.—Endemic.

EPIDENDRUM MONTIGENUM, Ridl. in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 282.

Roraima, upper slopes and summit, *McConnell & Quelch*, 37, 694 ; ledge and top, *im Thurn*, 322.—Endemic.

EPIDENDRUM VIOLASCENS, Ridl. in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 282, t. 46. fig. B.

Roraima, summit, 8600 ft., *McConnell & Quelch*, 81, 695 ; *im Thurn*, 300.—Endemic.

GALEANDRA JUNCEA, Lindl. Sert. Orch. sub t. 37.

Ireng Valley, *McConnell & Quelch*, 243.—Also found in Guiana and Brazil.

CYRTOPODIUM ANDERSONII, R. Br. in Ait. Hort. Kew. ed. 2, v. p. 216.

Kanuku Mts., *McConnell & Quelch*, 276.—Also found in Guiana, Brazil, and Paraguay.

CYRTOPODIUM CRISTATUM, Lindl. Bot. Reg. (1841), sub t. 8.

Kotinga Valley, amid stones on hill, *McConnell & Quelch*, 136. "Flowers yellow and brown."—Also found in Trinidad, Guiana, and Brazil.

ZYGOPETALUM BURKEI, Reichb. f. in Gard. Chron. xx. (1883), p. 684 ; Ridl. in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 282.

Z. Mackaii, R. Schomb. Reisen in Brit.-Guiana, ii. p. 266, iii. p. 1068 ; *Appun*, Unter den Tropen, ii. pp. 246, 292 (non Hook.).

Roraima, upper slopes, *McConnell & Quelch*, 26. "Flowers coffee-coloured." South side, at 5500 ft., *im Thurn*, 50. South and east sides, *Appun*, 1390.—Endemic.

ERIOPSIS SCHOMBURGKII, Reichb. f. in Bonplandia, iii. (1855), p. 67.

Pseudoeriosis Schomburgkii, Reichb. f. in Linnæa, xxii. (1849), p. 853; R. Schomb. Reisen in Brit.-Guiana, iii. p. 1123.

Roraima, upper slopes, *McConnell & Quelch*, 34. "Flowers brown." *Schomburgk*, 1679 a.—Also in the savannah of South Guiana.

HOULLETIA RORAIMENSIS, Rolfe, sp. n. Folia elliptico-lanceolata, magna. Scapus arcuatus, 7-9-florus. Bracteæ oblongæ, subobtusæ, concavæ. Sepala orbiculari-elliptica, concava. Petala orbiculari-elliptica, plana. Labellum profunde trilobum; lobi laterales late oblongi, truncati, oblique apiculatis; intermedius breviter unguiculatus, late trulliformis, apiculatus, callo transverso submembranaceo ad basin unguem sito. Columna clavata.

Leaf petiolate, elliptico-lanceolate, acute, $1\frac{1}{2}$ ft. long, $4\frac{1}{2}$ in. broad. Scape arching (base not seen); raceme 7-9-flowered. Bracts oblong, subobtuse, concave, 6-7 lin. long. Pedicels $1-1\frac{1}{4}$ in. long. Sepals orbicular-elliptical, concave, 9-11 lin. long, 7-9 lin. broad. Petals orbicular-elliptical, nearly flat, 9-10 lin. long, 7-8 lin. broad. Lip deeply 3-lobed; side lobes broadly oblong, nearly truncate and obliquely apiculate, 6 lin. long, $3\frac{1}{2}$ lin. broad, connected at the base by a suberect, transverse, stout, undulate membrane, and with a somewhat similar connecting callus at the junction of the front lobe, but more distinctly toothed; front lobe shortly and broadly unguiculate, broadly trulliform, abruptly apiculate, and with subacute lateral angles, 5 lin. long by nearly as broad. Column clavate, curved, 5 lin. long.

Roraima, upper slopes, *McConnell & Quelch*, 29.—Endemic.

A very distinct species, nearest *H. Lowiana*, Reichb. f., in structure, but very different in the broad nearly truncate side lobes of the lip, and more like *H. Brocklehurstiana*, Lindl., in habit.

CATASETUM DISCOLOR, Lindl. Bot. Reg. (1844), Misc. p. 34.

Roraima, upper slopes, *McConnell & Quelch*, 6.—Also found in Guiana, Brazil, and Paraguay.

MAXILLARIA CONNELLII, Rolfe, sp. n. Folia oblongo-lanceolata, acuta, basi in petiolum attenuata. Scapus foliis brevior, vaginis ovato-lanceolatis acutis subimbricatis obtectus. Bractea oblongo-lanceolata, acuta. Sepala subobtusa et minute apiculata; sepalum lateralia triangulari-oblonga; posticum oblongo-lanceolatum. Petala oblongo-lanceolata, subacuta. Labellum elliptico-oblongum, obtusum, subundulatum, apice recurvum, callo oblongo apice dilatato truncato. Columna brevis

Leaves oblong-lanceolate, acute, attenuate at the base, 6-8 in. long, 9-12 lin. broad. Scape $3\frac{1}{2}$ -4 in. long, covered with ovate-lanceolate acute sheaths, 6-9 lin. long and somewhat imbricate at the base. Bract oblong-lanceolate, acute, 9-12 lin. long. Sepals subobtuse and minutely apiculate, dorsal oblong-lanceolate, 7-8 lin. long; lateral triangular-oblong, 8-9 lin. long. Petals oblong-lanceolate, subacute, 7-8 lin. long. Lip elliptical-oblong, obtuse, somewhat undulate, recurved at the apex, rather fleshy,

5–6 lin. long; callus oblong, dilated and truncate at the apex. Column stout, free part scarcely 2 lin. long, with a much longer foot. Mentum rounded, obtuse, 3 lin. long.

Roraima: Kotinga Valley, *McConnell & Quelch*, 137.—Endemic.

Allied to the Colombian *Maxillaria melina*, Lindl., but the leaves smaller, sheaths of the scape shorter, and with various differences in the structure of the flower.

MAXILLARIA QUELCHII, Rolfe, sp. n. Subcaulescens, vaginis imbricatis multistriatis.

Pseudobulbi compressi, apice monophylli. Folia oblonga, subobtusata, conduplicata. Scapi foliis subæquales, vaginis lanceolatis subimbricatis obtecti. Bractea lanceolata, acuminata. Sepala oblongo-lanceolata, acuta. Petala lineari-lanceolata, acuta, sepalis breviora. Labellum trilobum, lobis lateralibus erectis rotundato-oblongis membranaceis, intermedio recurvo obovato-orbiculari obscure tricuspidato medio carnosio lateribus incurvo, callo oblongo obtuso concavo.

Subcaulescent, with the stems clothed with conduplicate, triangular-lanceolate, acute, closely striate, imbricating sheaths, 1–2 in. long, and partly enveloping the pseudobulbs. Pseudobulbs compressed, broadly oblong, shining and punctulate, wrinkled when old, 1 in. long, $\frac{3}{4}$ in. broad, monophyllous at the apex. Leaves conduplicate, lanceolate-oblong, subobtusate, 3–4 in. long, $\frac{3}{4}$ –1 in. broad. Scares 3–4 in. long, bearing 5–7 lanceolate, acute, striate, somewhat imbricating sheaths, 6–12 lin. long. Bract resembling the sheaths, 10–12 lin. long. Sepals oblong-lanceolate, acuminate, $1\frac{1}{2}$ in. long. Petals linear-lanceolate, acuminate, $1\frac{1}{4}$ in. long. Lip $\frac{1}{2}$ in. long, 3-lobed; side lobes erect, rounded-oblong, membranaceous, 1 lin. long, $1\frac{1}{4}$ lin. broad; middle lobe recurved, obovate-orbicular, obscurely tricuspidate at the apex, 2 lin. broad by nearly as long, thickened along the centre and incurved at the sides; crest oblong, obtuse, concave. Column clavate, incurved, 3 lin. long.

Roraima, summit, at 8600 ft. alt., *McConnell & Quelch*, 690.—Endemic.

Allied to the Peruvian *M. floribunda*, Lindl., but much less scandent, and smaller in all its parts.

ONCIDIUM ORTHOSTATES, Ridl. in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 283.

Kwaimatta Savannah, *McConnell & Quelch*, 281. Ireng River, *im Thurn*, 12.—Also found at Mimatta, on the Rupununi River.

SOBRALIA STENOPHYLLA, Lindl. Fol. Orch., Sobral. p. 2; Ridl. in Trans. Linn. Soc. Ser. II. ii. (1887), p. 283.

Kotinga River, *McConnell & Quelch*, 147. Ireng Valley, *McConnell & Quelch*, 329. Ipelemouta, Arapu River, *im Thurn*, 19. Roraima, *Appun*, 1076.—Endemic.

EPISTEPHIUM LUCIDUM, Cogn. in Mart. Fl. Bras. iii. pt. iv. p. 141, t. 30.

Roraima, upper slopes, *McConnell & Quelch*, 32.—Also found in Brazil.

EPISTEPHIUM PARVIFLORUM, Lindl. Gen. & Sp. Orch. p. 433.

Kotinga Valley, *McConnell & Quelch*, 160.—Also found in Trinidad and elsewhere in British Guiana.

SPIRANTHES BIFIDA, Ridl. in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 283.

Kotinga Valley, *McConnell & Quelch*, 200. Roraima, summit, 8600 ft. alt., *McConnell & Quelch*, 692. South side at 5500 ft., *im Thurn*, 342.—Endemic.

POGONIA PARVIFLORA, Reichb. f. Xen. Orch. ii. p. 90; Ridl. in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 283.

Roraima, upper slopes, *McConnell & Quelch*, 30. "Flowers pink and waxy." South side at 5500 ft., *im Thurn*, 115.—Endemic.

POGONIA TENUIS, Reichb. f. in Griseb. Fl. Brit. W. Ind. p. 637.

Roraima, summit, 8600 ft., *McConnell & Quelch*, 691.—Also found in Trinidad, Venezuela, and on the Upper Amazon.

HABENARIA MORITZII, Ridl. in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 284.

Roraima, upper slopes, *McConnell & Quelch*, 4; at 4000 ft., *im Thurn*, 367.—Also found in Venezuela.

HABENARIA RORAIMENSIS, Rolfe, sp. n. Planta erecta, foliosa. Folia caulina, basi lanceolata, acuta, superiora gradatim in vaginas bracteiformes decrescentia. Bracteae lanceolatae, acuminatae. Sepalum posticum erectum, ovatum, apiculatum; sepala lateralibus patentia, oblique ovato-oblonga, apiculata. Petala bipartita, lobo postico falcato-oblongo apiculato, antico filiformi-lineari arcuato. Labellum tripartitum; lobi laterales filiformi-lineares divergentes; lobus intermedius linearis obtusus; calcar clavatum.

Plant $\frac{1}{2}$ – $1\frac{1}{4}$ ft. high, somewhat leafy. Leaves cauline, the lower lanceolate, acute, 2– $2\frac{1}{2}$ lin. long, 4–5 lin. broad; upper smaller, decreasing upwards into bract-like sheaths. Bracts lanceolate, acuminate, 6–9 lin. long. Dorsal sepal erect, ovate, apiculate, 2– $2\frac{1}{2}$ lin. long, lateral spreading obliquely, ovate-oblong, apiculate, 2– $2\frac{1}{2}$ lin. long. Petals bipartite; upper lobe falcate-oblong, apiculate, $1\frac{1}{2}$ –2 lin. long; lower lobe filiform-linear, curved, shorter than the upper. Lip tripartite; side-lobes filiform-linear, diverging, $1\frac{1}{2}$ –2 lin. long; front lobe linear, obtuse, 2– $2\frac{1}{2}$ lin. long; spur linear-clavate, 4–5 lin. long. Anther-channels and stigmatic processes short.

Roraima, summit, 8600 ft., *McConnell & Quelch*, 698.—Endemic.

Allied to *H. pratensis*, Reichb. f., but readily distinguished by its much smaller flowers.

PHRAGMIPEDIUM KLOTZSCHIANUM, Rolfe, in Orch. Rev. iv. (1896), p. 332.

Cypripedium Klotzschianum, Reichb. f. in Linnæa, xxii. (1849), p. 811; Appun, Unter den Tropen, ii. p. 196.

Selenipedium Klotzschianum, Reichb. f. in Bonplandia, ii. (1854), p. 116; Ridl. in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 284.

Cypripedium Schomburgkianum, Klotzsch, in R. Schomb. Reisen in Brit.-Guiana, ii. p. 229; R. Schomb. Bot. Rem. Brit. Guiana, p. 59.

Arabapu River, *McConnell & Quelch*, 144. Aroie Creek, Kotinga River, *im Thurn*, 5. Ipelemouta, Arapu River, *im Thurn*, 31. Kako River, *Appun*, 1212.—Endemic.

BROMELIACEÆ.

BROCCHINIA REDUCTA, Baker, in Journ. of Bot. xx. (1882), p. 331; Baker, Handb. Brom. p. 88; Mez, in DC. Monogr. Phan. ix. p. 341. (Plate 12.)

Roraima Range, 3500 ft., *McConnell & Quelch*, 702.—Also found on the Kaieteur Savannah, *Jenman*, 873.

PUYA FLOCCOSA, E. Morren, in Belg. Hort. xxxv. (1878), p. 81; Mez, in DC. Monogr. Phan. ix. p. 478.

P. guianensis, Klotzsch, in R. Schomb. Reisen in Brit.-Guiana, iii. p. 1067.

P. sp., Baker, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 285.

Roraima Range, 3500 ft., *McConnell & Quelch*, 701.—Also in Venezuela, Colombia, and Bolivia.

CONNELLIA, N. E. Brown, gen. nov.

Flores hermaphroditi, regulares. Sepala libera, oblonga. Petala libera, lata, basi angustata, nuda, sepalis multo longiora. Stamina petalis breviora et iis basi breviter adnata, filamentis filiformibus, antheris oblongis basifixis. Ovarium superum, trigonum, loculis multiovulatis; stylus elongatus; stigmata linearia. Capsula in carpellis 3 septicide separata, carpellis introrsum dehiscentibus. Semina numerosa, parva, linearia, utrinque appendiculata.—Herbæ habitu *Tillandsiæ*. Folia rosulata, integra vel basi denticulata. Inflorescentia terminalis, spiciformis, simplex vel composita. Flores sub quaque bractea solitarii vel plures, pedicellati.

CONNELLIA AUGUSTÆ, N. E. Brown. (Plate 13.) Capsula 6-7 lin. longa, 3 lin. diam., trigona, oblonga, coriacea, glabra, in carpellis 3 septicide soluta, carpellis longe rostratis introrsum dehiscentibus. Semina $1\frac{3}{4}$ -2 lin. longa, lineari-teretia, curvata, utrinque membranaceo-appendiculata.—*Encholirium Augustæ*, R. H. Schomb. in Verh. des Ver. Beförd. Gartenb. in Preuss. (1846), p. 18, t. 2; in Bot. Zeitung (1846), p. 454; R. Schomb. Reisen in Brit.-Guiana, ii. p. 271, and iii. p. 1067; & in Bot. Rem. Brit. Guiana, p. 77; De Beer, Brom. p. 27. *Caraguata Augustæ*, Benth. & Hook. f. Gen. Pl. iii. p. 668. *Dyckia Augustæ*, Baker, Handb. Brom. p. 135. *Puya Augustæ*, Mez, in DC. Monogr. Phan. ix. p. 487, partim.

Summit of Mount Roraima, 8600 ft., *McConnell & Quelch*, 670. South slopes of Roraima at 6500 ft., *Schomburgk*, 687, 1021.

The discovery of fruiting specimens of this plant by Messrs. McConnell & Quelch demonstrates that it cannot properly be referred to either of the genera under which it has previously been placed. From *Encholirion* it differs totally in habit, large bracts, broad petals, and seeds. From *Caraguata* it differs in its free petals and in its seeds. From *Dyckia* it differs in habit, in the long-beaked entire (not bifid) carpels into which the ripe fruit separates, and in its seeds. And from *Puya* it differs in habit, in the petals not twisting when the flower fades, in the capsule not dehiscing loculicidally, and in the seeds. The only genus which approaches it in structure is *Lindmania*, Mez, but the habit, branching panicle, small bracts and very small flowers of the species of *Lindmania*

are so entirely different from those of *Connellia*, that they cannot naturally be placed in the same genus. Possibly the fruit of *Lindmania*, which I have not seen, and the dorsifixed anthers may afford technical characters to distinguish the two genera, especially if taken in conjunction with the difference in habit.

CONNELLIA QUELCHII, N. E. Brown. (Plate 14.) Folia rosulata $1\frac{1}{2}$ – $5\frac{3}{4}$ poll. longa, angusta, convoluta, integra, basi tantum minute denticulata, supra tomentosa, subtus glabra, vel rare utrinque tomentosa, marginibus albo-tomentosis. Scapus simplex, multibracteatus. Bracteæ imbricatæ, glabræ, nitidæ; inferiores steriles amplexicaules, abrupte subulato-cuspidatæ; florigeræ elliptico-ovatæ, apiculatæ, valde concavæ. Flores sub quaque bractea solitarii, pedicellati, speciosi, rosei.—*Tillandsia stricta*, var. ?, Baker, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 285. *Puya Augustæ*, Mez, in DC. Monog. Phan. ix. p. 487, partim.

Leaves rosulate, $1\frac{1}{2}$ – $5\frac{3}{4}$ in. long, 2– $4\frac{1}{2}$ lin. broad, linear, convolute-subulate, rather blunt, rigidly coriaceous, minutely denticulate at the base only, densely white-tomentose on the inner face and along the margins, usually glabrous on the convex back, rarely tomentose on both sides. Scape arising from the centre of the rosette, 2–5 in. long to the lowest flower of the 2– $3\frac{1}{2}$ in. long raceme, 1– $1\frac{1}{2}$ lin. thick, glabrous, clothed with imbricating bracts, which are $\frac{2}{3}$ – $1\frac{1}{4}$ in. long, 6–8 lin. broad, stem-clasping, broadly elliptic-oblong and abruptly contracted into a subulate leafy point at the apex, or broadly ovate and simply acute or acuminate, glabrous, with tomentose margins to the leafy point, brown in the dried state, shining. Raceme moderately dense, $1\frac{1}{4}$ – $1\frac{3}{4}$ in. diam., 7–13-flowered, its bracts ascending or somewhat spreading, 7–10 lin. long, 5–7 lin. broad, elliptic-ovate, acute or obtuse and apiculate, deeply concave, glabrous, brown, shining. Flowers solitary under each bract, pedicellate. Pedicels 2–8 lin. long, glabrous. Sepals free, 5– $5\frac{1}{2}$ lin. long, $2\frac{1}{2}$ –3 lin. broad, ovate-oblong, acute, glabrous, light brown, with thin rose-pink margins. Petals free, 9 lin. long, $5\frac{1}{2}$ lin. broad, orbicular, obtuse, narrowed into a short broad claw at the base, glabrous, entire, without a scale at their base, bright rose-pink. Stamens about 5 lin. long, glabrous; filaments very shortly adnate to the petals at their base, filiform; anthers $\frac{3}{4}$ lin. long, oblong, obtuse. Ovary superior, trigonous-ovoid, glabrous, narrowed into a glabrous style $3\frac{1}{2}$ lin. long, with 3 recurving compressed linear stigmas about 1 lin. long, slightly undulated along the stigmatose surface.

Summit of Mount Roraima, 8600 ft., *im Thurn*, 315; *McConnell & Quelch*, 107, 672.

In general appearance this plant bears a slight resemblance to *Tillandsia stricta*, Soland., but in all details is very different. It is considered by Mez to be identical with *C. Augustæ*; but although an undoubted congener of that plant, is most certainly very distinct from it specifically, being very much smaller in size, with the upper surface and margins of the leaves tomentose, solitary flowers under each bract, and a glabrous calyx; whilst *C. Augustæ* has leaves 8–14 in. long, quite glabrous on both sides, the flowers are in clusters of 3–7 under each bract, and the calyx is thinly covered with short hairs. I have not seen ripe seeds of *C. Quelchii*, but the ovules show that they will be appendaged at each end. It is a very pretty species and well worth cultivating.

TILLANDSIA RHODOCINCTA, Baker, in Journ. Bot. xxvi. (1888), p. 143, & Handb. Brom. p. 178; Mez, in DC. Monog. Phan. ix. p. 791. *Tillandsia sp.*, Baker, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 285.

Summit of Mount Roraima, *McConnell & Quelch*, 671.—Endemic.

IRIDACEÆ.

SISYRHINCHIUM ALATUM, Hooker, Icones Plant. t. 219, var.

Kotinga Valley, *McConnell & Quelch*, 139.

LILIACEÆ.

NIETNERIA CORYMBOSA, Klotzsch, in R. Schomb. Reisen in Brit.-Guiana, iii. p. 1066; in Thurn & Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), pp. 269, 285.

Upper slopes of Mount Roraima, *McConnell & Quelch*, 12, 36, 328. Summit of Mount Roraima, *McConnell & Quelch*, 83, 656.—Apparently endemic to the region.

TOFIELDIA SCHOMBURGKIANA, Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 285, t. 49. f. A.

Isidrogalvis guianensis, Klotzsch, in R. Schomb. Reisen in Brit.-Guiana, iii. p. 1065, name only.

Upper slopes and summit of Mount Roraima, *McConnell & Quelch*, 93, 326, 657; *Schomburgk*.—Endemic.

XYRIDACEÆ.

XYRIS SEUBERTII, Nilss. in K. Svensk Akad. Handl. xxiv. (1892), No. xiv. p. 51, t. 4. f. 1.

Upper slopes of Mount Roraima, *McConnell & Quelch*, 41.—The locality in which Schomburgk collected this plant is unknown to me, and the leaves of his specimens are much smaller than those of the specimens collected by McConnell & Quelch.

XYRIS CONCINNA, N. E. Brown, sp. n. Planta 3–6 poll. alta, perennis, glabra. Caudex $\frac{1}{4}$ – $\frac{1}{2}$ poll. longus. Folia disticha, 1–2 $\frac{3}{4}$ poll. longa, anguste linearia, acuta. Pedunculus subteres, uno margine prominente, basi vagina apice breviter foliata instructus. Spica ovoidea. Bracteæ oblongæ vel ellipticæ, obtusæ, castaneæ, late albo-marginatæ, laceratæ, glabræ. Sepala lateralibus lineari-lanceolata, acuta, carinata, glabra, carina integra vel ad medium minutissime scabrida. Petala elliptica, obtusa, integra, lutea. Staminodia bibrachiata, brachiis penicillatis luteis.

Perennial. Caudex $\frac{1}{4}$ – $\frac{1}{2}$ in. (or more?) long, $\frac{1}{2}$ lin. thick, naked below the leaves, emitting fibrous roots. Leaves distichous, 1–2 $\frac{3}{4}$ in. long, $\frac{1}{3}$ – $\frac{2}{3}$ lin. broad, linear, acute, abruptly dilated into a short, broad, bright brown, clasping base, not twisted, glabrous. Peduncles 1–6 to a stem, 2 $\frac{1}{2}$ –5 $\frac{1}{2}$ in. long, $\frac{1}{4}$ – $\frac{1}{3}$ lin. thick, subterete, with a raised line along one side, glabrous; their basal sheaths $\frac{3}{4}$ –1 $\frac{3}{4}$ in. long, produced at the apex into a short compressed or leafy point, keeled, glabrous, often minutely ciliate along the keel or at its apex. Spikes 2 $\frac{1}{2}$ –3 $\frac{1}{2}$ lin. long, ovoid, obtuse. Bracts 1 $\frac{1}{2}$ –2 $\frac{1}{2}$ lin. long, 1–1 $\frac{1}{2}$ lin.

broad, oblong or elliptic, obtuse, chestnut-brown, with a broad white hyaline border, which soon becomes lacerated, glabrous. Lateral sepals $2\frac{3}{4}$ lin. long, $\frac{1}{2}$ lin. broad, complicate, linear-lanceolate, acute, keeled, yellowish-brown, glabrous, the keel quite entire or very minutely scabrid along the middle. Petals 2 lin. long, 1 lin. broad, elliptic, obtuse, entire, yellow. Staminodes 2-armed, about $\frac{1}{2}$ lin. long, divided into tufts of yellow hairs.

Summit of Mount Roraima, 8600 ft., *McConnell & Quelch*, 496.

Allied to *Xyris Seubertii*, Nilss., but distinguished by its smaller size, lacerated bracts, and the glabrous or very minutely scabrid keel of the lateral sepals. The broad bright brown bases of the leaves give the base of the plant a somewhat bulbous appearance.

XYRIS WITSENIODES, Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 285, t. 50. f. B.

X. caulescens, Klotzsch, in R. Schomb. Reisen in Brit.-Guiana, iii. p. 1064, name only.

Summit of Mount Roraima, *McConnell & Quelch*, 95, 658.—Endemic.

ABOLBODA SCEPTRUM, Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 286, & pp. 258, 259, 262, 267, & 268.

Summit of Mount Roraima, *McConnell & Quelch*, 668.—Endemic on Roraima and the surrounding region.

The specimen collected by McConnell & Quelch consists of adult leaves and young plants, and that collected by im Thurn of the upper part of a flowering stem without leaves, so that this interesting plant is as yet very imperfectly represented in European Herbaria.

RAPATEACEÆ.

STEGOLEPIS GUIANENSIS, Klotzsch, *ex* Koern. in Linnæa, xxxvii. (1871-73), p. 481; im

Thurn & Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), pp. 265, 267 & 286.

Summit of Mount Roraima, *McConnell & Quelch*, 669.—Endemic.

ERIOCAULACEÆ.

ERIOCAULON TENUIFOLIUM, Klotzsch, in R. Schomb. Reisen in Brit.-Guiana, iii. p. 1116.

Savannahs generally, *McConnell & Quelch*, 307.

PÆPALANTHUS RORAIMÆ, Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 286.

Summit of Mount Roraima, *McConnell & Quelch*, 102, 315, 660.—Endemic.

PÆPALANTHUS FLAVESCENS, Koern. in. Mart. Fl. Bras. iii. pt. I. p. 423; im Thurn & Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), pp. 263, 286.

P. eriocephalus, Klotzsch in R. Schomb. Reisen in Brit.-Guiana, iii. p. 1064, name only.

Upper slopes of Mount Roraima, *McConnell & Quelch*, 9, 10, 327.—Also in Venezuela and Brazil.

PÆPALANTHUS FRATERNUS, N. E. Brown, sp. n. Acaulis. Folia dense rosulata, linearia, subacuta, basi late dilatata, supra canaliculata, rigida, supra pilis articulatis appressis

tenuiter oblecta et pilis longis paucis laxè ciliata, subtus vel rare utrinque glabra; in axillis dense lanato-villosa. Pedunculi 8-12 poll. longi, laxè villosi, rare glabri. Capitula $4\frac{1}{2}$ -5 lin. diam., albo-villosa. Bracteæ involucrantès 3-4-seriatæ, late ovatæ, acutæ, dorso appresse pubescentes, fuscæ. Bracteæ inter flores oblongæ, subacutæ, apice dense albo-barbatæ. Receptaculum pilosum. Sepala elliptica, obtusa, hyalino-olivacea, ciliata, apice dense albo-barbata. Petala floris masculini in tubum infundibuliformem stipitatum connata; floris fœminei libera, sessilia, angustè linearia, dense ciliato-barbata.

A stemless perennial. Leaves densely rosulate, $1-1\frac{3}{4}$ in. long, $1-1\frac{1}{2}$ lin. broad, linear, tapering to a blunt point, rather abruptly dilated into a broad open sheath at the base, rigid, concave and thinly covered with adpressed white jointed hairs on the face, convex and glabrous on the back, rarely glabrous on both sides, veins not prominent, scarcely striate, often, but not always, thinly ciliate with a few long hairs, densely villose-woolly with long fine hairs in the axils. Peduncles 1-3 to a plant, 8-12 in. long, slender, obscurely 6-7-angular, thinly covered with short spreading hairs or rarely glabrous; their basal sheaths $1-1\frac{3}{4}$ in. long, with a rigid oblique subacute ciliate mouth, otherwise glabrous. Heads $4\frac{1}{2}$ -5 lin. diam., at first hemispherical, afterwards globose, monœcious or rarely unisexual. Involucral bracts in 3-4 series, the innermost $1\frac{3}{4}$ -2 lin. long, $1\frac{1}{2}$ lin. broad, broadly ovate, acute, fuscous, adpressed-pubescent on the back, ciliate, the outer gradually smaller. Flowering-bracts $1\frac{1}{2}$ lin. long, $\frac{1}{2}$ lin. broad, oblong, obtuse or subacute, pale fuscous, densely bearded with white hairs on the apical part. Receptacle pilose with long white hairs, as are also the pedicels of the flowers. Male flowers pedicellate; sepals 3, $1\frac{1}{4}$ lin. long, $\frac{1}{2}$ lin. broad, elliptic, obtuse or subacute, fuscous, densely bearded with white hairs on the apical part; petals arising $\frac{1}{2}$ lin. above the sepals, all united into a hyaline funnel-shaped tube $\frac{1}{2}$ lin. long, truncate, with 3 very minute teeth at the apex, glabrous; stamens shortly exserted, with white anthers; rudimentary pistil represented by 3 clavate bodies, minutely papillate at the apex. Female flowers shortly pedicellate; sepals $1\frac{1}{2}$ lin. long, $\frac{2}{3}-\frac{3}{4}$ lin. broad, elliptic, subacute, beautifully reticulated with fuscous, ciliate from the base and densely bearded on the upper half with long white hairs; petals 3, free, arising close to the sepals, 1 lin. long, about $\frac{1}{6}$ lin. broad, ciliate and clothed on both sides with long white hairs; ovary ovoid, trigonous, glabrous, narrowed into a style about $\frac{1}{3}$ lin. long, with 3 linear bifid stigmas, and 3 stout appendages, papillate at the apex only.

Summit of Mount Roraima, 8600 ft., *McConnell & Quelch*, 96, 659.

Allied to *P. falcatus*, Koern., and *P. flavescens*, Koern. From the former it differs by its more rigid, channelled leaves, less woolly heads, and more acute involucral bracts, and from *P. flavescens*, Koern., by the narrower and more acute leaves, which have a different pubescence, and the very different flower-heads. It also appears to be near *P. Schomburgkii*, Klotzsch (which I have not seen), but that plant is described as having longer and broader leaves, striate, with prominent veins on the upper side, a subulate point to the peduncular sheath, much shorter peduncles, and glabrous involucral bracts.

PÆPALANTHUS SUBCAULESCENS, N. E. Brown, sp. n. Planta 4–8 poll. alta. Caulis simplex, dense foliosus. Folia lineari-lanceolata, acuta, glabra, juniora pilis longis laxè ciliata. Pedunculi 2–7-aggregati, obtuse 4-angulati, glabri, vaginis oblique fissis apice subulato-apiculatis glabris ore ciliatis. Capitula hemisphærica, albo-villosa. Bracteæ involucrantès suborbiculares vel late obovatæ, obtusissimæ, minutæ ciliatæ, fuscæ. Bracteæ inter flores oblongæ, obtusæ, apiculatæ, apice albo-barbatæ. Receptaculum glabrum, sed pedicellis laxè pilosis. Floris masculi sepala obovata, obtusa, apice albo-barbata; petala in tubum tridentatum longe stipitatum connata. Floris fœminei sepala elliptica vel obovata, obtusa, apice albo-barbata; petala libera, oblonga, obtusa, pilis longis dense ciliata, dorso glabra.

A perennial, with a simple densely leafy stem $\frac{1}{2}$ – $2\frac{1}{2}$ in. long. Leaves $\frac{3}{4}$ – $1\frac{1}{2}$ in. long, 1 – $2\frac{1}{2}$ lin. broad, linear-lanceolate, acute, glabrous, striate, green, the younger laxly ciliate with long white hairs, ascending, becoming deflexed with age. Peduncles 2–7 to a plant, $2\frac{1}{2}$ – $5\frac{1}{2}$ in. long, slender, obtusely 4-angled, glabrous; their basal sheaths 1 – $1\frac{1}{4}$ in. long, glabrous, with an oblique ciliate mouth, acute or subulate-pointed. Heads 2–3 lin. diam., hemispherical, monœcious, white-villous. Involucral bracts in 3–4 series; the innermost $\frac{3}{4}$ lin. long, $\frac{3}{4}$ – 1 lin. broad, suborbicular or broadly obovate, very obtuse, glabrous, minutely ciliate, fuscous, the outer gradually smaller. Flowering bracts $\frac{3}{4}$ – 1 lin. long, $\frac{1}{2}$ lin. broad, oblong, obtuse, apiculate, fuscous, densely bearded with short white hairs on the apical part. Receptacle apparently glabrous, but with the base of the pedicels pilose with long hairs. Male flowers: sepals $\frac{3}{4}$ lin. long, $\frac{1}{3}$ lin. broad, obovate, obtuse, brown, bearded with white hairs at the apex; petals connate into a hyaline, subtruncate, minutely 3-toothed tube, separated from the sepals by a stipes nearly $\frac{1}{2}$ lin. long, glabrous; anthers white. Female flowers: sepals $\frac{5}{8}$ – 1 lin. long, nearly $\frac{1}{2}$ lin. broad, elliptic or obovate, obtuse, fuscous or brownish, densely bearded with short white hairs on the apical part; petals 3, free, arising close to the sepals, $\frac{3}{4}$ lin. long, $\frac{1}{3}$ lin. broad, oblong, obtuse, light fuscous, densely fringed with very long hairs on the margins and borders of the inner face, glabrous on the back; ovary glabrous, trigonous; style elongated; stigmas linear, bifid to halfway down; appendages rather thick, papillate at the apex.

Kotinga Valley, *McConnell & Quelch*, 153. Savannahs generally, *McConnell & Quelch*, 300.

A very distinct species, bearing a resemblance to *P. uncinatus*, Gardn., in stem and foliage, but with very different flower-heads. It might be placed near *P. plantagineus*, Koern.

PÆPALANTHUS BIFORMIS, N. E. Brown, sp. n. Planta pusilla, acaulis, ut videtur annua. Folia rosulata, 3–4 lin. longa, linearia, acuta, supra arachnoideo-tomentosa, demum glabra. Pedunculi 3–4, filiformes, $1\frac{3}{4}$ – $2\frac{1}{4}$ poll. longi, laxè villosi vel subglabri, vaginis glabris ore obliquis acutis. Capitula hemisphærica, 2– $2\frac{1}{2}$ lin. diam., pallide luteo-alba. Bracteæ involucrantès oblongæ vel ovato-oblongæ, acutæ, glabræ. Bracteæ inter flores nullæ. Receptaculum et pedicelli lanati. Flores masculi quam fœminei multo minores, oblique deltoidei, acuti, sepalis inæqualibus ad medium barbatis, petalis in tubum minutum stipitatum connatis. Flores fœminei recti; sepala

eorum 3, libera, anguste lanceolata, acuminata, carinata, glabra, ad medium ciliata; petala 3, cuneato-obovata, prope apicem obtusum incurvum leviter connata, dorso pilis longis dense vestita.

A small stemless annual? Leaves rosulate, very spreading, 3-4 lin. long, $\frac{1}{4}$ - $\frac{1}{2}$ lin. broad, linear, acute, thinly cobwebby-tomentose above, becoming glabrous, glabrous beneath, woolly in the axils. Peduncles 3-4 to a plant, $1\frac{3}{4}$ - $2\frac{1}{4}$ in. long, filiform, thinly villose or nearly glabrous. Heads simple, monœcious, 2- $2\frac{1}{2}$ lin. diam., very pale straw-coloured in all parts. Involucral-bracts in 3-4 series, much shorter than the flowers, the innermost $\frac{3}{4}$ lin. long, $\frac{1}{4}$ - $\frac{1}{3}$ lin. broad, oblong or ovate-oblong, acute, quite glabrous, thin, semi-transparent. Flowering-bracts none. Receptacle and pedicels very woolly. Male flowers much smaller than the female flowers, with rather longer pedicels, very oblique; sepals 3, two of them $\frac{1}{2}$ lin. long, $\frac{1}{4}$ lin. broad, obliquely semiovate, straight along one margin, very much curved along the other, acute, the third smaller, rhomboid-lanceolate, acute, all bearded with short hairs at about the middle, transparent; petals connate into a very minute, stipitate, funnel-shaped cup, glabrous; anthers white. Female flowers straight, projecting much beyond the males; sepals 3, free, 1 lin. long, $\frac{1}{4}$ lin. broad, narrowly lanceolate, acuminate, boat-shaped, keeled, glabrous, ciliate at the middle only, transparent; petals arising a little above the sepals, scarcely $\frac{1}{2}$ lin. long, about $\frac{1}{8}$ lin. broad, cuneate-obovate, obtuse and inflexed at the apex, and apparently slightly connate there, densely clothed with long hairs on the back; ovary trigonous, glabrous, with a glabrous style, divided at the apex into 3 simple stigmas and 3 slender clavate appendages. Capsule scarcely $\frac{1}{4}$ lin. diam. Seeds ellipsoidal, faintly ribbed, glabrous, brown.

Kotinga Valley, near Mount Roraima, *McConnell & Quelch*, 126.

This species much resembles *Pæpalanthus simplex*, Miq., but is readily distinguished by the very unequal size and dissimilar form of the male and female flowers, the females conspicuously projecting much beyond the males, so that the heads have a somewhat echinate appearance.

PÆPALANTHUS UMBELLATUS, Kunth, Enum. Pl. iii. p. 537.

Kotinga Valley, *McConnell & Quelch*, 129.

PÆPALANTHUS SUBTILIS, Miq. Stirp Surin. Sel. p. 221.

Savannahs generally, *McConnell & Quelch*, 312.

PÆPALANTHUS CAPILLACEUS, Klotzsch, in R. Schomb. Reisen in Brit.-Guiana, i. p. 377, ii. p. 5, & iii. p. 1063; Koern. in Mart. Fl. Bras. iii. pt. i. p. 415, t. 53. f. 2.

Kukenaam River, *McConnell & Quelch*, 314. Roraima, *Appun*, 1217.—Endemic to the region.

CYPERACEÆ.

CYPERUS UNCINATUS, Poir. Encycl. vii. p. 247.

Kotinga Valley near Roraima, *McConnell & Quelch*, 128. Savannahs generally, *McConnell & Quelch*, 321.

FUIRENA UMBELLATA, Rottb. Deser. et Ic. Pl. p. 70, t. 19. f. 3.

Ireng Valley, *McConnell & Quelch*, 254.

HYPOLYTRUM PUNGENS, Vahl, Enum. ii. p. 283.

Kotinga Valley, *McConnell & Quelch*, 179.

EVERARDIA ANGUSTA, N. E. Brown, sp. n. Folia linearia, acuta, laxe pilosa. Culmi basi vaginati, aphylli, valde compressi, leviter concavo-convexi, parce pilosi vel fere glabri. Panicula elongata, linearia, stricta, ramis gracilibus erectis 1- (rare 2-)spiculatis in axillis vaginorum aggregatis. Spiculæ masculæ lanceolatae, glumis 5-7 coraceis oblongis acutis mucronatis brunneis sterilibus et glumis 4-6 membranaceis oblongis obtusis fertilibus instructæ; stamina sub quaque gluma 6-12. Spiculæ foemineæ omnes ad apicem paniculi dispositæ, glumis 5 oblongo-lanceolatis acutis mucronatis ovarium solitarium stipitatum circumdatis; stigmata 3, dense villosa; pili hypogyni copiosi.

Leaves in tufts of 3-6, distinct from the flowering culms, 8-12 in. long, $1\frac{1}{4}$ - $1\frac{3}{8}$ lin. broad, linear, acute, often folded, but when expanded channelled down the face, with an acute ridge on each side of the channel, flat on the back, thinly pilose, usually slightly recurving. Flowering culms 14-18 in. high, including the panicle, $\frac{1}{2}$ - $\frac{3}{4}$ lin. broad, flattened, slightly concave on one side, convex on the other, sparingly pilose or nearly glabrous, produced on distinct leafless shoots, embraced by a long glabrous leafless sheath at the base. Panicle 7-12 in. long, strictly erect, linear, composed of 4-5 distant tufts of erect, simple or once umbellately divided, filiform branchlets, clustered in the axils of close-fitting brown sheaths $\frac{1}{3}$ - $\frac{3}{4}$ in. long, which have spreading leafy points $\frac{1}{2}$ -1 in. long. Spikelets solitary or very rarely 2 at the end of the branchlets, light brown. Male spikelets 3-4 lin. long, $1-1\frac{1}{4}$ lin. thick, lanceolate, with 5-7 coriaceous, oblong, acute, mucronate, brown empty glumes, thinly and minutely pubescent on the back, and 4-6 membranous, oblong, obtuse, fertile glumes; stamens 6-12 under each glume; anthers 2 lin. long, linear, tipped with a minute tuft of stiff hairs. Female spikelets all at the apex of the panicle, more slender than the males, 3 lin. long, about $\frac{2}{3}$ lin. thick, with about 5 oblong-lanceolate, acute, mucronate, minutely ciliate glumes surrounding the solitary pubescent ovary; stigmas 3, densely villose; hypogynous hairs copious.

Summit of Mount Roraima, 8600 ft., *McConnell & Quelch*, 676.

This species is easily distinguished from the following by its very much narrower leaves and less floriferous panicles.

EVERARDIA MONTANA, Ridl. in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 287.

Summit of Mount Roraima, *McConnell & Quelch*, 674.—Endemic.

GRAMINEÆ.

PANICUM ELIGULATUM, N. E. Brown, sp. n. Folia erecta, linearia, attenuato-acuta, rigida, cinereo-glaucæ, supra ad basin longe villosa et ad basi vaginorum villosotomentosa; ligula nulla. Culmi 10-13 poll. longi, stricti, striati, glabri, ad nodos

subglauci, bivaginati, vagina inferiore foliosa. Panicula 2-3 poll. longa, oblonga, multiflora. Spiculæ ellipsoideæ, 1-2-floræ. Glumæ 3 inferiores vacuæ, glabræ, brunneæ vel atro-fuscæ; gluma infima parva; secunda et tertia subæquales ellipticæ, obtusæ, profunde concavæ 6-7-nerves. Gluma florens elliptica, obtusa, profunde concava, lævis, 5-nervis, glabra, albida. Palea elliptica, obtusa, concava, 2-nervis, glabra, albida.

Leaves, excluding the sheaths, 6-9 in. long, $2\frac{1}{2}$ -3 lin. broad, linear, tapering to a very acute subpungent point, with involute margins, rigid, erect, greyish-glaucous, striate, more or less densely pilose on the upper side at the base, and densely villose-tomentose at the very base of the sheaths; ligule none. Culms 10-13 in. long, straight, striate, glabrous, more or less glaucous at the nodes, bearing two sheaths, the lower of which bears a short leaf. Panicle 2-3 in. long, $\frac{3}{4}$ -1 in. diam., oblong-lanceolate, moderately compact, with semiverticillate, erect, glabrous branchlets and pedicels. Spikelets $1\frac{1}{4}$ - $1\frac{1}{2}$ lin. long, ellipsoidal, 1-2-flowered. Empty glumes 3, glabrous, brown or blackish; the lowest $\frac{3}{4}$ -1 lin. long, $\frac{1}{3}$ lin. broad, oblong or elliptic-oblong, obtuse, 2-nerved; the 2 inner $1\frac{1}{4}$ lin. long, $\frac{1}{2}$ lin. broad, elliptic, obtuse, deeply boat-shaped, 6-7-nerved. Flowering glume 1 lin. long, elliptic, obtuse, deeply concave, 5-nerved, glabrous, smooth, whitish. Pale similar to the flowering glume, but smaller and flatter, 2-nerved, whitish. Anthers $\frac{3}{4}$ lin. long, very dark purplish brown. Stigmas 2, densely plumose, brown or fuscous.

Summit of Mount Roraima, 8600 ft., *McConnell & Quelch*, 675.

Allied to *Panicum loreum*, Trin., but the leaves and panicle are shorter than they are in that species, the spikelets rather larger, and the leaves quite destitute of a ligule, the absence of which is its most remarkable character, since there are few grasses in which all trace of a ligule is absolutely wanting.

PANICUM CHNOODES, Trin. Gram. Panic. p. 211.

Roraima Range, 3500 ft., *McConnell & Quelch*, 709.—Also in Brazil.

ECHINOLÆNA HIRTA, Desv. Journ. Bot. i. (1813), p. 75.

E. scabra, H. B. & K. Nov. Gen. & Sp. i. p. 118, t. 38; Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 288.

Upper slopes of Mount Roraima, *McConnell & Quelch*, 74.—Throughout Guiana and Brazil. The only other species of this genus is a native of Madagascar.

OLYRA MICRANTHA, H. B. & K. Nov. Gen. & Sp. i. p. 199.

Roraima Range, 3500 ft., *McConnell & Quelch*, 708.—Widely dispersed in Brazil and Guiana.

ARUNDO RORAIMENSIS, N. E. Brown, sp. n. Planta $\frac{1}{2}$ -3 ped. alta, culmo ad apicem laxè folioso glabro. Folia linearia, complicata, apice oblique subacuta, subrigida, glabra vel minutissime scaberula, rare supra ligulam laxè pilosa, intra microscopice papillosa; ligula ad annulum dense pilosum reducta. Panicula 4-6 poll. longa, ramis pedicellisque scaberulis vel villosis. Spiculæ 3-5-floræ, rhachilla breviter pilosa inter flores articulata. Glumæ 2 inferiores vacuæ, subæquales, lineares, acutæ vel minute

trifidæ, 1-nerves, submembranaceæ, glabræ; glumæ florentes oblongæ, longe aristatæ, basi longe pilosæ, 5-nerves. Palea lineari-oblonga, bifida vel emarginata, 2-nervis, membranacea, marginibus inflexis.

Perennial, $1\frac{1}{2}$ –3 ft. (or more ?) high. Stems laxly leafy to the top, glabrous. Lower leaves 9–20 in. long, the upper gradually shorter, $1\frac{1}{2}$ –3 lin. broad, erect, slightly rigid, linear, complicate, very obliquely subacute, glabrous or occasionally very minutely and thinly scaberulous on the back, microscopically papillate on the upperside, sometimes pilose and ciliate with long hairs just above the ligule, which is reduced to a dense ring of hairs. Panicle slightly nodding, 4–6 in. long, $1\frac{1}{2}$ – $2\frac{1}{2}$ in. broad, not very dense, its branches and pedicels scaberulous or villose. Spikelets 8–9 lin. long, 3–5-flowered, the rachilla readily disarticulating between the flowers, which are hermaphrodite. Empty glumes 2, subequal, 3– $3\frac{1}{2}$ lin. long, linear, acute and mucronate or shortly bifid at the apex, 1-nerved, membranous, glabrous. Flowering glume (excluding the awn) 3 lin. long, oblong-lanceolate, very concave, villose on the basal part with white hairs 2 lin. long, 5-nerved, trifid at the apex, with the middle tooth (or nerve) running out into a straight scabrid awn about 4 lin. long. Pale $2\frac{1}{2}$ lin. long, oblong-linear, with inflexed margins, bifid or emarginate at the apex, 2-nerved, scabrous on the nerves, otherwise glabrous. Lodicules $\frac{1}{4}$ lin. long, broad and oblique, subentire or crenately 2-lobulate, glabrous. Stamens 3. Ovary about 1 lin. long, slightly fusiform, or subterete, glabrous. Styles 2, plumose for $\frac{2}{3}$ of their length, yellow or brown.

Summit of Mount Roraima, 8600 ft., *McConnell & Quelch*, 673.

Allied to *A. pilosa*, D'Urv.

ARUNDINARIA DEFLEXA, N. E. Brown, sp. n. Culmi superne ramosi, ramis erectis teretibus glabris foliosis. Folia deflexa, lanceolata, acuminata, glabra, marginibus scaberulis; ligula brevis, subtruncata; vagina ad apicem longe setosa. Panicula terminalis, erecta, angusta, laxa, glabra. Spiculæ angustæ, paucifloræ. Glumæ inferiores 3 vacuæ, gluma extima minimus, 2 sequentes florentibus subsimiles sed minores; glumæ florentes ovato-lanceolatæ, obtusæ, concavæ, 7-nerves, glabræ, minutissime ciliatæ. Palea angusta, brevissime bifida, bicarinata, carinis ciliatis.

Stems branching in the upper part. Branches very erect, close to and parallel with the main stem, terete, glabrous, leafy, nearly concealed by the leaf-sheaths. Leaves deflexed, 5– $5\frac{1}{2}$ in. long, 9–11 lin. broad, lanceolate, tapering to a very acute point, glabrous on both sides, scabrid on the margins, probably slightly glaucous beneath; primary veins not very distinct from the rest; sheaths closely embracing the stem, glabrous, striate, with a fringe of long bristles at the apex; ligule short, subtruncate, glabrous. Panicle 5–11 in. long, 1–2 in. broad, lax, with slender erect glabrous branches. Spikelets about 3-flowered, 5–6 lin. long, narrow, with the 3 lower glumes empty, the lowest of which is about 1 lin. long, the second and third $2\frac{1}{2}$ lin. long, oblong, obtuse, concave, glabrous, not ciliate, obscurely 5–7-nerved. Flowering glume $3\frac{1}{2}$ – $3\frac{3}{4}$ lin. long, ovate-lanceolate, obtuse, concave, glabrous, very minutely ciliate, 7-nerved. Pale 3 lin. long, linear-oblong, with inflexed sides, slightly bifid at the obtuse apex, with 2 keel-like ciliate nerves. Anthers $1\frac{1}{2}$ lin. long, linear.

Summit of Mount Roraima, 8600 ft., *McConnell & Quelch*, 678.

The spikelets on the specimens seen are imperfect, most of them consisting of empty bracts only, and none that I have examined contain a perfect ovary.

CHUSQUEA LINEARIS, N. E. Brown, sp. n. Culmi graciles, internodiis $\frac{3}{4}$ – $2\frac{1}{4}$ poll. longis, ramulis brevibus ad nodos aggregatis. Folia 6–10 lin. longa, 1 – $1\frac{1}{2}$ lin. lata, lanceolata, acutissima, glabra, secus unum marginem minute scaberula; vaginæ striatæ, pubescentes, ligula brevi subtruncati. Racemi ad nodos aggregati, lineares, spiculis 3–6, unifloris. Glumæ inferiores 4 vacuæ, gluma infima minima, sequentes majores pubescentes; gluma florens convoluta, obtusa, apicem tantum pubescens, 7–8-nervis. Palea oblongo-ovata, obtusa, convoluta, glabra, apice ciliata, 4-nervis.

Stems slender, terete, glabrous, with internodes $\frac{3}{4}$ – $2\frac{1}{4}$ in. long, concealed by the striated sheaths, bearing at the nodes clusters of 4–10 simple leafy or flowering branches $1\frac{1}{2}$ –3 in. long. Leaves distichous, those on the branches $1\frac{1}{2}$ –2 lin. distant, 6–10 lin. long, 1 – $1\frac{1}{2}$ lin. broad, lanceolate, tapering to a very acute mucronate point, abruptly narrowed into a very short petiole at the base, glabrous, minutely ciliate-scabrid along one margin, striate beneath from the numerous closely placed prominent nerves; sheaths striate, pubescent; ligule short, subtruncate, no bristles. Racemes linear, with 3–6 pedicellate, adpressed, 1-flowered spikelets, or the terminal one with 14–15 spikelets. Outer 4 glumes empty; the lowest $\frac{1}{2}$ – $\frac{3}{4}$ lin. long, broadly oblong, obtuse, 1-nerved; the second $1\frac{1}{4}$ – $1\frac{3}{4}$ lin. long, oblong, obtuse, with or without a short awn, 1-nerved; the two others subequal, $2\frac{1}{2}$ – $2\frac{3}{4}$ lin. long, oblong-lanceolate, acute, very concave, 5–6-nerved, all more or less pubescent on the back and ciliate, at least at the apex. Flowering glume $2\frac{3}{4}$ lin. long, ovate-lanceolate, obtuse, convolute, pubescent and ciliate at the apex only, with 7–8 slender nerves. Pale 2 lin. long, oblong-ovate, convolute, obtuse and ciliate at the apex, membranous, with 4 slender nerves. Lodicules 3, narrowly oblong, obtuse, ciliate at the apex, $\frac{2}{3}$ – $\frac{3}{4}$ lin. long. Stamens 3; anthers 1 lin. long, linear. Ovary narrowly ovoid, glabrous; styles 2, narrowly plumose nearly to the base.

Summit of Mount Roraima, 8600 ft., *McConnell & Quelch*, 677.

Allied to *C. abietifolia*, Griseb., but among other characters is easily distinguished by its linear racemes.

II. PTERIDOPHYTA.

By C. H. WRIGHT, A.L.S.

1. GLEICHENIA PUBESCENS, H. B. & K. Nov. Gen. & Sp. i. p. 29. *Mertensia pubescens*, Willd. ; R. Schomb. Reisen in Brit.-Guiana, ii. p. 272 : Appun, Unter den Tropen, ii. p. 594.
Roraima range, 3500 ft., *McConnell & Quelch*, 610, 619.
British Guiana : Kwating Creek ; also Mexico and West Indies to Chili.
2. CYATHEA VESTITA, Mart. in Denkschr. Regensb. ii. (1822), p. 146. *C. hirtula*, Mart. ; R. Schomb. Reisen in Brit.-Guiana, ii. p. 272 : Appun, Unter den Tropen, ii. p. 594.
Roraima range, 3500 ft., *McConnell & Quelch*, 605.
Humirida Mountains, Brazil.
3. HEMITELIA SUBINCISA, Kunze, in Bot. Zeit. ii. (1844), p. 296.
Roraima range, 3500 ft., *McConnell & Quelch*, 620.
British Guiana : Kwating Creek ; also Central and Tropical South America.
- ALSOPHILA MARGINALIS, Klotzsch, in Linnæa, xviii. (1844), p. 542 ; Appun, Unter den Tropen, ii. 594.
Roraima range, 3500 ft., *McConnell & Quelch*, 604.
Mexico.
5. ALSOPHILA VILLOSA, Presl, ex Hook. Sp. Fil. i. p. 43 ; Appun, Unter den Tropen, ii. p. 594 ; Baker, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 289. *Cyathea villosa*, H. B. & K. Nov. Gen. & Sp. i. p. 24, and vii. t. 670.
Roraima range, 3500 ft., *McConnell & Quelch*, 621.
British Guiana : Arapu River ; also throughout Tropical South America.
6. ALSOPHILA MICROPHYLLA, Klotzsch, in Linnæa, xviii. (1844), p. 541.
Roraima range, 3500 ft., *McConnell & Quelch*, 606.
Venezuela and New Granada.
7. DICKSONIA CONIFOLIA, Hook. Sp. Fil. i. p. 70, t. 24. fig. A.
Summit of Roraima, 8600 ft., *McConnell & Quelch*, 564.
Jamaica, Venezuela, New Granada, and South Brazil.
8. HYMENOPHYLLUM CRISPUM, H. B. & K. Nov. Gen. & Sp. i. p. 26.
Summit of Roraima, 8600 ft., *McConnell & Quelch*, 563.
Cuba, Jamaica, Mexico, Venezuela, New Granada, Ecuador, Peru, and Bolivia.

9. *HYMENOPHYLLUM DEJECTUM*, Baker, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 289; Hook. Ic. Pl. t. 1610.

Roraima: upper slopes, *McConnell & Quelch*, 46; summit, *McConnell & Quelch*, 110, 113.

Endemic. Mr. im Thurn gathered it on the summit, as stated on p. 269 of Trans. Linn. Soc. Ser. II. Bot. ii. (1887).

10. *HYMENOPHYLLUM POLYANTHOS*, Sw. Syn. Fil. p. 149; R. Schomb. Reisen in Brit.-Guiana, iii. p. 1044. *H. clavatum*, Sw.; R. Schomb. Reisen in Brit.-Guiana, ii. p. 272, iii. p. 1044; Appun, Unter den Tropen, ii. p. 594. *H. Pæppigianum*, Presl; R. Schomb. Reisen in Brit.-Guiana, iii. p. 1044; Appun, Unter den Tropen, ii. p. 594.

Roraima range, 3500 ft., *McConnell & Quelch*, 611; upper slopes, *McConnell & Quelch*, 50 p.p., 73; summit, 8600 ft., *McConnell & Quelch*, 624.

British Guiana: Kwating Creek, Potaro River and Mazaruni River; also West Indies, Central America, the entire South American continent, Tropical and Subtropical Asia, New Zealand, Mascarene Islands, and Tropical Africa.

11. *HYMENOPHYLLUM MICROCARPUM*, Desv. in Mém. Soc. Linn. Paris, vi. (1827), p. 333.

Roraima: upper slopes, *McConnell & Quelch*, 44, 69.

Portorico, Guatemala, New Granada, Ecuador, Peru, Bolivia, and South Brazil.

12. *HYMENOPHYLLUM SERICEUM*, Sw. Syn. Fil. p. 146.

Summit of Roraima, 8600 ft., *McConnell & Quelch*, 628.

West Indies, Guatemala, French Guiana, New Granada, Ecuador, Peru, Bolivia, and South Brazil.

13. *HYMENOPHYLLUM LINEARE*, Sw. Fl. Ind. Occ. iii. p. 1749.

Roraima: upper slopes, *McConnell & Quelch*, 55 pro parte; summit, 8600 ft., *McConnell & Quelch*, 562.

Tropical America from the West Indies and Mexico southward to Peru and Brazil, Tropical and South Africa, and Mascarene Islands.

14. *HYMENOPHYLLUM* sp., *H. multifido*, Sw., affinis.

Mazaruni, between 250 and 3000 ft., *McConnell & Quelch*, 596, 598.

15. *TRICHOMANES ELEGANS*, Rudge, Pl. Guian. Rar. Ic. p. 24. *Hymenostachys elegans*, Presl; R. Schomb. Reisen in Brit.-Guiana, ii. p. 272; Appun, Unter den Tropen, ii. p. 594.

Roraima range, 3500 ft., *McConnell & Quelch*, 615; Mazaruni River, under 3000 ft., *McConnell & Quelch*, 594.

Central America, Trinidad, New Granada, Venezuela, and Northern Brazil.

In no. 594 the upper part of one of the fertile fronds is barren, and about six pairs of pinnae have grown out in a similar manner to those of the barren fronds.

16. *TRICHOMANES MACILENTUM*, Van den Bosch, *Hymenophyll. Nov.* p. 12; Baker in *Trans. Linn. Soc. Ser. II. Bot. ii.* (1887), p. 289.

Mazaruni River, 250 ft., *McConnell & Quelch*, 578; between 250 and 3000 ft., *McConnell & Quelch*, 597.

British Guiana: upper slopes of Roraima; forests near the Kaieteur Savannah, Potaro River; Amutu, below the Kaieteur Fall, Potaro River; Macouria River; also in Trinidad and Brazil, especially the northern provinces.

This is regarded by Mr. Baker (*Trans. Linn. Soc. loc. cit.*) as a variety of *T. Bancroftii*, Hook. & Grev.

17. *TRICHOMANES BANCROFTII*, Hook. et Grev. *Ic. Fil. t.* 204; R. Schomb. *Reisen in Brit.-Guiana*, iii. p. 1043; Appun, *Unter den Tropen*, ii. p. 594.

Mazaruni River, between 250 and 3000 ft., *McConnell & Quelch*, 587.

British Guiana: Amutu and below the Kaieteur Fall on the Potaro River; also in the West Indies, Guatemala, French Guiana, Northern Brazil, and Peru.

18. *TRICHOMANES BRACHYPUS*, Kunze, in *Linnæa*, ix. (1834), p. 105; R. Schomb. *Reisen in Brit.-Guiana*, ii. p. 272, iii. p. 1043; Appun, *Unter den Tropen*, ii. p. 594. *T. Ankersii*, Parker; R. Schomb. *loc. cit.*; Appun, *loc. cit.*

Roraima range, 3500 ft., *McConnell & Quelch*, 609.

British Guiana: in the forest just below the Kaieteur Fall, Potaro River; Amutu, Potaro River; Tumbana-Capra; also in St. Vincent, Trinidad, Guatemala, French Guiana, Venezuela, New Granada, and Brazil.

19. *TRICHOMANES PYXIDIFERUM*, Linn., var. *EMARGINATUM*, Hook. et Baker, *Syn. Fil.* p. 81.

Mazaruni, between 250 and 3000 ft., *McConnell & Quelch*, 599.

British Guiana: upper slopes of Roraima; also in the Andes of Quito. The type occurs throughout the tropics of both hemispheres.

20. *TRICHOMANES RADICANS*, Sw. *Fl. Ind. Occ.* iii. p. 1736.

Roraima range, 3500 ft., *McConnell & Quelch*, 612.

Extending from the Southern United States, through Central America and the West Indies, to Southern Brazil and Bolivia; also in Western Europe, Atlantic Islands, West Africa, Mascarene Islands, Northern India, China, Japan, and Polynesia.

21. *TRICHOMANES RORAIMENSE*, Jenm. in *Gard. Chron. Ser. III. xx.* (1896), p. 716.

Summit of Roraima, 8600 ft., *McConnell & Quelch*, 108, 561.

22. *TRICHOMANES CRISPUM*, Linn. *Sp. Pl. ed. 2*, p. 1560; Baker, in *Trans. Linn. Soc. Ser. II. Bot. ii.* (1887), p. 289.

Roraima: upper slopes, *McConnell & Quelch*, 63; Roraima range, 3500 ft., *McConnell & Quelch*, 616.

British Guiana: Essequibo River; Berbice (coast); Corentyne River; forests near Kaieteur Savannah; Kaieteur Ravine; also West Indies, Central America, French Guiana, Venezuela, New Granada, Ecuador, Peru, Bolivia, Paraguay, throughout Brazil, and West Tropical Africa.

23. *TRICHOMANES PINNATUM*, Sw. Syn. Fil. p. 142. *Neurophyllum pinnatum*, Presl; R. Schomb. Reisen in Brit.-Guiana, iii. p. 1044.

Mazaruni, between 250 and 3000 ft., *McConnell & Quelch*, 585; Roraima range, 3500 ft., *McConnell & Quelch*, 614.

British Guiana: Potaro River, below the Kaieteur Fall; Kanaku Mountains; Karawanu Mountains; Corentyne River; Berbice; also in the West Indies, Guatemala, French Guiana, Venezuela, New Granada, Peru, Northern Brazil, and Chili.

24. *TRICHOMANES PRIEURII*, Kunze, Analect. Pterid. p. 48; R. Schomb. Reisen in Brit.-Guiana, ii. p. 272, iii. p. 1043. *T. rigidum*, Appun, Unter den Tropen. ii. p. 594.

Roraima range, 3500 ft., *McConnell & Quelch*, 613.

British Guiana: Kaieteur Ravine; Potaro River, below the Kaieteur Fall; also in the West Indies, French Guiana, Venezuela, New Granada, Ecuador, and Brazil from the Amazon region southward to Santa Catherina.

25. *TRICHOMANES GEMMATUM*, J. Sm. in Hook. Journ. Bot. iii. (1841), p. 417. *T. cellulolum*, Klotzsch in Linnæa, xviii. (1844), p. 531; R. Schomb. Reisen in Brit.-Guiana, ii. p. 272, iii. p. 1043.

Roraima range, 3500 ft., *McConnell & Quelch*, 608.

British Guiana: Kanaku Mountains; Amutu, Potaro River; Pacatout, below the Kaieteur Fall, Potaro River; also in Venezuela, Northern Brazil, Polynesia, the Philippines, Borneo and the Malay Peninsula.

26. *LINDSAYA DUBIA*, Spreng. Syst. Veg. iv. p. 79; R. Schomb. Reisen in Brit.-Guiana, iii. p. 1048.

Mazaruni, between 250 and 3000 ft., *McConnell & Quelch*, 593.

British Guiana: Roraima; Pacatout, below the Kaieteur Fall, Potaro River; Amutu, Potaro River; also in French Guiana.

27. *LINDSAYA TRAPEZIFORMIS*, Dryand. in Trans. Linn. Soc. iii. (1797), p. 42, var.

Roraima range, 3500 ft., *McConnell & Quelch*, 617.

Intermediate between var. *laxa*, Baker, and var. *arcuata*, Baker (in Mart. Fl. Bras. i. ii. p. 355). The type is widely diffused in Tropical America, Asia, and Australia.

28. *LINDSAYA STRICTA*, Dryand. in Trans. Linn. Soc. iii. (1797), p. 42; in Thurn and Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), pp. 269, 290.

Roraima: upper slopes, *McConnell & Quelch*, 55, 62, 76; summit, 8600 ft., *McConnell & Quelch*, 497, 557.

British Guiana: Kaieteur Savannah; Head of Iteibisci Lake; also in the West Indies, Central America, French Guiana, Venezuela, New Granada, Peru, Bolivia, and throughout Brazil.

29. *PTERIS AQUILINA*, Linn., var. *CAUDATA*, Hook. Sp. Fil. ii. p. 196; in Thurn, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 266.

Summit of Roraima, 8600 ft., *McConnell & Quelch*, 553.

Humirida Mountains; Florida, West Indies, Central America, Venezuela, Brazil, China, Malay Peninsula, and East Africa. The type is cosmopolitan.

30. *PTERIS PALMATA*, Willd. Sp. Pl. v. p. 357.

Kanuku Mountains, 1000 ft., *McConnell & Quelch*, 273.

British Guiana: Rupununi and Watu Ticaba; also in the West Indies, Central America, Venezuela, New Granada, Ecuador, Peru, Bolivia, Brazil, and New Guinea.

31. *PTERIS INCISA*, Thunb. Fl. Cap. ed. Schult. p. 733; Baker, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 290.

Roraima: upper slopes, *McConnell & Quelch*, 75.

Also at the bottom of the cliff, Roraima. West Indies, Central America, Venezuela, New Granada, Galapagos Islands, Ecuador, South Brazil, Juan Fernandez, Chiloe, Polynesia, Himalaya, Formosa, Ceylon, Tasmania, Mascarene Islands, South Africa, and West Tropical Africa.

32. *LOMARIA PROCERA*, Spreng. Syst. Veg. iv. p. 65; Baker, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 290.

Summit of Roraima, 8600 ft., *McConnell & Quelch*, 124, 566.

Throughout Tropical America from the West Indies and Mexico to Chili; also in Polynesia, Malaya, New Zealand, Tasmania, Southern Australia, Mascarene Islands, and South Africa.

33. *LOMARIA BORYANA*, Willd. Sp. Pl. v. p. 292; Baker, in Trans. Linn. Soc. Ser. II. ii. (1887), p. 290. *Lomaria Schomburgkii*, Klotzsch, in R. Schomb. Reisen in Brit.-Guiana, ii. p. 272, iii. p. 1050, and in Linnæa, xx. (1846), p. 346; Appun, Unter den Tropen, ii. p. 594.

Summit of Roraima, 8600 ft., *McConnell & Quelch*, 121.

West Indies, Venezuela, New Granada, Ecuador, Bolivia, South Brazil, Chili, Patagonia, Tristan d'Acunha, Tropical and South Africa, and the Mascarene Islands.

34. *ASPLENIUM SERRATUM*, Linn. Sp. Pl. ed. 1, p. 1078.

Roraima range, 3500 ft., *McConnell & Quelch*, 601.

British Guiana: Berbice; Kuyuni Creek; Oreala; also in Florida, West Indies, Central America, French Guiana, Venezuela, New Granada, Ecuador, Galapagos Islands, Peru, Bolivia, Paraguay, Brazil, and the Society Islands.

35. *ASPLENIUM LUNULATUM*, Sw., var. *ERECTUM*, Baker, in Mart. Fl. Bras. i. ii. p. 435 Oliver, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 290. *A. harpeodes*, Kunze, in Linnæa, xviii. (1844), p. 329; R. Schomb. Reisen in Brit.-Guiana, iii. p. 1051; Appun, Unter den Tropen, ii. p. 595.

Upper slopes of Roraima, *McConnell & Quelch*, 45.

West Indies, Central America, Venezuela, New Granada, Ecuador, Peru, Bolivia, Argentine, Paraguay, Brazil, New Caledonia, India, Mascarene Islands, Tropical Africa, South Africa, and the Atlantic Islands.

36. *ASPLENIUM SERRA*, Langsd. et Fisch. Pl. Voy. Russes, i. p. 16, t. 19; R. Schomb. Reisen in Brit.-Guiana, ii. p. 272, iii. p. 1051; Appun, Unter den Tropen, ii. p. 595.

Summit of Roraima, 8600 ft., *McConnell & Quelch*, 623.

British Guiana: Kwating Creek; also in Brazil, Venezuela, New Granada, Ecuador, Peru, New Caledonia, Tropical and South Africa, and the Mascarene Islands.

37. *ASPLENIUM AURITUM*, Sw. Fl. Ind. Occ. iii. p. 1616; R. Schomb. Reisen in Brit.-Guiana, ii. p. 272, iii. p. 1051.

Roraima range, 3500 ft., *McConnell & Quelch*, 607.

West Indies, Central America, Venezuela, New Granada, Ecuador, Galapagos Islands, Peru, Bolivia, Brazil, South India, Madagascar, and East Tropical Africa.

38. *ASPIDIUM MENISCIODES*, Willd. Sp. Pl. v. p. 218.

Roraima range, 3500 ft., *McConnell & Quelch*, 618.

British Guiana: Mazaruni Bush; Potaro River; Kabalebo River; Berbice; Essequibo River; also in Trinidad, French Guiana, Brazil, and Peru.

39. *NEPHRODIUM CONTERMINUM*, Desv. in Mém. Soc. Linn. Paris, ii. (1827), p. 255; Baker, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 290.

Summit of Roraima, 8600 ft., *McConnell & Quelch*, 627.

British Guiana: upper slopes of Roraima; also in Florida, West Indies, Central America, and South America southward to Chili and the Argentine.

This form has the ultimate segments more obtuse than the type.

40. *NEPHRODIUM DENTICULATUM*, Hook. Sp. Fil. iv. p. 147; Baker, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 290.

Summit of Roraima, 8600 ft., *McConnell & Quelch*, 622.

Roraima: on the upper slopes and near the base of the cliff; also in the West Indies, Central America, Venezuela, New Granada, and Brazil.

41. *POLYPODIUM* (§ *GRAMMITIS*) *CONNELLII*, Baker, n. sp.; ad *P. gramineum*, Sw., accedit; differt frondibus glabris longioribus margine revolutis, soris haud contiguis regulariter biseriatis.

A densely caespitose herb. Rhizome scarcely creeping; paleæ lanceolate, brown,

membranous. Frond linear, simple, entire, 3-4 in. long, 1-1½ lin. wide, rigidly coriaceous, glabrous, gradually contracted into a very short, but distinct, stipe; veins immersed, concealed. Sori oblong, superficial, parallel with the midrib, often confluent.

Summit of Roraima, 8600 ft., *McConnell & Quelch*, 111, 118, 570.

42. *POLYPODIUM MARGINELLUM*, Sw. Fl. Ind. Occ. iii. p. 1631; Baker, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 291. *Mecosorus marginellus*, var. *major*, Klotzsch; R. Schomb. Reisen in Brit.-Guiana, iii. p. 1054.

Roraima: upper slope, *McConnell & Quelch*, 51; summit, 8600 ft., *McConnell & Quelch*, 117, 119, 568.

West Indies, Venezuela, Peru, Brazil, Samoa, and St. Helena.

Swartz (Fl. Ind. Occ. iii. p. 1631) says this species has "venis bifidis"; subsequent authors describe the veins as simple, which is the case in all the specimens at Kew.

43. *POLYPODIUM LEPTOPODON*, C. H. Wright, n. sp.; ad *P. marginellum*, Sw. accedit; differt fronde membranacea, marginibus ciliatis, venis unifurcatis, stipite elongato tenui piloso.

Rhizome shortly creeping, 2 lin. in diam., with brown lanceolate scales near the apex. Frond oblanceolate, obtuse, up to 2 in. long by 3 lin. broad, glabrous except for a few cilia on the margin when young, edged with a black line; veins once forked, conspicuous; sori few, near the apex of the frond, often confluent; stipe up to 2½ in. long, slender, pilose.

Summit of Roraima, 8600 ft., *McConnell & Quelch*, 569.

44. *POLYPODIUM FURCATUM*, Mett. in Abhandl. Senck. naturf. Ges. ii. (1858), p. 34; Baker, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 291.

Mazaruni, 250 ft., *McConnell & Quelch*, 576; between 250 and 3000 ft., *McConnell & Quelch*, 584.

British Guiana: upper slope of Roraima; below and above the Kaieteur, Potaro River; also in French Guiana, North Brazil, and the Island of Grenada.

45. *POLYPODIUM SERRULATUM*, Mett. Fil. Hort. Bot. Lips. p. 30; Baker, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 291. *Xiphopteris serrulata*, Kaulf.; R. Schomb. Reisen in Brit.-Guiana, ii. p. 272, iii. p. 1056.

Mazaruni, between 250 and 3000 ft., *McConnell & Quelch*, 592.

British Guiana: upper slopes of Roraima; also in the West Indies, Central America, Venezuela, New Granada, Ecuador, Peru, Bolivia, Brazil, Sandwich Islands, Mascarene Islands, and West Tropical Africa.

Var. *STRICTISSIMUM*, Hook. Sp. Fil. iv. p. 175.

Summit of Roraima, 8600 ft., *McConnell & Quelch*, 555.

Ecuador.

46. *POLYPODIUM TOVARENSE*, Klotzsch, in *Linnæa*, xx. (1846), p. 374; Baker, in *Trans. Linn. Soc. Ser. II. Bot. ii.* (1887), p. 291.
Roraima: upper slopes, *McConnell & Quelch*, 67.
Venezuela, New Granada, and Ecuador.
47. *POLYPODIUM MONILIFORME*, Lag. in *Sw. Syn. Fil.* p. 33; Baker, in *Trans. Linn. Soc. Ser. II. Bot. ii.* (1887), p. 291.
Roraima: upper slopes, *McConnell & Quelch*, 47, 66; summit, 8600 ft., *McConnell & Quelch*, 558
West Indies, Central America, Andes of New Granada, Ecuador, and Bolivia.
48. *POLYPODIUM HARTII*, Jenm. in *Journ. Bot.* xxiv. (1886), p. 272.
Mazaruni, between 250 and 3000 ft., *McConnell & Quelch*, 591.
Jamaica, Dominica, and Island of Grenada.
49. *POLYPODIUM TRICHOMANOIDES*, Sw. *Prodr. Veg. Ind. Occ.* p. 131; R. Schomb. *Reisen in Brit.-Guiana*, iii. p. 1052; Appun, *Unter den Tropen*, ii. p. 595; Baker, in *Trans. Linn. Soc. Ser. II. Bot. ii.* (1887), p. 291.
Roraima: upper slopes, *McConnell & Quelch*, 49; Mazaruni, *McConnell & Quelch*, 575.
British Guiana: Kaieteur Ravine and Savannah; also in the West Indies, Central America, French Guiana, Brazil (Prov. Bahia), Ecuador, Peru, Juan Fernandez, Ascension Island, Northern India, and East Tropical Africa.
50. *POLYPODIUM CULTRATUM*, Willd. *Sp. Pl.* v. p. 187; R. Schomb. *Reisen in Brit.-Guiana*, ii. p. 272, iii. p. 1052; Baker, in *Trans. Linn. Soc. Ser. II. Bot. ii.* (1887), p. 292.
Roraima: upper slopes, *McConnell & Quelch*, 54, 56, 71.
British Guiana: Kaieteur Savannah; also in the West Indies, Central America, Venezuela, New Granada, Peru, Bolivia, Brazil, West Tropical Africa, and the Mascarene Islands.
51. *POLYPODIUM XANTHOTRICHUM*, Klotzsch, in *Linnæa*, xx. (1846), p. 376; Baker, in *Trans. Linn. Soc. Ser. II. Bot. ii.* (1887), p. 292.
Roraima: upper slopes, *McConnell & Quelch*, 64, 68.
Endemic.
52. *POLYPODIUM CAPILLARE*, Desv. in *Berl. Ges. naturf. Freunde, Mag.* v. (1811), p. 316; Baker in *Trans. Linn. Soc. Ser. II. Bot. ii.* (1887), p. 292.
Roraima: upper slopes, *McConnell & Quelch*, 53, 58.
British Guiana: in woods near the Kaieteur Savannah; also in the West Indies, Venezuela, Ecuador, Peru, and Brazil.
53. *POLYPODIUM KALBREYERI*, Baker, in *Trans. Linn. Soc. Ser. II. Bot. ii.* (1887), p. 291.
Roraima: upper slopes, *McConnell & Quelch*, 65, 72.
New Granada.

54. *POLYPODIUM RIGESCENS*, Bory, in Willd. Sp. Pl. v. p. 183; Baker, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 292.

Roraima: upper slopes, *McConnell & Quelch*, 48, 70; summit, 8600 ft., *McConnell & Quelch*, 559.

British Guiana: Kwating Creek; also in the West Indies, Venezuela, New Granada, Ecuador, Peru, Brazil, Solomon Islands, Mascarene Islands, and Tropical Africa.

55. *POLYPODIUM ELASTICUM*, Bory, in Willd. Sp. Pl. v. p. 183.

Makarapan Mt. on the north of Kwaimatta, 2000 ft., *McConnell & Quelch*, 278.

British Guiana: Kwating Creek; Berbice; also in Florida, the West Indies, Central America, New Granada, Ecuador, Peru, Bolivia, Paraguay, and Brazil.

56. *POLYPODIUM MERIDENSE*, Klotzsch. in Linnæa, xx. (1846), p. 380.

Summit of Roraima, 8600 ft., *McConnell & Quelch*, 498, 499, 560.

Venezuela, New Granada, South Brazil.

7. *POLYPODIUM INCANUM*, Sw. Syn. Fil. p. 35.

Kanaku Mts., 500 ft., *McConnell & Quelch*, 271.

British Guiana: Savannah Bush; Rupununi; Corentyne River; also in the Southern United States, West Indies, Central America, French Guiana, Venezuela, New Granada, Peru, Brazil, Paraguay, Uruguay, Chili, South Africa, and the Zambesi.

58. *POLYPODIUM CRASSIFOLIUM*, Linn. Sp. Pl. ed. 1, p. 1083; R. Schomb. Reisen in Brit.-Guiana, iii. p. 1053; Appun, Unter den Tropen, ii. p. 595.

Roraima range, 3500 ft., *McConnell & Quelch*, 603.

West Indies, Central America, French Guiana, Surinam, Venezuela, New Granada, Ecuador, Galapagos Islands, Peru, Bolivia, Brazil.

59. *GYMNOGRAMME CYCLOPHYLLA*, Baker, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 293, t. 53.

Summit of Roraima, 8600 ft., *McConnell & Quelch*, 116, 120, 567.

Endemic.

60. *GYMNOGRAMME ELAPHOGLOSSOIDES*, Baker, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 293, t. 54.

Summit of Roraima, 8600 ft., *McConnell & Quelch*, 112, 554.

Endemic.

61. *GYMNOGRAMME PUMILA*, A. Spreng. Tent. Suppl. ad Syst. Veg. p. 31; R. Schomb. Reisen in Brit.-Guiana, iii. p. 1054.

Mazaruni, 250 ft., *McConnell & Quelch*, 491.

British Guiana: Rupununi River; also in the West Indies, Central America, New Granada, and Brazil.

62. GYMNOGRAMME FLEXUOSA, Desv. in Mém. Soc. Linn. Paris, ii. (1827), p. 215^l; Baker, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 293.

Roraima: upper slopes, *McConnell & Quelch*, 59; summit, 8600 ft., *McConnell & Quelch*, 629.

Central America, Venezuela, New Granada, Ecuador, Peru, Bolivia, and Brazil.

63. ACROSTICHUM SIMPLEX, Sw., var. MARTINICENSE, Hook. et Baker, Syn. Fil. ed. 1, p. 400.

Mazaruni, between 250 and 3000 ft., *McConnell & Quelch*, 581.

British Guiana: Essequibo River and on the coast; also in the West Indies, New Granada, Ecuador, Peru, and Brazil.

64. ACROSTICHUM CONFORME, Sw. Syn. Fil. pp. 10, 192, t. 1. fig. 1.

Mazaruni, 250 ft., *McConnell & Quelch*, 577.

British Guiana; Essequibo River and on the coast; also in the West Indies, Central America, Venezuela, New Granada, Ecuador, Peru, Chili, Brazil, Polynesia, Queensland, India, Mascarene Islands, Tropical and South Africa.

The form of this polymorphic species represented in this collection has been named *Acrostichum alatum*, Fée (Mém. Fam. Fougèr. ii. p. 35, t. 5. fig. 2).

65. ACROSTICHUM LATIFOLIUM, Sw. Fl. Ind. Occ. iii. p. 1589; Baker, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 294.

Summit of Roraima, 8600 ft., *McConnell & Quelch*, 571.

British Guiana: upper slopes of Roraima; Asabaru Creek; Sheenabowa and Kaieteur Ravine, Potaro River; Demerara River and on the coast; also throughout the West Indies and Tropical America, and widely diffused in Polynesia, Java, Mascarene Islands, Tropical and South Africa.

66. ACROSTICHUM SQUAMOSUM, Sw. in Schrad. Journ. Bot. iv. (1800), p. 11; Baker, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 295.

Roraima: upper slopes, *McConnell & Quelch*, 57.

British Guiana: Kaieteur Savannah and Essequibo River; also in the West Indies, Central America, Venezuela, Ecuador, Brazil, Sandwich Islands, Ceylon, South India, Mascarene Islands, Tropical Africa, and the Atlantic Islands.

67. ACROSTICHUM DECORATUM, Kunze, in Linnæa, ix. (1834), p. 25; Baker, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 294.

Roraima range, 3500 ft., *McConnell & Quelch*, 602.

West Indies, Peru, and South Brazil.

68. ACROSTICHUM PELTATUM, Sw. Fl. Ind. Occ. iii. p. 1593; Appun, Unter den Tropen, ii. p. 595; Baker, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 295.

Roraima: upper slopes, *McConnell & Quelch*, 61; Mazaruni, between 250 and 3000 ft., *McConnell & Quelch*, 582.

West Indies, Central America, Venezuela, New Granada, Ecuador, and Brazil.

69. *ACROSTICHUM OLIGARCHICUM*, Baker, in Hook. et Baker, Syn. Fil. ed. 1, p. 418.
Roraima range, 3500 ft., *McConnell & Quelch*, 600.
North Peru.
70. *SCHIZÆA FLUMINENSIS*, Miers, in Mart. Fl. Bras. i. pars II. p. 184, t. 15. fig. 2.
Mazaruni, between 250 and 3000 ft., *McConnell & Quelch*, 580.
British Guiana : Kukenaam, and woods at the side of the Kaieteur Savannah ; also in Venezuela and Brazil.
71. *SCHIZÆA ELEGANS*, Sw. Syn. Fil. p. 151 ; Baker, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 295.
Mazaruni, between 250 and 3000 ft., *McConnell & Quelch*, 589.
British Guiana : Essequibo River ; Epiro Creek, Corentyne River ; Demerara River, and on the coast ; also in the West Indies, Central America, Surinam, Venezuela, New Granada, and Brazil.
72. *SCHIZÆA PENNULA*, Sw. Syn. Fil. pp. 150, 379. *Actinostachys pennula*, Hook. ; R. Schomb. Reisen in Brit.-Guiana, iii. p. 1045.
Mazaruni, between 250 and 3000 ft., *McConnell & Quelch*, 588.
British Guiana : Roraima ; Kukuya Creek ; Kaieteur Savannah ; Serra Mey and Demerara River ; also in the West Indies, Surinam, Brazil, and New Caledonia.
73. *ANEMIA TOMENTOSA*, Sw. Syn. Fil. p. 157 ; Baker, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 295.
Kanuku Mountains, 1000 ft., *McConnell & Quelch*, 272, 274.
- Var. *FULVA*, Hook. et Baker, Syn. Fil. ed. 1, p. 433.
Ireng Valley, *McConnell & Quelch*, 201.
British Guiana : Roraima and Rupununi ; also in the West Indies, Central America, Venezuela, New Granada, Peru, Paraguay, Brazil, and Uruguay.
74. *LYCOPodium ALOPECUROIDES*, Linn. Sp. Pl. ed. 2, p. 1565 ; Baker, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 295 ; Jenman, in Timehri, v. (1886), p. 44.
Roraima : upper slopes, *McConnell & Quelch*, 60.
Southern United States, Venezuela, New Granada, Peru, Brazil, Bolivia, and Uruguay.
75. *LYCOPodium CERNUUM*, Linn. Sp. Pl. ed. 2, p. 1566 : Jenman, in Timehri, v. (1886) p. 45.
Ireng Valley, *McConnell & Quelch*, 206.
British Guiana : Corentyne, Pomeroon and Macouria Rivers and at Berbice ; also throughout the tropics and extending a short distance on either side.
76. *LYCOPodium CONTIGUUM*, Klotzsch, in Linnæa, xviii. (1844), p. 519.
Summit of Roraima, 8600 ft., *McConnell & Quelch*, 572.
Higher parts of the Andes in Venezuela, New Granada, Ecuador, and Bolivia.

77. *LYCOPODIUM CLAVATUM*, Linn. Sp. Pl. ed. 2, p. 1564; Jenman, in Timehri, v. (1886), p. 44.

Roraima range, 3500 ft., *McConnell & Quelch*, 707.

Arctic and alpine zones of both hemispheres, also on the mountains throughout the tropics.

78. *LYCOPODIUM CAROLINIANUM*, Linn. Sp. Pl. ed. 2, p. 1567; Jenman, in Timehri, v. (1886), p. 43.

Roraima : upper slopes, *McConnell & Quelch*, 52.

British Guiana: Kaieteur Savannah; Corentyne River; Kako Creek; Demerara; also in Florida, Guadeloupe, Brazil, Paraguay, Hongkong, Ceylon, Mascarene Islands, and South Africa.

79. *SELAGINELLA VERNICOSA*, Baker, in Trans. Linn. Soc. Ser. II. Bot. ii. (1887), p. 295, t. 56. f. A.

Summit of Roraima, 8600 ft., *McConnell & Quelch*, 574, ex parte.

Also found by im Thurn at the base of the cliff.

Var. *OLIGOCLADA*, Baker, *l. c.* fig. B.

Summit of Roraima, 8600 ft., *McConnell & Quelch*, 574, ex parte.

80. *SELAGINELLA FLABELLATA*, Spring, Monogr. Lycopod. p. 174.

Essequibo River, *McConnell & Quelch*, 115.

British Guiana : foot of the Kaieteur. Tropics and subtropics of America, Asia, and Polynesia.

81. *SELAGINELLA ANCEPS*, A. Br. in Ann. Sc. Nat. Sér. V. iii. (1865), p. 278.

Roraima range, 3500 ft., *McConnell & Quelch*, 706.

British Guiana : Potaro River; also in Guatemala, Venezuela, New Granada, Ecuador, and Peru.

III. BRYOPHYTA.

MUSCI by V. F. BROTHÉRUS, Ph.D.

DICRANACEÆ.

DICRANUM LONGISETUM, Hook. Musc. Exot. t. 139.

Mount Roraima, *McConnell & Quelch*, 343.

Found in the mountains of Venezuela, New Granada, and Quito.

Var. *LAXIFOLIUM*, Broth. n. var. Folia laxius disposita, apice argutius serrulata.

Mount Roraima, Ledge, 7500-8000 ft., *McConnell & Quelch*, 344.

DICRANODONTIUM PULCHRO-ALARE, Broth., n. sp. *Dioicum*, caespitosum, caespitibus densiusculis lutescentibus nitidis; *caulis* 8 cm., sterilis usque ad 12 cm. altus, arcuato-flexuosus, fusco-tomentosus, densiuscule foliosus, sæpe innovationibus elongatis gracilibus. *Folia* falcata, canaliculato-concava, e basi lanceolata longe subulata, usque ad 1 cm. longa, superne dense et argute serrulata, nervo basi folii latitudinis tertiam partem vel paulum ultra occupante et superne dorso serrulato ibidemque folium totum occupante; cellulæ superiores angustæ; basilares ad nervum laxæ, rectangulares, hyalinae; exteriores angustissimæ, limbum multi-seriatum efformantes; alares in ventrem magnum fugacem dispositæ, laxæ, teneræ, fuscae. *Bractea perichætii* internæ e basi vaginante subito in subulam integram attenuatæ, cellulis omnibus elongatis angustis teneris. *Seta* 2·5 cm. alta, tenuis, serpentino-flexuosa, lutea, apice cygnea, rubra. *Theca* sicca erecta, plicata, humida ob setam apice cygneam horizontalis, lævis, oblonga, c. 2·5 mm. alta, leptodermis, fuscidula. *Peristomium* simplex; *exostomii* dentes 16, aurantiaci, c. 0·37 mm. longi et c. 0·075 mm. lati, usque ad basin in cruribus duobus filiformibus inæqualibus articulatis dense papillosis divisi; *spori* 0·017 mm., fusci, papillosoi. *Cætera* ignota.

Mount Roraima, Ledge, 7500–8000 ft., *McConnell & Quelch*, 345, 346.

Endemic.

CAMPYLOPUS CHIONOPHILUS, Mitt. in Journ. Linn. Soc., Bot. xii. (1869), p. 81.

Dicranum chionophilum, C. Muell. Syn. i. 398.

Mount Roraima, Ledge, 7500–8000 ft., *McConnell & Quelch*, 342.

Found in the mountains of New Granada.

CAMPYLOPUS RORAIMÆ, Broth., n. sp. *Dioicus*, caespitosus, caespitibus densis nigrescentibus superne lutescentibus etiamque nitidiusculis; *caulis* ad 3 cm. usque altus, tomentosus, dense et per totam longitudinem æqualiter foliosus, ramosus. *Folia* homomallula, erecto-patentia, canaliculato-concava, e basi oblongo-lanceolata, breviter subulata, c. 5 mm. longa et c. 0·75 mm. lata, marginibus summo apice serratis, nervo basi dimidiam partem folii latitudinis occupante, lamina usque ad apicem folii distincta; cellulæ basilares subrectangulares, inter se porosæ, marginem versus angustiores, dein rhomboideæ; superiores rhombeæ, incrassatæ; alares numerosæ, subquadratae, rubræ, basin totam occupantes. *Cætera* ignota.

Mount Roraima, Ledge, 7500–8000 ft., *McConnell & Quelch*, 347.

Endemic.

Species *C. concolori*, Brid. affinis, sed foliis multo brevioribus primo visu dignoscenda.

CAMPYLOPUS ATRATUS, Broth., n. sp. *Dioicus*, robustus, ater, apice lutescens, nitidiusculus. *Caulis* 10 cm. altus, erectus, haud tomentosus, dense et per totam longitudinem æqualiter foliosus, parce ramosus; *folia* sicca suberecta, humida recurvulo-patentia, canaliculato-concava, superne tubulosa, e basi lanceolata longe et anguste subulata, pilo stricto tenui hyalino longiusculo serrato terminata,

c. 10 mm. longa et c. 1.1 mm. lata, marginibus integris superne conniventibus, nervo basi tertiam partem folii latitudinis occupante, lamina usque ad apicem folii distincta; cellulæ elongatæ, angustæ, marginem versus incrassatæ, lumine angustissimo, limbum inferne latiusculum hyalinum superne tenuiorem, ultra medium evanidum efformantes, alares numerosæ, subquadratae, rubræ, in ventrem distinctum dispositæ.

Cætera ignota.

Summit of Mount Roraima, 8600 ft., *McConnell & Quelch*, 527.

Endemic.

Species cum *Campylopode griseo*, Hornsch., comparanda, sed statura multo robustiore, foliis strictipilis, integris, tubulosis, limbatis facillime dignoscenda.

LEUCOBRYACEÆ.

LEUCOBRYUM MEGALOPHYLLUM, Mitt. in Journ. Linn. Soc., Bot. xii. (1869), p. 112.

Dicranum megalophyllum, Raddi, Critt. Bras. p. 3. *Leucobryum giganteum*, C. Muell. Syn. i. p. 79, ii. p. 536. *Leucobryum robustum*, Sull. in Proc. Amer. Acad. (1861), p. 279.

Mount Roraima, Ledge, 7500–8000 ft., *McConnell & Quelch*, 351 (forma).

Found in Cuba, Jamaica, Trinidad, Barbados, New Granada, Venezuela, and Brazil.

LEUCOBRYUM CRISPUM, C. Muell. Syn. i. p. 78.

Roraima range at 3500 ft., *McConnell & Quelch*, 546.

Found in the West-Indian Islands, Venezuela, New Granada, and Brazil.

LEUCOBRYUM LÆVIFOLIUM, Broth. *Dioicum*, robustiusculum, cæspitosum, cæspitibus mollibus dilabentibus lutescenti-albidis ætate fusciscentibus nitidis; *caulis* usque ad 6 cm. altus, erectus, dense foliosus, plerumque fasciculatim ramosus, ramis fastigiatis. *Folia* sicca et humida suberecta, stricta vel rarius homomallula, canaliculato-concava, e basi oblongo-lanceolata sensim acuminata, acuta, 5–6 mm. longa et c. 0.66 mm. lata, dorso sicca et humida lævissima, marginibus superne involutaceis integerrimis, limbata, limbo hyalino, inferne e seriebus cellularum 7–8 formato, superne angustiore, apice obsoleto, lamina e stratis cellularum æqualium duobus composita. *Cætera ignota.*

Summit of Mount Roraima, 8600 ft., *McConnell & Quelch*, 505, 536.

Species *L. Martiano*, Hampe, affinis, sed statura robustiore, nitore foliisque strictis, raro indistincte homomallulis oculo nudo jam dignoscenda.

OCTOBLEPHARUM MITTENII, Jaeg. Adumbr. i. p. 169. *Octoblepharum longifolium*, Mitt. in Journ. Linn. Soc., Bot. xii. (1869), p. 110.

Summit of Mount Roraima, 8600 ft., *McConnell & Quelch*, 517.

Found in Brazil.

TORTULACEÆ.

LEPTODONTIUM CIRRHIFOLIUM, Mitt. in Journ. Linn. Soc., Bot. xii. (1869), p. 52.

Summit of Mount Roraima, 8600 ft., *McConnell & Quelch*, 511.

Found in the mountains of Quito.

GRIMMIACEÆ.

MACROMITRIUM MUCRONIFOLIUM, Schwaegr. Suppl. ii. II. i. p. 167, t. 172.

Roraima range, 3500 ft., *McConnell & Quelch*, 544.

Found in Florida, in the West-Indian Islands, Guiana, Surinam, and Brazil.

ZYGODON SUBDENTICULATUS, Hampe, Ann. Sc. Nat. Sér. V. iv. (1863-67), p. 326.

Summit of Mount Roraima, *McConnell & Quelch*, 540.

Found in New Granada.

FUNARIACEÆ.

FUNARIA CALVESCENS, Schwaegr. Suppl. i. II. p. 77, t. 65.

Mount Roraima, Ledge, 7500-8000 ft., *McConnell & Quelch*, 338.

Found in South Europe and in the tropical and subtropical regions of Asia, Africa, America, and Australia.

BARTRAMIACEÆ.

BREUTELIA SCOPARIA, Besch. in Ann. Sc. Nat. Sér. VI. iii. (1876), p. 209. *Bartramia scoparia*, Schwaegr. Suppl. iii. I. i. t. 211.

Summit of Mount Roraima, 8600 ft., *McConnell & Quelch*, 514.

Found in the West-Indian Islands.

MNIACEÆ.

RHIZOGONIUM LINDIGII, Mitt. in Journ. Linn. Soc., Bot. xii. (1869), p. 328. *Mnium Lindigii*, Hampe, in Ann. Sc. Nat. Sér. V. iv. (1863-67), p. 345.

Summit of Mount Roraima, 8600 ft., *McConnell & Quelch*, 502, 537.

Found in Costa Rica and in the mountains of New Granada.

MNIUM ROSTRATUM, Schrad. in Linn. Syst. Nat. 13 ed. Gmel. ii. pars II. p. 1330.

Roraima, Ledge, 7500-8000 ft., *McConnell & Quelch*, 339.

Found in all parts of the world.

NECKERACEÆ.

RHACOCARPUS HUMBOLDTII, Lindb. in Öfvers. Svensk. Vet.-Ak. Förh. (1862). *Har- risonia Humboldtii*, Spreng. Syst. Veg. iv. i. p. 145.

Mount Roraima, Ledge, 7500-8000 ft., *McConnell & Quelch*, 336, and summit of Mount Roraima, 519, 534 (var. *fusco-viridis*).

Found in the mountains of New Granada.

PILOTRICHUM BIPINNATUM, Brid. Bryol. Univ. ii. p. 283.

Roraima range, 3500 ft., *McConnell & Quelch*, 548.

Found in the West-Indian Islands, in Guiana, Brazil, and in the mountains of Quito and Peru.

HOOKERIACEÆ.

HOOKERIA (§ **HYLOTAPIS**) **PILOTRICHELLOIDES**, Broth., n. sp. *Robustiuscula*, mollis, fusco-aurea, nitida; *caulis* usque ad 15 cm. longus, flexuosus, complanatus, dense foliosus, per totam longitudinem pinnatim ramosus, ramis vix ultra 1 cm. longis patentibus strictis vel arcuatulis complanatis dense foliosis obtusis. *Folia* sicca imbricata, humida suberecta, cymbiformi-concava, oblonga vel ovato-oblonga, in apiculum acutum, recurvulum subito contracta, marginibus inferne revolutis, integerrimis vel apice minutissime crenulatis, nervis binis elongatis vix divergentibus longe infra apicem evanidis dorso lævibus; cellulæ angustissime lineares, seriatim papillosæ; basilares læves; infimæ laxæ, breves, fusco-aureæ. *Cætera* ignota.

Mount Roraima, Ledge, 7500–8000 ft., *McConnell & Quelch*, 337.

Endemic.

Species pulcherrima, mollitie necnon caule longissimo *Pilotrichellis* nonnullis similis, cum nulla alia commutanda.

HOOKERIA (§ **OMALIADELPHUS**) **RORAIMÆ**, Broth., n. sp. *Dioica*, robusta, lutescenti-viridis, ætate fusco-aurea, nitida; *caulis* usque ad 20 cm. longus, complanatus, dense foliosus, dense pinnatim ramosus, ramis patulis, 1–1.5 cm. longis, strictis, complanatis, dense foliosis, obtusis; *folia* sicca laxè imbricata, valde undulata, humida erecto-patentia, concaviuscula; *lateralia* oblongo-lanceolata, breviter acuminata, acutiuscula, marginibus inferne revolutis dein erectis minute serrulatis, nervis binis tenuibus parce divergentibus infra apicem evanidis dorso superne denticulatis; cellulæ elongatæ, angustissime lineares; basilares infimæ laxæ, fusco-aureæ, omnes lævissimæ; *bracteæ perichætii* internæ e basi ovata sensim subulatæ, minute denticulatæ, enerves; *seta* 1.5 cm. vel paulum ultra alta, sat tenuis, rubra, lævissima; *theca* horizontalis, e collo brevi ovalis, sicca deoperculata sub ore haud constricta, fusca; *peristomium* duplex; *exostomii* dentes e basi lanceolata longissime subulati, c. 0.95 mm. longi et c. 0.14 mm. lati, latiuscule exarati, dense et alte lamellati, aurantiaci; *endostomium* sordide luteum, minute papillosum; *processus* carinati, vix perforati; *operculum* longe et anguste subulatum; *calyptra* ad medium thecæ producta, basi multifida, apice fusca, glabra.

Roraima range, 3500 ft., *McConnell & Quelch*, 490, 550.

Endemic.

Species pulcherrima, statura robusta ab omnibus speciebus sectionis oculo nudo jam dignoscenda.

STEREODONTACEÆ.

ECTROPOTHECIUM AMABILE, Mitt. in Journ. Linn. Soc., Bot. xii. (1869), p. 514.

Mount Roraima, Ledge, 7800–8000 ft., *McConnell & Quelch*, 341, and summit of Mount Roraima, 8600 ft., 516, 542.

Found in the mountains of New Granada.

CTENIDIUM MALACODES, Mitt. in Journ. Linn. Soc. Bot. xii. (1869), p. 509.

Mount Roraima, Ledge, 7500–8000 ft., *McConnell & Quelch*, 340.

Found in the mountains of New Granada and Quito.

LESKEACEÆ.

THUIDIUM ANTILLARUM, Besch. in Ann. Sc. Nat. Sér. VI. iii. (1876), p. 244.

Roraima range, 3500 ft., *McConnell & Quelch*, 549.

Found in Costa Rica and the West-Indian Islands.

THUIDIUM PSEUDO-PROTENSUM, Mitt. in Journ. Linn. Soc., Bot. xii. (1869), p. 578.

Hypnum pseudo-protensum, C. Muell. in Bot. Zeit. vi. (1848), col. 779.

Mount Roraima, Ledge, 7500–8000 ft., *McConnell & Quelch*, 348.

Found in Venezuela.

SPHAGNACEÆ.

SPHAGNUM SANGUINALE, Warnst. in Bot. Centralbl. lxxvi. (1898), p. 385.

Mount Roraima, Ledge, 7500–8000 ft., *McConnell & Quelch*, 350, 541 in part.

Endemic.

SPHAGNUM MEDIUM, Limpr. in Bot. Centralbl. vii. (1881), p. 313.

Mount Roraima, Ledge, 7500–8500 ft., *McConnell & Quelch*, 349, 541 in part.

Found in Europe and in America from Labrador to Patagonia.

HEPATICÆ. By F. STEPHANI.

The collection of Liverworts made by Messrs. McConnell and Quelch is a small one, but is of particular interest from a geographical point of view. Many plants were found which hitherto had been only observed in the Andes of South America; their unexpected appearance on the top of Mount Roraima is quite startling: the curious *Frullania mirabilis*, Jack et Steph., is of particular interest, as well as the very rare and beautiful *Pleurozia paradoxa*, Jack, both of which up to this time had not been elsewhere collected.

It is possible that some of these plants may have intermediate stations of which we are at present ignorant; but a similar surprising collection was made by Mr. Ule (of the Botanic Garden of Rio Janeiro) in the Serra do Mar and the Serra Itatiaia, where Andine forms were collected, though stations forming connecting-links are altogether missing.

There are also several new species, amongst which *Metzgeria inflata* is one of the most curious in the genus, being quite hairless and almost without any rootlets, lying like a small inflated cylindric pouch amongst other mosses.

As cryptogamic plants, and in particular Liverworts, with few exceptions, have very small spores and live in sheltered and moist places from where the wind cannot easily

carry away the spores, they are a better indication than phænogamic plants as to the Flora we must regard as the remnant of another, which, like that of South America, once covered a wide area, and is now separated by intermediate barren plains or treeless Llanos and Pampas of enormous extension where no Liverworts can live.

That the spores are not able to propagate a Liverwort beyond, perhaps, a small area, no better example can show than the very interesting flora of Killarney, where—doubtless for many centuries—a few species of Hepaticæ of tropical origin have been preserved without being able to reach the Continent.

The present condition of the literature of Liverworts seldom allows one to make geographical speculations; but when my “Species Hepaticarum” is finished, and the number and relation of known liverworts, as also their distribution, settled, we may be able to draw conclusions which, I think, will be of value to all who study the geographical distribution of plants.

The following Liverworts have been collected, viz. :—

A. ANACROGYNÆ.

1. DUMORTIERA HIRSUTA, Nees, Hep. Eur. iv. p. 163. *Marchantia hirsuta*, Sw. Prodr. Fl. Ind. Occ. p. 145.
Roraima, Ledge, 7500–8000 ft., *McConnell & Quelch*, 334/6.
Common in Tropical and Subtropical America, England, Italy, Africa.
2. ANEURA SCHWANECKEI, Steph. in Hedwigia xxvii. (1888), p. 278.
Summit of Roraima, *McConnell & Quelch*, 512.
Found also in the West-Indian Islands.
3. ANEURA ALGOIDES, Steph. in Bull. Herb. Boiss. vii. (1899), p. 682. *Metzgeria algoides*, Taylor, in Hook. Lond. Journ. Bot. v. (1846), p. 410.
Summit of Roraima, *McConnell & Quelch*, 529.
An Andine plant, collected by Jameson near Quito.
4. ANEURA RORAIMENSIS, Steph., n. sp. *Dioica*, minor, pallide olivacea, aliis hepaticis consociata. *Frons* ad 5 mm. longa, exalata, irregulariter bipinnata, sæpe subfasciculata. *Truncus* primarius angustus, biconvexus, ramis trunco latoribus minus convexus. *Cuticula* ubique lamellata, lamellis denticulatis margine bene prominulis. *Cellulæ* frondis internæ corticalibus multo majores, fronde in adspectu itaque optime reticulata. *Rami feminei* in trunco solitarii, breves, margine latissime alati, alis profunde inciso-lobatis, lobis lanceolatis vel ligulatis obtusis, squama basali similiter lobata ramulum ♀ a tergo tegente. *Reliqua* desunt.

A very good and most distinct species, easily to be recognized by the rough cuticula of the cortical cells.

In the monograph of the genus *Aneura* ('Species Hepaticarum,' p. 736) it is to be placed after No. 81, *Aneura scabra*, Steph.

Summit of Roraima, *McConnell & Quelch*, 350. Mixed with *Micropterygium pterygophyllum*, Spruce.

5. ANEURA BREUTELII, Steph. in Bull. Herb. Boiss. vii. (1899), p. 759.
Summit of Roraima, *McConnell & Quelch*, 533.
Collected before by Breutel in the West-Indian Island of St. Kitts.

6. ANEURA FUCOIDES, Steph. in Bull. Herb. Boiss. vii. (1899), p. 680. *Jungermannia fucoides*, Sw. Prodr. fl. Ind. occ. p. 45.
Roraima, Ledge, 7500–8000 ft., *McConnell & Quelch*, 334/6 in part.
Common in Tropical America.

7. METZGERIA INFLATA, Steph., n. sp. *Dioica*, spectabilis, brevis, flaccida, pallide lurida, optime nitida, muscis consociata. *Frons* furcata, ad 3 cm. longa, omnino nuda, basi tantum paucis radicellis affixa. *Costa* tenuis, cellulis corticalibus utroque latere biseriatis tecta et in sectione transversa late ovalis; cellulæ antice et postice magnæ, æquales, internæ multo minores triseriatae. *Alæ* apice cucullatae, ubique maxime irregulariterque revolutæ, in sectione transversa valde asymmetricæ, uno latere quadruplo latiores, re vera itaque sinuatim lobatae, lobis bulloso inflatis ob flexuram tamen haud discretis. *Cellulæ* alarum 36–54 μ , parietibus validis, trigonis parvis. *Rami feminei* parvi, fere conduplicati, margine breviter setosi.
Summit of Roraima, *McConnell & Quelch*, 524.
This plant is well adapted to retain the water which it takes up from the mosses on which it is growing in rather exposed places on the very summit of the mountain; it may be compared to the Tasmanian species *M. saccata*, Mitt., which prefers to grow on the uppermost twigs of trees, where it has a similar exposed situation and shows a similar adaptation to retain the water within the inflated lobes of the thallus.

8. METZGERIA HAMATA, Lindb. Monogr. Metzg. p. 25.
Roraima, Ledge, 7500–8000 ft., *McConnell & Quelch*, 334/4 and 5.
Common in all parts of the globe.

9. MONOCLEA GOTTSCHKEI, Lindb. in Rev. Bryol. xiii. (1886), p. 102, in adnot.
Roraima, at 3500 ft., *McConnell & Quelch*, 551.
West-Indian Islands and the northern part of South America, Chile, Peru.
This plant has been distributed by Gottsche and others under the name of *Monoclea Forsteri*, a very different plant and altogether antarctic (New Zealand and Patagonia).

10. PALLAVICINIUS WALLISII, J. B. Jack & Steph. in Hedwigia, xxxi. (1892), p. 23.
Summit of Roraima, *McConnell & Quelch*, 501.
Collected before by Wallis in the mountains of New Granada near Antioquia, 8000 ft.

B. ACROGYNÆ.

11. JAMESONIELLA COLORATA, Spruce, in Journ. Bot. xvi. (1876), p. 30. *Jungermannia colorata*, Lehm. in Linnæa, iv. (1829), p. 366.
Summit of Roraima, *McConnell & Quelch*, 539 in part.
America, Australia, South Africa. Very common.

12. SYZYGIELLA QUELCHII, Steph., n. sp. *Sterilis* magna, fusco-purpurascens, tenera et flaccida, aliis hepaticis sparsim consociata. *Caulis* ad 8 cm. longus simplex (semper?). *Folia* opposita, conferta, assurgentia, si aliam decurrentem anticam excipis oblique falcato-ovata, apice duplo angustiora quam basi, oblique truncata vel lenissime emarginata, omnino integerrima, angulis solum dentiformibus; *marginē postico* valde arcuato, basi folii oppositi brevissime coalito ibidemque *hamatim auriculato*, *marginē antico* substricto breviter recurvo, longe decurrente, folia tamen antice haud attenuata sed oblique truncata et abrupte desinentia, folio opposito minime coalita. *Cellulæ* foliorum nodulose incrassatæ, nodulis maximis nusquam confluentibus, multo majores quam in *S. manca*, Steph.
Roraima, Ledge, 7500–8000 ft., *McConnell & Quelch*, 334/14 in part.
A most beautiful and very curious plant; the two opposite leaves are quite erect and touch each other, the stem being perfectly hidden when seen from above. At the postical base these two leaves are united for a very short space; just above this point they are rounded off and much projecting, and have a large incurved acuminate lobule.

13. SYZYGIELLA PERFOLIATA, Spruce, in Trans. Edinb. Bot. Soc. xv. (1885), p. 500, in adnot. *Jungermannia perfoliata*, Sw. Prodr. Veg. Ind. Occ. p. 143.
Summit of Roraima, *McConnell & Quelch*, 524 in part.
Found in Venezuela and the West Indies.

14. PLAGIOCHILA AËREA, Taylor, in Hook. Lond. Journ. Bot. v. (1846), p. 263.
Roraima, Ledge, 7500–8000 ft., *McConnell & Quelch*, 334.
Found before in Venezuela and Mexico.

15. PLAGIOCHILA GAVANA, Steph., n. sp. *Dioica*, spectabilis, pallide glauco-virens, ætate flavo-virens, effuse cæspitans. *Caulis* ad 10 cm. longus, longe remoteque pauciramosus, ramis late divergentibus, apice sæpe fasciculatis ubique æqualiter foliatis. *Folia* magna, tenera, remotiuscula, basi tantum imbricata, subrecte patula, breviter inserta, *i. e.* basi antica et postica fere opposita, breviter decurrentia, ceterum ovato-oblonga, apice fere triplo angustiora quam basi, *marginē antico* substricta recurva subintegerrima, *marginē postico* arcuata versus basin ampliata, rotundata ibidemque cum folio opposito ad cristam erectam conniventia, longe spinosa, spinis sub 30 magnis approximatis angustis longe acuminatis recte patulis 7 cellulas longis basi 2–4 cellulas latis. Folia *cellulæ* $34 \times 50 \mu$, trigonis majusculis; basales $34 \times 80 \mu$, parietibus æqualiter incrassatis. *Perianthia* in caule terminalia, geminatim innovata, foliis parum longiora, plus duplo angustiora quam longa, antice late alata,

apice recte truncata ut in ala longe spinosa, spinis ut in folio sed magis confertis. *Folia floralia* caulinis simillima. *Andræcia* in caule terminalia, longe spicata, spicis ad 2 vel 3 subfasciculatis erectis; bractæ ad 16 jugæ, e basi longe saccata parum recurvo-patulæ, apice truncatæ et paucidenticulatæ.

Roraima, Ledge, 7500–8000 ft., *McConnell & Quelch*, 334/9 in part.

In *Plagiochila superba*, Wahlenb., the leaf-cells are much larger, the spines very stout, very much longer and less numerous. Lindenb. says (Monogr. Plag. p. 80) that Sieber had given him this species as coming from Australia; but this is a mistake of Sieber, who had not marked his packets and did not know on his return whence he had procured the plants; many errors of the same kind have been previously pointed out by me.

16. *PLAGIOCHILA REMOTIFOLIA*, Hampe & Gottsche in Linnæa, xxv. (1852), p. 340.

Roraima, Ledge, 7500–8000 ft., *McConnell & Quelch*, 334/3.

Pretty common in the West Indian Islands.

17. *PLAGIOCHILA RUTILANS*, Lindenb. Spec. Hep. ii. & iii. (Monogr. Hep. Gen. Plag.) p. 47.

Roraima, Ledge, 7500–8000 ft., *McConnell & Quelch*, 334/2 in part.

Common in Tropical America.

18. *LEIOSCYPHUS FRAGILIS*, J. B. Jack & Steph. in Hedwigia, xxxi. (1892), p. 20.

Summit of Roraima, *McConnell & Quelch*, 524 in part.

Also in New Granada.

19. *LOPHOCOLEA BREUTELII*, Gottsche, in Syn. Hep. p. 154.

Roraima, Ledge, 7500–8000 ft., *McConnell & Quelch*, 334/14 in part.

Common in the West Indian Islands.

20. *MASTIGOBRYUM RORAIMENSE*, Steph., n. sp. *Dioica*, mediocris, apice dilute flavo-virens, inferne brunneola, erecta, laxè intricata. *Caulis* ad 4 cm. longus, repetito furcatus, postice flagellis numerosis capillaceis instructus. *Folia* conferta, valde decurva, in plano subrecte patula, oblonga, parum falcata, antice parum ampliata, caulemque haud superantia, apice triplo angustiora quam basi, acute bidentula sinu lunato parum profundo vel truncata, angulis acutis. *Cellulæ* foliorum magnæ, apice 27 μ , trigonis maximis contiguis vel late confluentibus, medio infero multo majores, 37–56 μ , incrassatio parietum grosse trabeculata. *Amphigastria* caulina caule triplo fere latiora, imbricata, transverse inserta, late obtusata, tertio supero inciso biloba, lobis similiter sed brevius bilobis; *cellulæ* amphigastriæ caulinis simillimæ, parum minores. *Folia floralia* quadrijuga, intima caulinis æquilonga, profundissime bifida, laciniis late lanceolatis, marginibus externis crenatis vel crenato-dentatis, internis subintegerrimis. *Amphigastrium florale* foliis suis æquale. *Perianthium* (juvenile) ore longe fimbriatum, ciliis triangulariter articulatis hic illic ramosis. Reliqua desunt.

Summit of Roraima, *McConnell & Quelch*, 523.

21. *MASTIGOBRYUM DISSODONTUM*, Steph. n. nomen; *Bazzania bidens* var. *dissodonta*, Spruce, in Trans. Edinb. Bot. Soc. xv. (1885), p. 371.
Summit of Roraima, *McConnell & Quelch*, 532.
Found in Colombia.
22. *MASTIGOBRYUM PORTORICENSE*, Hampe & Gottsche in Linnæa, xxv. (1852), p. 348.
Roraima at 3500 ft., *McConnell & Quelch*, 545.
Common in the West Indian Islands and northern South America.
23. *MASTIGOBRYUM VINCENTINUM*, Lehm. & Lindenb. Syn. Hep. p. 226.
Roraima, Ledge, 7500–8000 ft., *McConnell & Quelch*, 334/10, 15, & 16.
Common in the West Indian Islands.
24. *MASTIGOBRYUM KRUGIANUM*, Steph. in Hedwigia, xxvii. (1888), p. 300.
Roraima, Ledge, 7000–8000 ft., *McConnell & Quelch*, 334/14 in part.
Found before in the island of Santo Domingo.
25. *MASTIGOBRYUM GRACILE*, Hampe & Gottsche, in Linnæa, xxv. (1852), p. 346.
Roraima, Ledge, 7500–8000 ft., *McConnell & Quelch*, 334/11 in part.
Known from different places in the West Indian Islands.
26. *MICROPTERYGIUM GRANDISTIPULUM*, Steph., n. sp. *Sterilis*, minor, dilute brunnea vel flavo-rufescens, muscicola, valde intricatim cæspitans. *Caulis* tenuis, ad 2 cm. longus, irregulariter ramosus. *Folia* conferta, decurvula, semiamplexicaulia, ex angusta basi ovato-oblonga, longe acuminata, apice abrupte attenuata ibidemque paucidenticulata vel integerrima. *Folii* lobulus posticus integerrimus, brevi spatio (ad $\frac{1}{4}$) carinatim coalitus, folium subinde oblique percurrens, attenuatus, longe sub folii apice desinens. *Cellulæ* foliorum læves, $8 \times 12 \mu$, basales $8 \times 17 \mu$, subrectangulatæ, maxime æqualiterque incrassatæ. *Amphigastria* magna, imbricata, foliis parum breviora, transverse inserta, plana, integerrima, medio infero fere circularia, superne abrupte angustata, lanceolata, acuta.
Roraima, Ledge, 7500–8000 ft., *McConnell & Quelch*, 334/14 in part.
There is not a species of this genus which does not have some of its leaves merely conduplicate, the wing being entirely wanting; where the latter is developed it is part of the antical lobe, which can easily be observed if a transverse cut of the leaf is made; the same occurs sometimes in *Scapania*, and is best developed in *Schistochila*.
27. *MICROPTERYGIUM PTERYGOPHYLLUM*, Spruce, in Trans. Edinb. Bot. Soc. xv. (1885), p. 384. *Jungermannia pterygophylla*, Nees, in Mart. Fl. Bras. vol. i. pars prior, Algæ, Lichenes, Hepaticæ, p. 377.
Summit of Roraima, *McConnell & Quelch*, 530 in part.
Common in Tropical America.
28. *LEPIDOZIA COMMUTATA*, Steph. in Hedwigia, xxvii. (1888), p. 293.
Summit of Roraima, *McConnell & Quelch*, 521.
Very common in the West Indian Islands.

29. *LEPIDOZIA LAXEPINNATA*, Spruce, in Trans. Edinb. Bot. Soc. xv. (1885), p. 360.
 Roraima at 3500 ft., *McConnell & Quelch*, 552.
 An Andine species not found anywhere else before.
30. *SCHISMA JUNIPERINUM*, Nees, Hep. Eur. iii. p. 575. *Jungermannia juniperina*, Sw.
 Veg. Ind. Occ. iii. p. 1855.
 Summit of Roraima, *McConnell & Quelch*, 522, 528, 538, 539; in the last associated to *Jamesoniella colorata*, Spruce.
 Very common in Tropical America.
31. *SCHISMA PENSILIS*, Steph., n. nomen. *Sendtnera pensilis*, Taylor, in Hook. Lond. Journ. Bot. v. (1846), p. 372.
 Roraima, Ledge, 7500–8000 ft., *McConnell & Quelch*, 334/2 in part.
 A pretty rare Andine species.
32. *SCHISMA DURANDII*, Steph., n. sp. *Sterilis* longissima, gracilis, lurida, longe lateque stratificata et dense cæspitosa. *Caulis* ad 20 cm. longus, fragilis, simplex, superne sæpe fasciculatim ramosus, apice breviter circinatus, postice flagellifer, flagellis brevibus approximatis parvifoliis. *Folia* pro planta parva, triplo longiora quam lata, ad $\frac{2}{3}$ bifida, laciniis lanceolatis apice longe setaceis valde hamatis valdeque divergentibus. *Cuticula* grosse verrucosa. *Cellulæ* foliorum subapicales $17 \times 25 \mu$, parietibus longioribus et marginalibus trabeculatim incrassatis. *Cellulæ* basales $25 \times 70 \mu$, oblongo-hexagonæ, trigonis magnis sæpe trabeculatim confluentibus. *Flagellorum* folia multoties minora, ovato-oblonga, laxè accumbentia, porrecta, ad $\frac{2}{3}$ bifida, laciniis late lanceolatis vix setaceis.
 Roraima, Ledge, 7500–8000 ft., *McConnell & Quelch*, 334/2, 334/11 & 334/14, all in part.
 This plant has been found before by Pittier in the higher mountains of Costa Rica.
33. *SCHISMA SUBDENTATUM*, Steph., n. sp. *Sterilis*, mediocris, fusco-purpurascens, optime nitens, densissime depresso-cæspitosa. *Caulis* ad 6 cm. longus, simplex vel furcatus, furcis divergentibus. *Folia* confertissima, maxime decurvo-hamata, angusta, plus 3-plo longiora quam lata, ad $\frac{2}{3}$ bifida, laciniis parum divergentibus, longe acuminatis, acutis. *Cellullæ* foliorum subapicales $17 \times 25 \mu$, parietibus longioribus valde incrassatis; incrassatio marginalis, præcipue ad angulos cellularum, grosse nodulose-vel subdentatim prominula. *Cellulæ* basales folii longe hexagonæ, $25 \times 68 \mu$, parietibus longioribus maxime trabeculatim incrassatis.
 Roraima, Ledge, 7500–8000 ft., *McConnell & Quelch*, 334/1.
34. *TRICHOCOLEA SPHAGNOIDES*, Steph., n. nomen. *Leiomitra sphagnoides*, Spruce, in Trans. Edinb. Bot. Soc. xv. (1885), p. 350.
 Roraima, Ledge, 7500–8000 ft., *McConnell & Quelch*, 334/9 in part, 11 in part, and 13.
 Andes of Quito.

35. *SCAPANIA PORTORICENSIS*, Hampe & Gottsche, in *Linnaea*, xxv. (1852), p. 342.
Summit of Roraima, *McConnell & Quelch*, 573.
Not rare in the West Indian Islands, and found by Spruce also on the top of Mount Abitagua, 2000 m. (his *Scapania splendida*).
36. *PLEUROZIA PARADOXA*, Steph., n. nomen.—*Physotium paradoxum*, J. B. Jack, in *Hedwigia*, xxvii. (1886), p. 85.
Summit of Roraima, *McConnell & Quelch*, 543.
Known before from Pasto, collected by F. C. Lehmann.
37. *HARPALEJEUNEA TENAX*, Steph., n. sp. *Sterilis*, minor, fusco-olivacea, rigida, dense caespitosa. *Caulis* ad 15 mm. longus, capillaceus, parum ramosus. *Folia* remotiuscula, caule vix latiora, parum patula, cauli fere parallela, valde concava, in plano cordiformia, tertio supero abrupte angustata acuminata acuta vix incurva, *lobulo* duplo angustiore, valde inflato, in acumen folii attenuato. *Cellulae* foliorum parvae, ad 12 μ , trigonis late confluentibus. *Amphigastria* parva, transverse inserta, caulem vix superantia subrotunda, appressa, integerrima.
Roraima, Ledge, 7500–8000 ft., *McConnell & Quelch*, 334/.
A much reduced form, rigid and evidently well adapted to a dry climate, the leaves being very small, little projecting from the stem, the cell-walls very thick. This plant is almost unique in the great number of species of *Lejeunea*, which always live in damp and well sheltered places, and disappear where a dry atmosphere can reach them.
38. *FRULLANIA MIRABILIS*, J. B. Jack & Steph. in *Hedwigia*, xxxi. (1892), p. 15.
Summit of Mount Roraima, *McConnell & Quelch*, 510.
Collected before by Wallis in New Granada, near Antioquia.
39. *FRULLANIA ATRATA*, Nees, in *Syn. Hep.* p. 463. *Jungermannia atrata*, Sw. *Prodr. Fl. Ind. Occ.* p. 144.
Summit of Roraima, *McConnell & Quelch*, 535.
Very common in Tropical America.
40. *FRULLANIA LONGICOLLIS*, Lindenb. & Gottsche, in *Syn. Hep.* p. 783.
Summit of Roraima, *McConnell & Quelch*, 335.
An Andine species, found also in Mexico.
-

IV. THALLOPHYTA.

FUNGI. By G. MASSEE, F.L.S.

PHOMA PSAMMISLÆ, Massee, n. sp. Maculæ nullæ; perithecia epidermide tecta, gregaria, 500 μ diam., atra, ostiolo papillato pertusa; sporæ ellipticæ, 2-guttulatæ, 10–11 $\times$ 3 μ .

Spots absent; perithecia gregarious, covered by the grey raised epidermis, lenticular, averaging 500 μ in diameter, black, glabrous; ostiolum minute, papillate; texture parenchymatous, dense, olive; spores hyaline, elliptical, ends obtuse with two minute polar guttulæ, 10–11 $\times$ 3 μ .

On dead branch of *Psammisia coriacea*, N. E. Brown. Summit of Roraima, *McConnell & Quelch*.

Allied to *P. stictica*, Berk. et Broome.

ECHINBOTRYUM ROSEUM, Massee, n. sp. Mycelium tenuissimum vix manifestum, album; hyphæ fertiles erectæ, septatæ, hyalinæ, 4 μ crassæ. Conidia in apicibus hypharum fertilium densius aggregata, sessilia, fusoidea, 15–17 $\times$ 6 μ , roseo-tincta.

Sterile hyphæ prostrate, hyaline, septate, branched, about 4 μ thick, loosely interwoven into a very thin white layer; fertile hyphæ erect, septate, simple, hyaline, 4 μ thick, apex minutely nodulose and producing numbers of sessile spindle-shaped, pale rose-coloured conidia measuring 15–17 $\times$ 6 μ .

Parasitic on *Octomeria parvifolia*, Rolfe. Summit of Roraima, *McConnell & Quelch*, 696.

Allied to *E. læve*, Sacc., from which the present differs in the form and colour of the spores.

STEMPHYLIUM ERICOCTONUM, A. Br. & de Bary, Krankh. d. Pflanz. (1854), p. 18, tab. 2.

On living branches of *Leitgebia Imthurniana*, Oliver. Summit of Roraima, *McConnell & Quelch*.

Europe.

MACROSPORIUM RAMULOSUM, Sacc. Fung. Ital. tab. 854 (1881).

On fading leaves of *Tofieldia Schomburgkiana*, Oliver. Summit of Roraima, *McConnell & Quelch*.

Europe, and in the Andes of Colombia.

CAPNODIUM FIBROSUM, Berk. in Hook. Fl. Nov. Zel. ii. (1855), p. 209.

Forming a black bristly pile on living twigs of *Psammisia*. Summit of Roraima, *McConnell & Quelch*, 662.

N. Zealand.

RHIPIDONEMA MEMBRANACEUM, Sacc. Syll. vi. (1888), p. 688.

On bark. Roraima, 8000 ft., *McConnell & Quelch*, 508, 526.

Amazon Valley; Marianne Islands.

LICHENES. By G. MASSEE, F.L.S.

SPHÆROPHORON COMPRESSUM, Ach. Meth. Lich. (1803), p. 135.

On stones. Summit of Roraima, *McConnell & Quelch*, 506.

Cape Horn; Chili; W. Indies; Europe; India; Ceylon.

CLADONIA RANGIFERINA, Hoffm. Flor. Germ. (1795), p. 114.

On the ground. Summit of Roraima, *McConnell & Quelch*, 507 & 520.

Cosmopolitan.

PARMELIA DICTYORHIZA, Massee, n. sp. Thallus membranaceo-cartilagineus, substellatus, repetito dichotomo-laciniatus, cretaceus, subtus niger, spongiosus. Apothecia ignota.

Thallus thin, cartilaginous, radiating from a centre in an irregularly stellate manner, repeatedly dichotomously divided, chalk-white above, black and spongy below, the spongy stratum composed entirely of closely septate brown hyphæ 11–12 μ thick, arranged in the form of an irregular network. Apothecia unknown. Laciniæ about 2 mm. wide.

On the ground. Summit of Roraima, *McConnell & Quelch*, 53.

Allied to *P. hypotropa*, Nyl., but separated from this and every other described species in the reticulate structure of the spongy lower surface of the thallus.

PARMELIA PERFORATA, Ach. Lich. Univers. (1810), p. 410.

On branches. Summit of Roraima, *McConnell & Quelch*, 503 & 513.

Central and N. America; Central Africa; Polynesia.

ALECTORIA OCHROLEUCA, Nyl. Prod. Lich. (1857), p. 46.

On branches. Summit of Roraima, *McConnell & Quelch*, 504.

Mexico; United States; Europe; Himalaya; Japan; Australia.

EXPLANATION OF THE PLATES.

PLATE 1.

Ilex apicidens. (Figs. 1 to 6.)

Figs. 1 & 2. Branches with lax and dense inflorescences, natural size.

Fig. 3. Flower, $\times 4$.

4. Flower with petals and stamens removed, $\times 4$.

5. Stamen, dorsal view, $\times 4$.

6. Stamen, front view, $\times 4$.

Cyrilla brevifolia. (Figs. 7 to 16.)

Figs. 7 & 8. Two branches, natural size.

Fig. 9. Fragment of a raceme with one flower, $\times 5$.

10. Sepal, seen from within, $\times 7$.

11. Petal, seen from within, $\times 7$.

12. Stamens, front and side views, $\times 7$.

13. Flower, with the outer parts removed to show the ovary, $\times 7$.

14. A trifold stigma, $\times 7$.

15. Ovary, longitudinal section, $\times 12$.

16. Ovaries, transverse sections, $\times 12$. The ridges on the ovaries may be due to shrinkage, but they are represented as seen after long boiling.

PLATE 2.

Macairea aspera.

Fig. 1. Branch, natural size.

2. Fragment of leaf, upper surface, $\times 4$.

3. Flower, with the petals and six stamens removed, $\times 5$.

4. Petal, $\times 5$.

5. Transverse section through the calyx-tube and ovary, $\times 5$.

Figs. 6 & 7. Long and short stamens from another specimen (no. 24, *McConnell & Quelch*), $\times 5$.

PLATE 3.

Passiflora Quelchii.

Fig. 1. Branch, natural size.

2. Inflorescence, natural size.

3. Longitudinal section of a flower, $\times 2$.

4. Filament of the lower corona, $\times 5$.

PLATE 4.

Henriquezia Jenmani.

- Fig. 1. Leaf and inflorescence, natural size.
 2. Flower, with the corolla removed, natural size.
 3. Corolla, laid open, natural size.
 4. Ovary after the sepals have fallen, $\times 3$.
 5. Ovary, longitudinal section, $\times 3$.
 6. Ovary, transverse section, $\times 3$.

PLATE 5.

Didymochlamys Connellii. (Figs. 1 to 9.)

- Fig. 1. Part of a plant seen from above, natural size.
 2. Part of a plant seen from beneath, natural size.
 3. Inflorescence, with one bract removed, $\times 2$.
 4. Flower with 2 sepals and the corolla removed, $\times 5$.
 5. Corolla, laid open, $\times 2$.
 6. A lobe of the corolla, $\times 4$.
 7. Ovary, with glands and calyx-lobes, longitudinal section, $\times 5$.
 8. Ovary, longitudinal section, showing placentas, $\times 5$.
 9. Ovary, transverse section, $\times 5$.

Chalepophyllum Connellii. (Figs. 10 to 17.)

Figs. 10 & 11. Two branchlets showing the range of variation in different specimens, natural size, but the stems are not stout enough.

- Fig. 12. Calyx and ovary, natural size.
 13. Part of a corolla laid open, natural size.
 14. Stamens, dorsal and front views, $\times 2$.
 15. Ovary, longitudinal section, $\times 4$.
 16. Ovary, transverse section, $\times 4$.
 17. Ovule, $\times 12$.

PLATE 6.

Retiniphyllum laxiflorum.

- Fig. 1. Branch, natural size.
 2. Flower, $\times 2$.
 3. Apex of a pedicel, $\times 5$.
 4. Longitudinal section through the lower part of a flower, $\times 5$.
 5. Three views of a stamen, $\times 5$.
 6. Ovary, transverse section, $\times 5$.
 7. Longitudinal section of an ovarian cell, $\times 20$.
 8. Transverse section of an ovarian cell, $\times 20$.
 9. Oblique dorsal view of the cap-shaped placenta and ovules, $\times 20$.
 10. Oblique front view of the placenta and one ovule, the other ovule removed, $\times 20$.
 11. Fruit, $\times$ about $2\frac{1}{2}$.
 12. Fruit, transverse section, $\times 3$.

PLATE 7.

Heterothalamus densus. (Figs. 1 to 7.)

- Fig. 1. Branch of a male plant, natural size.
 2. Branch of a female plant, natural size.
 3. Involucre of a male head of flowers, $\times 5$.
 4. Involucre of a female head of flowers, $\times 5$.
 5. Male floret, $\times 5$.
 6. Bract from the receptacle of a female head, $\times 5$.
 7. Female floret, $\times 5$.

Quelchia conferta. (Figs. 8 to 14.)

- Figs. 8 & 9. Two branches, showing variation in the length of the peduncles, natural size.
 Fig. 10. A flower-head, $\times 4$.
 11. A flower-head, longitudinal section, $\times 4$.
 12. Corolla laid open, with four stamens removed, $\times 4$.
 13. Anthers, dorsal view, with three filaments cut off, $\times 4$.
 14. Fruit, $\times 2$.

PLATE 8.

Stiftia Connellii.

- Fig. 1. Branch, natural size.
 2. Bract from the receptacle, $\times 2$.
 3. Floret, with part of the pappus removed, $\times 2$.
 4. Stamens, $\times 2$.
 5. Apex of style, $\times 2$.

PLATE 9.

Bonyunia minor. (Figs. 1 to 5.)

- Fig. 1. Branch, with flowers and fruit, natural size.
 2. Branchlet of an inflorescence, $\times 4$.
 3. Part of a corolla laid open, $\times 4$.
 4. Longitudinal section of a flower, with the corolla removed, $\times 4$.
 5. Transverse section of an ovary, $\times 12$.

Lisianthus Quelchii. (Figs. 6 to 9.)

- Figs. 6 & 7. Portions of different plants, natural size.
 Fig. 8. Flower with two sepals and the corolla removed, natural size after boiling.
 9. Corolla laid open, natural size.

PLATE 10.

Utricularia Connellii. (Figs. 1 to 6.)

- Fig. 1. Three entire plants, natural size.
 2. Part of inflorescence, showing bract and bracteoles, $\times 5$.
 3. Flower, $\times 2$.
 4. Calyx, dorsal view, $\times 5$.
 5. Stamens, $\times 5$.
 6. Pistil, $\times 10$.

Utricularia concinna. (Figs. 7 to 11.)

Fig. 7. Three entire plants, natural size.

8. Inflorescence, $\times 3$.

9. Flower, viewed from above, with the dorsal sepal turned back, $\times 3$.

10. Stamens, $\times 10$.

11. Pistil, $\times 10$.

Utricularia Quelchii. (Figs. 12 to 16.)

Fig. 12. Two entire plants, natural size.

13. Bract, bracteoles, and calyx, natural size.

14. Corolla, natural size.

15. Stamens, $\times 5$.

16. Pistil, $\times 5$.

PLATE 11.

Utricularia roraimensis. (Figs. 1 to 4.)

Fig. 1. Group of plants, natural size.

2. Part of a raceme, with the corolla removed, $\times 5$.

3. Corolla, $\times 5$.

4. Stamen, $\times 10$.

Genlisea roraimensis. (Figs. 5 to 12.)

Fig. 5. Plant, natural size.

6. Fragment of the base of a stem, with leaf (+ +), false roots, and an imperfect utricle standing erect among the leaves, $\times 10$.

7. Perfect utricle, $\times 5$.

8. Imperfect utricle, transition form between the ordinary false root and the perfect utricle, $\times 5$.
This form has a very minute pore at the apex.

9. Imperfect utricle, erect form, $\times 5$.

10. Part of a raceme, with the corolla removed, $\times 5$.

11. Corolla, $\times 5$.

12. Stamen, $\times 5$.

PLATE 12.

Brocchinia reducta.

Fig. 1. Rosette of leaves, natural size.

2. Inflorescence, natural size.

3. Flower, with a sepal and two petals removed, $\times 5$.

4. Sepal, $\times 5$.

5. Petal, $\times 5$.

6. Style and stigmas, $\times 5$.

7. Transverse section of unripe capsule, seeds not shown, $\times 5$.

8. Seed, $\times 5$.

PLATE 13.

Connellia Augustæ.

- Fig. 1. Leaf, natural size.
2. Inflorescence in flower, natural size.
3. Inflorescence in fruit, natural size.
4. Sepal, natural size.
Petal, natural size.
6. Ovary, the other parts of the flower being removed, natural size.
7. Ovule, $\times 30$.
8. Ripe fruit and bracteole, natural size.
9. Seeds, $\times 10$.

Figs. 1, 3, 8, & 9 are from McConnell & Quelch's specimen, the rest are from Schomburgk's specimen.

PLATE 14.

Connellia Quelchii.

- Figs. 1 & 2. Plants natural, size.
Fig. 3. A large tuft of leaves, natural size.
4. A leaf, tomentose on the back, natural size.
5. Sepal, $\times 2$.
6. Petal, $\times 2$.
7. Ovary, $\times 2$.
8. Young fruit, $\times 2$.
9. Transverse section of an ovary, $\times 5$.

J.N. Fitch del. et lith. N.E. Brown anal.

W.E. Hemsley dir.

1-6 *ILEX APICIDENS*, N.E. Brown. 7-16 *CYRILLA BREVIFOLIA*, N.E. Brown.

J.N. Pritch del. et lith. N.E. Brown anal.

W.B. Hemsley dir.

MACAIREA ASPERA, N.E. Brown.

J.N.Fitch del.et.lith.NE.Brown anal.

W.B.Hemsley dir.

HENRIQUEZIA JENMANI, K. Schum.

J.N. Fitch del et lith. N.E. Brown anal.

W.B. Hemsley dir.

J.N.Fitch del. et lith. N.E.Brown anal.

W.B. Hemsley dir.

1-7 HETEROTHALAMUS DENSUS, N.E.Br. 8-14 QUELCHIA CONFERTA, N.E.Br.

J.N.Fitch del. et lith. N.E. Brown anal.

W.B.Hemsley dir.

STIFTIA CONNELLIⁱⁱ, N.E. Brown.

J.N. Fitch del. et lith. N.E. Brown anal.

W.B. Hemsley dir.

1-5. *BONYUNIA MINOR*. N.E. Brown. 6-9, *LISIANTHUS QUELCHII*. N.E. Brown.

J. N. Fitch del. et lith. N. E. Brown anal.

W. B. Hemsley dir.

J.N. Fitch del. et lith. N.E. Brown anal.

W.B. Hemsley dir.

BROCCHINIA REDUCTA, Baker.

J.N. Fitch del. et lith. N.E. Brown anal.

W.B. Hemsley dir.

CONNELLIA AUGUSTÆ, N.E. Brown.

J.N. Eitch del. et lith N.E. Brown anal.

W.B. Hemsley dir.

CONNELLIA QUELCHII, N.E.Br.