topsies, historical research into the past history of the disease, botanical and zoological research, particularly as regards the possible control of malaria by means of trees, plants, fishes, etc., and, finally, the ascertainment of the approximate economic loss; particularly in the case of plantation labor, in consequence of malaria, more or less in conformity with the methods adopted by the U. S. Public Health Service and the International Health Board.

In support of the plea and plan for malaria eradication Dr. Hoffman, in conclusion, presented a brief analysis of the results of a comprehensive statistical study of malaria frequency throughout the world. He directed attention to the remarkable decline in malaria in Southern states as well as in certain Northern states where formerly the disease prevailed to a not inconsiderable extent. In New Jersey, for illustration, the malaria death rate had been reduced from 37 per 100,000 in 1881 to only 0.3 during 1914; in Connecticut, from a maximum of 21.3 per 100,000 during 1881 to 1.0 during 1914; in Memphis the rate had been reduced from 227 per 100,000 in 1885 to 55 during 1914; in New Orleans, from 163 per 100,000 in 1885 to 9 during 1914; and in Jacksonville, Fla., from 278 per 100,000 in 1901 to 6 during 1914. These gratifying changes in the malaria death rate, North and South, he said, could not, however, be made to apply to the rural sections, particularly of the Southern states, where, as for illustration, in certain portions of southeastern Missouri, malaria continued to prevail to an alarming degree. He directed attention to the fact that the malaria death rate of Dunklin County, Missouri, was higher than the corresponding malaria death rate of well-known fever centres in the West Coast of Africa and the West Indies. Considering, however, the statistical evidence of the results of antimalaria measures at Panama, the Straits Settlements, the West Coast of Africa, and elsewhere, he said it was self-evident that similar results could be achieved throughout the Western hemisphere wherever malaria was known to prevail, but only through the active and hearty co-operation of federal, state and local government authorities, assisted by corporate and

other bodies interested in the prevention of disease and the prolongation of human life.

Such efforts, he felt sure, would not in the long run prove excessively expensive, but, much to the contrary, it was practically certain that even heavy local expenditures in behalf of malaria eradication would yield a return many times in excess of the original cost in consequence of an increase in land values and better opportunities for the more effective utilization of local natural resources. In any event, he said, the Nation was under the most solemn obligation as a matter of international agreement to assist in every reasonable way in the proposed effort at malaria eradication, and since such an effort was entirely practicable and of enormous public interest, it was to be hoped that the plan proposed by the National Committee on Malaria would be approved and receive the necessary financial and other support and co-operation on the part of the Federal government, the states and the localities directly concerned.

INFANTILE PARALYSIS

BY W. A. EVANS, M.D.,
Health Editor, Chicago Tribune,
Chicago, Ill.

Editor's Note.—Dr. Robert Wilson, Jr., President of the Southern Medical Association, in introducing Dr. Evans, said: "The lecturer whom we shall have the pleasure of hearing this evening, although coming to us from Chicago, is really of the South, having been born in Alabama, brought up in Mississippi, and educated in Louisiana; but for the last twenty-five years he has been living in Chicago. Formerly, he was Health Commissioner of Chicago, and is now Health Editor of the Chicago Tribune. A few days ago he was honored by being elected President of the American Public Health Association."

Last May, after a winter during which there had been an occasional case of infantile paralysis in New York City, there began to develop an unusually large number of cases in the western part of Brooklyn, in a little district just south of Greenwood Cemetery. These cases were overlooked. The majority of them were unrecorded and probably unrecognized. Some time about the middle of June a

*Address, Public Meeting, Southern Medical Association, Tenth Annual Meeting, Atlanta, Ga., Nov. 13-16, 1916.

sufficient number of cases of this disease had been reported to the New York Department of Health to excite its interest and curiosity, and an investigation was begun which disclosed the fact that there had been an unusually large number of cases of this disease in Brooklyn during the preceding month. But by the time the investigation had been concluded it was evident to those who were in charge that an epidemic existed, which proceeded through the summer with an increasing number of cases. The epidemic waged, reaching its maximum in the week of August 12, during which week there were 1,200 cases of this disease recognized and reported by the New York Department of Health. It was found that a focus would begin, would rapidly increase for about six weeks, would reach its maximum at the end of six weeks, whereupon there would ensue a period of gradual decline. But by reason of the fact that in this great community there were thousands of smaller communities, naturally the epidemic would travel from one focus to another, and from a new focus a number of cases would multiply until at the end of about six weeks' duration that focus would reach its maximum, whereupon it would begin to decline.

During the summer and fall months in that City there were approximately 12,000 cases of infantile paralysis. Within fifty miles of the lower end of Manhattan Island, in these four or five months, there were more than 15,000 cases of the disease. Probably in no previous year in the entire world's history had there been 15,000 cases of infantile paralysis. We have, therefore, just gone through the most extensive and the most malignant epidemic of infantile paralysis that the world has ever known. However, since August 12 the disease has been on the decline, a decline at first somewhat gradual and slow, but later more rapid until at the present time we may say that there is nothing more than the embers of that epidemic. It is a thing of the past. And if that be true, then why is it that we are taking up your time in considering this disease? If it is a matter of history, however modern the history may be, there is on the face of things no reason why you should give over an evening to learning about infantile paralysis, and particularly

why you should be appealed to, as I shall appeal to you during the course of the evening, to do your duty in connection with this disease. Now, why? This is the reason why.

It is the history of this disease that it has never begun violently in the same community two seasons in succession, but that in a succeeding season in which it has been violent in one part of the country, or one section, or one district, it is violent in some other part of the country. The theory is that when it invades a community in epidemic form, that practically every person in that community who has not a natural immunity acquires immunity either by an attack of the disease or by becoming infested with the micro-organisms of the disease, and infested in such a way that atypical illness will result, but nevertheless immunity is acquired. Those people who have either had the disease, or have otherwise acquired immunity in some degree, remain as carriers and can be the cause of the disease and later spread it to others.

You will understand that it is not a something created *de novo*. It is not a something that originates out of nothing. It is a living, active thing, and it must be kept alive by some agency capable of supporting life in it. There is no reason for believing,—in fact there is every reason for disbelieving,—the possibility that it can remain alive in water, in milk, in air, in dust, and in dirt. There is no reason for believing that it can winter over from season to season in an insect or in the body of any animal other than man. There is every reason for believing that the agencies that are harboring this hibernating organism in this winter season are human beings, some in New York City, some in Atlanta, some in San Francisco, and some in Chicago, harboring this organism and ready to feed it, when the time is propitious, back to those who are susceptible of having it.

The reason that it is worth your while considering infantile paralysis tonight is that the teaching of experience is that during the year 1917 this disease will be epidemic in some part of the United States, probably not in New York City, or in its immediate environment; probably in some part of the country distant from New York City, and possibly in At-

lanta. Quite certainly, men who are within the hearing of my voice, men who are in this hall tonight, before the year 1917 shall have ended will be brought in contact with epidemics of infantile paralysis. Men who are listening to me here this evening will be called upon to control an epidemic of this disease concerning which they know nothing, and other men will be called upon to diagnose and treat and to give expert attention to and care for cases of a disease with which they have had no experience. It behooves you, therefore, to learn of it before the oncoming of the next season.

Now, a word or two briefly as to infantile paralysis to the men who practice curative medicine. In all probability, the bacterium which is responsible for this disease is known. In 1912 or 1913, Flexner succeeded in growing the coccus of infantile paralysis and he succeeded in inoculating animals with this coccus and producing the disease in the inoculated animal. That information has been in hand for three years, and yet for practical purposes in this epidemic it has been of no service, and why? Because in the stage of the discovery, the information about it was only valuable in laboratories which were well equipped with apparatus and with men, and as a diagnostic procedure it has been of no help in this epidemic.

Within the last few weeks there have been additional discoveries which have come out of this epidemic which place the laboratory procedures at the service of the departments of health. The latest of these discoveries was announced in the *Journal of the American Medical Association* on Saturday of last week. Dr. Nuzum, of Chicago, collaborating with Dr. Herzog, has rediscovered the Flexner coccus of infantile paralysis. The additional discoveries that he has made, or they have made, have made Flexner's discovery available for practical diagnostic laboratory purposes, consisting in this: they have succeeded in finding the organism in the cerebrospinal fluid by prolonged centrifugalizing and staining of sediment. That, of course, means an immediate diagnostic procedure. By growing this organism from the cerebrospinal fluid in acetic broth or broth containing a little bit of kidney tissue, they have been able

to get a good growth within twenty-four to thirty-six hours, and that is a practical laboratory procedure. It is grown in the oven at 35° C. or at 95° F. It appears as a cloudiness in the bouillon, and at the end of another 12 to 24 hours this cloudy material settles to the sides and bottom of the test tube. The longer the organism grows, the larger it becomes. At its beginning, it is in a great measure an ultramicroscopic micro-organism or coccus which Flexner discovered. At the end of twenty-four hours a moderate proportion of the cocci are large enough to see with ordinary microscopic magnification. At the end of forty-eight hours a considerable proportion of them are visible with the ordinary immersion lens. A swab culture grown on blood agar slants will grow as a large coccus, easily seen under microscopic magnification.

This organism can be and has been injected into lambs, into rabbits, and into monkeys, producing flaccid paralysis, which is quite the same disease clinically as infantile paralysis in the human subject.

The organism secures entrance to the human body through the mouth or nose, and from this atrium travels to the spinal canal; some authorities say by the lymph routes, others say by the blood stream, when it reaches the cerebrospinal canal. It affects first, the spinal meninges in the region of the posterior roots, and that is an explanation of the pain and tenderness which in the usual case is a forerunner of the paralysis. It extends from there to the anterior part of the cord, and later involves in a destructive way the motor cells of the anterior horn. There is a definite sequence according to most authorities between the beginning of the disease in the posterior horn, or rather in the vicinity of the posterior nerve roots, and the later destructive development of the disease in the motor cells of the anterior horn. I shall tell you something about the percentage of certain clinical phenomena and certain principles of treatment that are dependent upon this lapse of time. So much for the pathology of the disease and something of its symptomatology.

I shall not occupy your time in detailing to you all the symptoms of the disease in that form where there is a violent onset,

with rapidly developing paralysis or convulsions, or other evidence of violent disease. I want to concentrate your attention upon the atypical form of infantile paralysis, a disease which begins much as does an ordinary attack of "summer complaint," a mild degree of fever, and some intestinal disturbance; and the keenest of diagnosticians, with nothing save these symptoms in mind, will fail to make the diagnosis more frequently than he will succeed. But if a man is alert, he will note presently the development of an unusual degree of tenderness and pain in joints and in limbs, a degree of pain that is not present in the ordinary distemper or ordinary febrile processes in the summer season.

I want to caution you to be on your guard for those cases of apparent distemper which is the cause of a mild febrile attack in summer that develop an unexpected and otherwise inexplicable amount of tenderness and pain in joints. In a certain percentage of cases, somewhere between the first and sixth day of the disease, paralysis ensues, starting gradually in a certain sense, and yet in another sense not gradually, for the disease at its very beginning, from careful analysis and careful search, presents more of paralysis than it does at a later time. In the more violent cases paralysis develops on the first day. In a considerable percentage of cases it does not develop until the sixth day, and other cases are in between. The importance of that lies in this: that before the development of the paralysis it is exceedingly difficult and oftentimes impossible for any practitioner of medicine, however alert he may be, to diagnose the disorder.

Our terror of this disease is not dependent upon the death rate, for the death rate is not high. In the Massachusetts epidemic there was a death rate of 8%, which means that eleven-twelfths of all people who develop infantile paralysis recovered more or less completely. The highest death rate from infantile paralysis in this country is the death rate in the recent epidemic in New York City. It seems that in that epidemic there was a death rate of 22%, and that means that of those who developed the disease in New York City, three out of every four recovered more or less completely. Then, our

terror of infantile paralysis is not due to its high death rate, for there are many diseases that have a higher mortality. It is due to the terrible results that follow in those who recover from the disease. It is due to the fact that we have people constantly before our eyes dragging their limbs around. It is due to this that this disease strikes so much of terror, and yet not all of the cases of infantile paralysis develop paralysis of those that recover. Somewhere between 40 and 50% recover without any permanent paralysis or wasting of the limbs. In the New York City epidemic 81% of those who died, died in the first week of the disease, 11% in the second week, 3% in the third week, and 3% after the third week.

A word or two as to the treatment. A few years ago it was said that we should treat infantile paralysis with urotropin. It has been demonstrated by the work of Hanslick and others that urotropin can not possibly be of any service in this connection. In the first place, it is not an antiseptic as urotropin; it does not become an antiseptic until it has been decomposed and formalin has been formed; and it does not produce formalin except in an acid medium. None of the tissues in which the coccus of infantile paralysis is present is acid, and therefore it is not antiseptic in this disease.

A while ago it was noticed that a considerable percentage of the people who died of infantile paralysis succumbed early, before there was time for destructive degeneration of the cells of the anterior horn. It was noticed, furthermore, that there was a primary period during which the symptoms were violent, and not infrequently during which death would ensue from embarrassment of respiration, and if the person could live through that period he would get better and remain better for a certain length of time until there began to appear that train of symptoms that resulted from the breaking down of the cells in the anterior horn. Out of that observation there grew the recommendation by Meltzer that adrenalin be injected into the cerebrospinal canal, the idea being that he would prevent in this way congestion of the meninges and pressure that was responsible for the embarrassment to respiration, an embarrassment which not infrequently

terminated in death; that if he could tide his patients over that period, there was a possibility of their withstanding destructive effects of the disease, and so he made the proposal that adrenalin be injected. A little later he believed there might be some specific action from adrenalin. His remedy has been more or less tried out during this epidemic. There is a report by Hoyne indicating that it may be of some value. The majority of observations, however, during this epidemic, so far as I have been able to learn, do not indicate that this treatment is of any particular value.

In 1910, Flexner demonstrated that it was possible to prevent the development of paralysis in monkeys having infantile paralysis by injecting into their spinal canals blood serum from persons or from monkeys who had recovered from the disease. A little later Nettar, in France, applied this observation practically in the treatment of human cases of infantile paralysis. The first case in which it was employed was a man with a severe attack of the disease, one in whom there seemed no possible outcome save death. However, that man recovered, and then he tried it in others. It has been employed in this epidemic in more than thirty-five cases, but it does not seem to be giving such good results in the United States as it was indicated that it gave in France. What is the treatment, and what is the philosophy of it?

There are developed in people who have infantile paralysis certain antibodies which begin to make their appearance within five days after the onset of the disease. They remain in the cerebrospinal fluid for an indefinite time after the disease has begun. They have been found in the cerebrospinal fluid forty years after the onset of the disease. They are found in the cerebrospinal fluid of any individual who has been made immune by a severe attack, with paralysis, or by a light attack without paralysis. What Nettar did was to employ the blood serum of people who had recovered from the disease. Some had but recently recovered, and one individual had recovered thirty-eight years before. He injects 10 c. c. of fluid a day into the cerebrospinal canal; he makes six injections, making a total of 60 c. c. of cerebrospinal fluid used. The

information which we have is that there have not been such good results from the use of this remedy in New York City as in the use of the remedy in Paris, so that, as the matter now stands, a physician would not be justified in treating the disease during its acute stage by putting in operation any treatment save what may be termed symptomatic treatment,—meeting symptoms as they arise.

And what is the acute stage? It is held that the acute stage runs from its onset to the time when tenderness disappears. The significance of that is this: manipulation, massage, and passive motion should not be begun until the acute stage has passed, but please note that for the recognition of the passing of the acute stage we are not dependent upon any technical procedure or any test which must be applied or can only be applied with great difficulty. We have the plainest of all guides, and that is this: does it hurt? If it hurts, or if it tires, or if it exhausts, you are beginning your passive motion and your exercise too early. If it does not hurt, then it is time to begin the manipulation of the limbs.

The second stage runs through the end of the period of tenderness to the end of two years. This period of convalescence is somewhat an arbitrary period. It is during this time that passive motion and massage are of most service. It is also during this period that it may be most harmful; for the muscles are easily tired or readily fatigued, and if they be stimulated beyond their point of endurance more harm than good results. And then comes the third stage, according to Lovett's classification of stages, the chronic stage, and it formerly was held that during this chronic stage there is no possibility of further improvement. In other words, whatever improvement takes place will be in the first two years, and after a period of two years has elapsed there is no further possibility of improvement. It has been proven that this conclusion is incorrect; that while improvement is difficult and slow, the possibility of improvement abides, and perhaps permanently abides. That is of interest to all practicing physicians as well as to health officers.

Infantile paralysis was first described, or was first recognized as a disease entity

in Austria, in 1840. The first report of an epidemic was in Lusanne, in 1841. Practically nothing was known of infantile paralysis until the beginning of this century. It is for all practical purposes a disease of the twentieth century. Since the year 1900 we have never been without infantile paralysis in the United States, and we can further say that no part of the United States has been free from some cases of infantile paralysis in the last sixteen years. However, during these sixteen years there has been some recompense for the wide distribution of the disease. There have been a multitude of careful and productive studies by Whipple, by Levaditti, by Peterson in Sweden, and by others. I would suggest that those of you who have access to the available records carefully study the report that was made to the Fifteenth International Congress of Hygiene and Demography on the epidemic of 1910-11 in Sweden, rendered into excellent English and found in most of the libraries in the United States.

In 1910, it was stated, and I believe correctly, that two-thirds of all the infantile paralysis in the world was in the United States, and I believe it is generally accepted at the present time that for the last sixteen years the United States has been a hot-bed of infantile paralysis and has been some recompense here, too. There have been a number of most excellent studies of its epidemiology made by the various departments of health and others in the United States. There have been excellent studies of the Buffalo epidemic, in 1912; of the three-year epidemic in Massachusetts, of the epidemic in Vermont, studies of the epidemic in New York City, in 1907 and 1910, and again this year, and a number of others that might be mentioned.

Probably the first valuable contribution to the epidemiology of infantile paralysis was that of Wickham. But we have made in this country two notable contributions, one by Caverly of Vermont, who demonstrated the fact that a great many of the people who are infected with this disease were either not sick at all or were mildly and most atypically sick. Caverly proved to the world that the largest factor in the spread of infantile paralysis was the person who was atypically sick,

the person who was undiagnosed or wrongly diagnosed, or held to be a well person, a carrier of the disease, if you please. And the second great contribution is that made by the laboratory workers, the first being the contribution of a health officer and a clinician—Caverly. The contribution of the laboratory workers was first the discovery by Flexner of the coccus of infantile paralysis, its artificial cultivation, the production of the disease experimentally, and the late additions to the bacteriology of the disease, and perhaps the addition recently reported by Rosenow and his group, and more recently the additions made by Nuzum and Herzog, to which reference has already been made.

It has been demonstrated that infantile paralysis remains in the body of certain persons who have been infected with the disease for a period of five months; of monkeys that have been infected for a period of six months, and there is not only the possibility but the probability that it can remain in the noses of people who are acting as carriers for still longer periods. However, it has been proven that the period of infectivity in the average case does not last longer than six weeks, and therefore a six weeks quarantine period is long enough, for it fulfills all indications.

Every effort to control infantile paralysis is based upon the theory that it is spread by contact and carriers; and yet it is with exceeding difficulty that we recognize the facts in relation to this theory. For instance, in New York City in this epidemic, in the report made about a month ago, it was stated that in 97% of the cases there was but one case to the family. In practically 3% of the cases there were two cases in a family; in three-tenths of 1% there were three cases in one family. In a very few cases—too few to figure in terms of percentage—there were four or five cases to the family. This conclusion has been criticised by the statement that the investigation did not determine how many people there were in those families who were capable of having the disease; yet there is the observation made in the Buffalo epidemic that answers that criticism.

In the Buffalo epidemic 263 cases were studied, occurring in 263 houses. One

hundred and forty-two of those cases were in families where more than one family was exposed to the disease. There was a total of 350 families who were exposed to these 142 cases. How many of them developed the trouble? There were four contact cases in three families. In one family there were two contact cases; in two families in each there was one contact case. How different that is, as you well know, from, let us say, yellow fever, in which White and Carter demonstrated that the disease would travel from house to house; that it was exceedingly loath to jump across the street or to invade the houses on the yonder side of any open space. How different that is, as you all know, from scarlet fever, in which you expect cases to develop in the same school room with other cases; in which you know your method of search for the origin of the disease is to go to the children that have been in contact with the sick children and to search there for peeling of the limbs or other evidence of scarlet fever. Infantile paralysis will show itself here, and the next case will be in an adjoining county or some far-away community.

Please note this observation: In Illinois during the past summer, in a study of 335 cases, it was found that they occurred in 146 communities. Approximately half of the cases were in one community, Chicago. But note this: 99 cases occurred in communities in which there was but one case, and 31 in communities in which there were 12 cases. What is the explanation? One case of infantile paralysis developed in the country neighborhood and the only case in the entire community, no cases developing from it, and no possibility of demonstrating the relationship between that case and any other.

Again, the age, the distribution of infantile paralysis is not indicative of contact infection. Nine cases in the Illinois investigation were under six months of age, and 22 cases between 6 and 12 months of age. In the New York epidemic 10% of the cases were under one year of age; 44% of the cases were between 2 and 4 years of age, and 76% of the cases were under 6 years of age, or 76% of the cases were under school age. Furthermore, the disease tends to end with the opening

of schools, so that there is much reason for believing that it is not a disease that is due to contact infection.

Another theory is that it is due to infection from insects. Part of the Massachusetts study was made by Rosenau. They noticed in Massachusetts that it was a rural disease; that it spread along routes that horses travel rather than along routes of railroads. They concluded it must be transmitted by insects that had to do with horses, and they got some proof that it was spread by the bite of stable flies. They were able to produce the disease by letting stable flies infect monkeys and then bite susceptible monkeys. That observation was verified by Anderson, but since that time no observer has been able to sustain the conclusion, and the tendency at the present time is to believe that it is not spread by insects.

The recently-announced theory of Richardson, of Boston, that it is spread by flea bites; that fleas in turn spread it by rats, has not met any particular approval. There is much reason for believing that the disease is carried into the intestinal tract and is spread by food substances. It is a disease of the summer time. Its season is the fly season. It has the same seasonal prevalence as typhoid fever. Furthermore, it has been found occasionally that there were ulcers in the intestinal tract, and not infrequently enlarged lymph nodes in the intestines have been demonstrated. Those are the reasons why many men believe that our efforts to control this disease should be exerted along food lines rather than along contact lines in the years to come.

Now, there is a brighter side of the picture, and it is this: in the first place, I want to call your attention to the fact that in the year 1916 they have had in the City of New York the lowest baby death rate they have ever had in the history of that City. There have been more than 900 fewer deaths of babies in New York City during the year 1916 than in any previous year. In the three months when this terrible, devastating epidemic was prevailing in that community, there were practically 400 fewer deaths in babies from all causes than during the year 1915. Nineteen hundred and fifteen was their record year. In 1915 New York City had one of the lowest death rates of

any of the baby death rates of any large city on the face of the globe. What is the explanation? The explanation is that the effort which has been directed toward the control of infantile paralysis in that community during the year 1916 is carried to other diseases, and in spite of the fact that nearly 3,000 people have died from infantile paralysis, and the majority of them were children, in New York City during the year 1916 there has been a decrease in the number of deaths from all causes, and why? Because the Health Department has been more active in bringing about sanitation, in producing cleanliness, in controlling flies, in caring for the milk supply, than it has ever been before, partly because it has more men and more money than it has ever had, but in greater part because it has had more co-operation and has had more help from the people themselves than it ever had before, and that is one bright spot in that very dark picture.

Then, there is another thing they have demonstrated there: that while they have not been able to control the disease in its entirety until cold weather came on, it has been possible to control in part the epidemic focus in about six weeks, and that means that the same measures applied ought to control an epidemic in an isolated community, in a country community, in the same length of time.

Hitherto, we have held there was no help; that practically it had to run until frost came. They have demonstrated with their imperfect machinery that it is possible to control the disease in about six weeks; and now that they are at the end of this disease, we are on the verge of another season. We are here in this meeting laying our plans to the end that there may be no epidemic of infantile paralysis in 1917; and what measures shall be undertaken? In the first place, the disease must be reported; and in the second place, it must be diagnosed. Wherever possible, every case must be hospitalized; quarantine all cases and all carriers, all contacts, all those who have been in contact with the cases as well as those who have been sick, to be inaugurated and to be maintained for about six weeks. There must be mediary disinfection. I do not believe that anything is gained from terminal disinfection. What is the distinction?

The distinction is this: there is far greater need for mediary than for terminal disinfection. By mediary disinfection we mean the disinfection of discharges during the course of the illness as distinguished from that disinfection which leaves everything until the illness has terminated and then assumes to wipe the slate clean by liberating some formalin or burning some sulphur.

Were I a health officer and charged with the responsibility of controlling this disease, did I feel that behind me there stood women and men depending upon me to buttress them against this specter that stalks in our land, I would place my emphasis upon the prohibition of the use of uncooked food. That is my judgment. Perhaps it is a guess, incapable of demonstration, but I think there lies a method of control that offers us much of hope, and if the responsibility for protecting the people were with me, I would do everything within my power to see to it that those people protect not only all food that was uncooked, but raw milk, raw fruits, or raw vegetables, or any other form of raw food, and into this injunction I would throw every bit of kinetic drive, every bit of determination, every bit of personal power and persuasion that lay within my power to make my people unanimously put into operation that as a method of preventing the disease.

And now, finally, what can the different groups who are represented here tonight do as their contribution toward the control of infantile paralysis? I believe that we are confronted by a duty that is essentially a duty of the American people. We have done many things in the way of health conservation and disease prevention of which we may well be proud. We have driven back from our shores cholera. I dare say there is not within the hearing of my voice one person who has personal recollection of the fact that not more than two generations ago our land was devastated by cholera, and we have rolled it back, have perpetually banished it from our shores. And if there be none of those here who remember the menace of cholera, there are certainly those present who recall how our people were constantly in dread of the menace of yellow fever. Many of you know now of the security with which we go about in our daily work

in summer time by reason of the fact that scientists and health officers have seen their duty and done it.

There are other accomplishments of which we may well be proud. Our army on the Rio Grande for five years has been absolutely free from typhoid fever by reason of the efficiency of men whose duty it is to guard public health. I learned that the First Illinois Cavalry that recently returned from the Rio Grande report that in their tenure of service of practically four months there has not developed in that command a case of venereal disease. We are proud of our accomplishments. We are proud of the fact that we went to Cuba and drove from that land the relic of the old world monarchy. We are prouder still of the fact that we went to Cuba and demonstrated that American intelligence and American efficiency could make that pest hole of yellow fever and malaria a place fit for the habitation of men. We are proud of that fact. We arrogate to ourselves a certain superiority to other people because we went to Panama, where a great nation had attempted to do a great task, and had failed,—failed because they had not planned along health lines as completely and as scientifically as they had planned along the engineering line. We went there and made it one of the healthiest spots of the world and made it possible for the black man from Tahiti and San Domingo to work in comfort and efficiency along the Chagres River. Why? Because of the capacity of American scientists, because of the efficiency of American health officers. We have arrogated to ourselves a certain possession of capacity in the control of health problems, and when certain people were struggling in the days of despair to put down typhoid fever, it was to America that the cry came to save us from the enemy that oppresses us. We stand, therefore, in the eyes of the world as a nation of accomplishments in disease prevention. Therefore, we have a duty in regard to infantile paralysis because by all the world we are regarded as the pest hole of infantile paralysis.

What are the responsibilities of the different members of our groups in relation to this disease? As practitioners of curative medicine, what can you do? You can control the key to the situation. We will

control the disease if you are capable of recognizing it early and acting upon correct diagnoses. The difficulty lies in this: the majority of you have never had an opportunity to see cases of infantile paralysis. I want to call attention to the fact that in the State of Vermont in 1912, subsequent to their epidemic, the State Board of Health was given the sum of \$10,000 to spend in evolving methods for controlling this disease, and how did they spend it? They spent part of it to bring laboratory workers into the State to attempt to find a method of diagnosing the disease and producing a serum for its cure. They spent part of it in bringing clinicians into the State to locate, to hunt out and ferret out the cause or causes of infantile paralysis. Then they spent another part of the money for what? To see that every man practicing medicine in the State of Vermont had an opportunity to see some cases of infantile paralysis, remembering this, that nobody is quite so much interested in competent, trained, experienced doctors in the midst of the community as the people themselves; that it is worth paying for to give these men free of expense to themselves an opportunity to learn practically about infantile paralysis. And this is the duty, as I see it, of the men who are practicing curative medicine—to avail themselves of every opportunity to learn the disease, so that when they come in contact with such cases they will be able to ferret them out, and having ferreted them out, to bring them to the attention of health authorities.

Now, what is the function of departments of health, as I see it? They have a duty outside of their duty to care for the cases as they develop. I believe it is their duty to plan to the end that better principles, better opportunities for diagnosis of infantile paralysis may be available even than those in the summer of 1916. I believe that we have come to the point where it is feasible, as a practical health department procedure, to put laboratories for the practical, clinical diagnosis of infantile paralysis at the service of all doctors of a state and in the various communities, and that I believe to be a part of their duty.

In spite of our discoveries, there are things that yet remain undiscovered. We

know comparatively little of the methods by which infantile paralysis is spread. I believe they have a duty in relation to this disease in trying to ferret out the relation between cases of infantile paralysis, the laws of spread of one case to another.

Now what is the duty of the laboratories? I trust that every one here understands that there is a relationship between the man who works in cloister, who works in a laboratory, who works isolated from the things of every day, and the work of the market place. What is the duty of the laboratory in this great co-operative work? I believe we have developed methods for practical every-day diagnosis, but we are far from having developed methods of cure that are satisfactory. The methods of diagnosis that we now have can be materially improved, but the best method we have of curing cases by the injection of serum taken from people who have recovered from the disease is far from being a practical procedure.

In New York City they advertised in the midst of their epidemic for people who had had the disease, who were willing to contribute blood serum for the cure of other cases. I learned that people came forward in great numbers, and some of those who were cured donated to the Department of Health considerable sums of money to be given to those people who were in need of it who had contributed blood serum. I heard they have had great difficulty in disposing of that money, that those who donated their blood serum insisted that they did it from the best of motives, patriotic motives, humanitarian motives, motives of fellowship and right, and that they refused to be compensated for the service they had rendered. Yet that does not alter the main fact that any therapeutic procedure that depends for the source of supply of a remedy upon the voluntary donation of people will not permanently work as a therapeutic procedure. Laboratories, therefore, are under the necessity of perhaps perfecting this serum and certainly of discovering some method by which it or some other similar substance can be recovered from the bodies of large animals, for instance, horses, cows, sheep, etc. The laboratories have a duty to discharge in connection with this disease. I would not

have the people understand that they are free of responsibility, for they are not free from responsibilities toward this situation. The duties can not be carried and should not be carried alone by health departments or by practicing physicians without departments, or by laboratories, practicing physicians and health departments, but the people themselves have a duty to discharge in relation to this disease, and what is that duty? It has been said that the greatest need of a democracy is not an intelligent leadership, but is rather an intelligent fellowship. There is the need that the people should understand that if their representatives are to protect them against preventable disease, they are charged with the duty of supporting these health departments not only with money, with material means, with taxation given them by governmental bodies, and other monies furnished by philanthropists, but they must support them by intelligent public sentiment; they must stand behind them and uphold them; that they must furnish the bulwark and the solid foundation upon which health departments can build and without which health departments will not build.

THE WASSERMANN TEST IN PUBLIC HEALTH WORK*

BY F. A. COWARD, M.D.,
Columbia, S. C.

The writer wishes to present the following remarks upon his experience with, and trials of, the Wassermann test during the past six and one-half months, as used in the Laboratory of the South Carolina State Board of Health. The test was performed free of charge on specimens sent in from any source, the only requirement being that certain information as to the social status of patients be given. It is our opinion that the value of the Wassermann test no longer requires discussion. In public health work it has the same standing as other well-recognized tests such as those for tuberculosis, typhoid fever, hookworm, malaria, etc.

As Dr. Martin, our Chairman, stated in

*Read in Section on Public Health, Southern Medical Association, Tenth Annual Meeting, Atlanta, Ga., Nov. 13-16, 1916.