The point that he had wished to emphasize was the existence of a group of cases developing under intensive military training, with pathological pupillary response to light, coarse tremors of the face, tongue and hands and a slight dysarthria closely simulating that of early paresis. He had tried to make clear that the mental and general cerebral symptoms were not characteristic of paresis, but in conjunction with the somatic symptoms were suggestive and corroborative. Furthermore, the disappearance of all symptoms, general cerebral, mental and somatic, after a period of rest, and the normal serological findings, left no other interpretation than that of fatigue or exhaustion of the central nervous system.

Dr. Hunt again stated what he had previously emphasized, that he had never encountered the condition reported in civil life, and any importance or significance which attached to this fatigue syndrome was, he thought, limited to the exigencies of military life. He thought it not unlikely that cases of this description which showed thus early a tendency to exhaustibility of the sensory nervous system might be the type of case which later, under greater provocation of stress and strain, might pass into the more severe neuroses or neuropsychoses of war.

These men were not recommended for discharge for the reason that they were quite willing to resign voluntarily from the camp, realizing from their experience that they were unfitted for the mental and physical strain of the life. Under conditions where the candidate was unwilling to resign and yet presented such a striking picture of fatigue, the question of the recommendation for discharge would have to be settled on the individual merits of each case.

CHICAGO NEUROLOGICAL SOCIETY

NOVEMBER 15, 1917

The President, DR. ARTHUR W. ROGERS, in the Chair

A CASE OF "QUADRANTIC BLINDNESS"

By Brown Pusey, M.D.

The man consulted Dr. Pusey complaining of defective vision. He was able to read ordinary print slowly. Examination of the fields of vision showed blindness of each eye in the lower right quadrant. The fields otherwise were normal. The refracting media and fundi were normal. The case was shown because of the very great rarity of the condition. Dr. Pollock described the case further.

History.—(October 29, 1917.) The patient stated that he was apparently well until a week ago, when he complained of severe pain in the head. He went to work, but returned at four o'clock, his headache having been worse. He rode a block past his house and then walked back. From then on he was unable to see distinctly. He vomited once that afternoon and was unable to work the next day because of severe frontal headache. His memory was so affected that he had forgotten where he lived.

Previous history negative. Denied lues; Wassermann reaction negative. Did not drink, smoked moderately.

Physical Examination.—A well-nourished male, aged fifty years, 5½ feet tall. Chest negative; abdomen and contents negative; extremities and genitalia negative. No gross loss of power in any of the extremities; no tremor. Motor and cranial nerves all normal. Heat and cold, touch and

pain sense normal; no ataxia, no dizziness. Deep reflexes equal and normal; knee jerks variable, wrist jerks equal, triceps equal, normal. Jaw jerk normal. Plantar on the right frank, left less so. Cremasteric slightly sluggish on left, as was the abdominal. Pupils equal and reacted to light and in accommodation. Ocular movements normal except that perhaps convergence was a little bit poor. Sensory and cranial nerves normal, aside from the optic nerves. In speaking there was a slight tendency to misunderstand spoken words. He complained of forgetting numbers; could not remember the name of his first wife or of his present wife. He could not tell where Marshall Field's was located; said the City Hall was where the LaSalle Hotel is. He could not remember 2468 after two other numbers had been given him. Although he stated that he was a school teacher and was accustomed to doing rapid calculation, he could not do simple problems. The patient's blood pressure systolic was 140. He complained of headache, which he said was severe, but he was up and about and when his attention was distracted from his head he apparently had no pain.

He was sent to the dispensary clinic, where he had to go down an incline from the hospital; when he came out of the clinic he thought he had to go down stairs and lost his way in that manner on a number of occasions. At first he could not recognize any writing, but on a number of other occasions following Dr. Pusey's examination he could read his own name and on other occasions could read rather small type. He was given a deck of cards and told to deal himself a full house; he looked through the deck several times, passing three queens, muttered to himself "eight" and finally with great difficulty was able to pick out three eights. There was some defect in writing; he could write to dictation those things with which he was familiar—his name, address, Chicago, but when asked to write President Wilson, started with p-r-e-s-, and had great difficulty in getting further. There was an agraphia which was present not only on dictation but also on volitional writing. Relative to the agraphia, it might be interesting to recall the fact that a similar case was described by Casamajor in 1907 in the *JOURNAL OF NERVOUS AND MENTAL DISEASE*, except that that was a case of homonymous hemianopia with agraphia in which the question arose as to the localization of the lesion in the cases with an agraphia. Attention was called to Dejerine's dictum, that those lesions which were in the angular gyrus were associated with agraphia, whereas those which were subcortical were not affected with agraphia and in these a pure alexia was present.

Dr. Hugh T. Patrick said he had seen several of these cases of quadrantal blindness, but a clean-cut quadrant was very rare and very interesting in relation to the structure of the occipital cortex. Most of the peculiar fields were due to occipital lesions. Just at present these were even more interesting than they had been because a great many had been observed since the war began. Marie and Foix had reported a number of cases of gunshot or projectile wounds of the occipital lobes and had found some most extraordinary fields. The thing was of interest because of the question of cortical representation for color aside from form and a most interesting feature was the association of other symptoms with the peculiar visual defects. These related first to those symptoms connected with the various aphasias, alexias, agraphias and so on. In his experience the cases if seen early were associated with some form or degree of aphasia, and he believed the difficulty which the patient presented had in reading was due to the man's failure to recognize what he really saw. When Dr. Patrick was told that the patient had poor central vision, he said he must have something else besides an occipital lesion and it turned out, as in so many of the cases, that the trouble lay not in lack of visual acuity, but in lack of recognition, just as in ordinary word blindness.

The gentleman with quadrantic blindness for color only also had difficulty with his speech as well as with vision, which Dr. Patrick believed was due to a difficulty of recognizing the significance of what he saw. This applied not only to written characters but also to objects, which brought up another symptom which he thought was nearly related, and that was the point Dr. Pollock spoke of—failure to recognize surroundings. The patient did not orient himself properly, his recognition of places was defective.

Dr. Patrick cited a case seen several years ago in which the patient had an irregular, incomplete horizontal homonymous hemianopia. That man would get lost in his own house where he had lived for many years and could not go many yards beyond the front gate without losing himself. That would seem to have no relation at all to aphasia, but he believed it was closely related to visual aphasia in that it was an object blindness—a failure to recognize and appreciate what was seen, just as a person with word blindness sees perfectly but fails to recognize the printed characters he sees.

In another case seen recently there was an indication of the same sort of thing, which had been misinterpreted as mental confusion. A woman with a homonymous hemianopia was supposed to be mentally off because she acted so peculiarly. The trouble was that she did not see things on the right-hand side and at the table she would put out her hand without turning her head to look and fail to get what she reached for; second, she did not recognize all that was said to her; third, she did not recognize what she read and, fourth, she did not express herself fluently. All these collateral symptoms were due to the fact that the lesion was neither strictly cortical nor strictly occipital but extended forward into the speech centers or the fibers from them.

PSEUDOSCLEROSIS

By Lewis J. Pollock, M.D.

Dr. Pollock read a paper on this subject and presented a patient with this disease.

After reviewing the literature on the subject the speaker stated that in many respects pseudosclerosis was closely related to Parkinson's disease, and particularly to the so-called juvenile form. Many of the cases reported as juvenile paralysis agitans had really been examples of other diseases and the pure clinical cases had been very few in number. Many of the so-called juvenile paralysis agitans were certainly cases which would fit in the description of pseudosclerosis.

In the case shown, which had been diagnosed as juvenile paralysis agitans on several occasions, it was the tremor and rigidity which led to the diagnosis. The marked asymmetry of the involvement, and lack of conformation to type of the muscle rigidity, the extent of the tremor all spoke against paralysis agitans, as the chronic course, absence of emaciation and dysphagia, spoke against the inclusion of the case with a categorically defined Wilson's disease.

If juvenile paralysis agitans was to be considered as a clinical entity, conforming in its symptomatology to the adult type, it had better be sharply differentiated from the cases which differed in any respect from the recognized type. These cases, as the one shown, must be included in a group of diseases of the lenticular nucleus either complicated by diffuse changes elsewhere or affecting one or another area of the lenticular nucleus. Some of the cases might further be grouped, from the similarity of the cause or general pathology, as in the cases of pseudosclerosis and Wilson's disease showing the concomitance, the lenticular nucleus degeneration and liver disease.