The frequency of the interruption of pregnancy in epidemic pneumonia is very high.

Abortion occurs with relative ease and lack of pain.

In the majority of cases, death follows within twenty-four hours after emptying of the uterus.

The percentage of recoveries after interruption of pregnancy is small.

THE PROPHYLACTIC VALUE OF LEARY'S VACCINE

HARRY LEE BARNES, M.D.

WALLUM LAKE, R. I.

As an institution offers unusual advantages for the study of an epidemic and the measures for its control, a record of our experience with influenza vaccine at the state sanatorium may not be without value. The institution had about 225 patients and ninety-four employees. During the last two weeks of September, 1918, four employees and five patients contracted influenza. These patients were all in one ward of the hospital and were apparently infected from a patient who visited Woonsocket. They were isolated in rooms and a cubicle of the ward, and the disease did not spread.

INCIDENCE OF INFLUENZA AMONG VACCINATED AND
UNVACCINATED

Wards	Vaccinated			Unvaccinated		
	Number of Persons	Cases of Influenza Num- ber	Per Cent.	Number of Persons	Cases of Influenza Num- ber	Per Cent.
East.....	32	11	34	26	12	46
West.....	31	6	19	10	3	30
Hospital 1.....	16	0	0	24	1	4
Hospital 2.....	14	1	7	24	0	0
Children.....	32	0	0	1	0	0
Employees.....	27	7	26	28	7	25
Total.....	152	25	16	113	23	20

The four employees were isolated in their rooms. September 29, all patients and employees, with few exceptions, were forbidden to receive visitors or leave the institution. We were free from influenza, which was at the height of the epidemic in the surrounding country, until October 10, when a woman patient admitted on October 9 developed the disease. October 14, a patient who slept near this patient developed influenza. October 22, the first case developed in the men's ward of the sanatorium, and from there the disease spread to the employees, until there had been a total of eighty-two cases, or 25 per cent. of the total population. All of the 114 young adults, including some expatient employees sleeping in the east and west wards, were probably well exposed to the disease and furnish the best opportunity to estimate its morbidity. Counting all cases that occurred before and after vaccination, 45, or 40 per cent., developed influenza. October 22 and the few days succeeding, influenza vaccine furnished through the courtesy of Dr. Timothy Leary of the Tufts Medical School was given to 172 employees and patients. The doses were 400, 800 and 1,200 million at twenty-four hour intervals.

In computing the incidence of the disease for comparison between vaccinated and unvaccinated, no cases were counted that developed before the vaccine was given, and no patient has been counted as vaccinated unless he received the three doses. Before the arrival

of the vaccine, eight cases that terminated fatally had developed.

In estimating the value of the vaccine, it would seem clear that the children should be deducted from the total because they were quarantined and so far as known were not exposed to the disease. If the children are deducted, the influenza incidence was 20 per cent. among both vaccinated and unvaccinated. It would also appear preferable to deduct the two hospital wards which had only one case each which were so promptly isolated that the exposure probably was insufficient to test the vaccine. If these wards are deducted, ninety vaccinated persons furnished twenty-four cases, or an incidence of 26 per cent., while sixty-four unvaccinated persons furnished twenty-two cases, or an incidence of 34 per cent. In twenty-five influenza cases following vaccination, four patients, or 16 per cent., died. In fifty-seven influenza cases among the unvaccinated, the mortality was nine, or 15.8 per cent.

CONCLUSION

The morbidity was only slightly lower among the vaccinated, and the mortality among those who developed influenza was practically the same whether vaccinated or unvaccinated.

Military Medicine and Surgery

THE PANDEMIC IN THE ARMY CAMPS *

GEORGE A SOPER, PH.D., M.D.

Major, S. C., U. S. Army

WASHINGTON, D. C.

The pandemic in the Army camps was a part of the great outburst that swept over the United States in September and October, 1918. This, in turn, was a manifestation of the still larger visitation that was to cover most, if not all, of the world.

As soon as the pandemic appeared, each camp and other station of troops was required to keep the Surgeon-General informed of the appearance and progress of the disease. Daily and weekly telegraphic reports were sent in giving the number of new cases of influenza and pneumonia and other details. Most of the statistical data presented here are taken from these reports.

Were it not for the pulmonary complication which has accompanied it, few would doubt that the disease was influenza. Its epidemic character, its clinical aspects and its bacteriology, so far as they have been studied, all point to that disease. It will consequently be called influenza here.

As to the lung complication, that is a similar matter. Objection has been made to the use of the term "pneumonia" to describe it. As yet no other official designation has been given it. It may or may not be a part of the original disease. Pending further investigation, and since it must be given some name in this report, it will be called pneumonia here.

Had it not been for the pneumonia, the pandemic would not have attracted much attention. Epidemics of so-called "colds" and other respiratory diseases occur every fall, and if the typical influenza of today does not bear a close resemblance to these

* From the Division of Infectious Diseases and Laboratories, Medical Department, U. S. Army.

familiar infections, there have been many cases in which the difference was not discernible. It took the pneumonia with its high mortality to command attention. The pneumonia places this outbreak among the most striking pandemics of modern times.

Excluding the pneumonia, the influenza of September and October seems to be identical with the influenza that visited the Army training camps in the United States last March and April. That it attracted little notice then was due to its mild character.¹ The influenza which until recently has been reported from Europe appears to occupy a position between these two so far as severity is concerned.

Apparently there have been three visitations, or waves, each succeeding one being more severe, more widespread and more fatal than its predecessors. This fact may or may not have significance for the future. It is possible that there will be another wave. There may be one or more. It is equally possible that there will be none. There is no firm ground on which to base a prediction one way or the other. Considering the gravity of the matter, it would be unwise to assume that we are entirely done with the disease.

During the period from Sept. 12 to Nov. 1, 1918, inclusive, there were 306,719 cases of influenza reported among the troops in America. During the same time there were 48,079 cases of pneumonia and 19,429 deaths. The total strength of troops may be taken as one and a half million. Therefore it is approximately correct to say that about one in every five had influenza, that of these about one in six developed pneumonia, and that of the pneumonia patients about two out of five died.

The country as a whole suffered severely but not so greatly in proportion to the population as did the camps. Up to and including October 26, a total of 86,045 deaths from influenza and pneumonia had been reported to the United States Public Health Service. This figure is incomplete, since it includes no report from some states and only partial reports from others; in many states deaths for cities only were included.

In the forty-five principal cities for which statistics are available, 3.24 per thousand are reported to have died. There were 127,654 deaths from all causes. Had there been no epidemic of influenza-pneumonia, it is estimated that there would have been 47,967 deaths. The difference therefore represents the deaths due to the pandemic: 79,687. If this rate prevailed throughout the continental United States, there would have been 341,021 deaths. To this must be added the 19,429 in the camps, making a total of 360,450 deaths. These figures are not estimates—they are not even good guesses; but they serve to show the scale on which the pandemic operated.

I. THE STATISTICAL FACTS

THE OUTBREAK

The first report that a serious epidemic existed in any camp came from Devens, at Ayer, Mass. September 16, the camp sanitary inspector, reporting through the camp surgeon to the Surgeon-General of the Army, announced that an epidemic of so-called

"Spanish influenza" had broken out at Devens as a part of a general epidemic that had attacked Boston and the neighboring states and towns some weeks before.

The Devens epidemic is supposed to have commenced, Sept. 7, 1918, in D Company, Forty-Second Infantry. On that date a case of supposed meningitis was sent to the hospital from this company; on the following day twelve cases were sent for observation. These proved to be influenza. By the 16th, thirty-seven cases had gone from the same company. One death from pneumonia had occurred. Almost simultaneously, other cases appeared in other organizations. By September 12 the total number of patients that had been admitted was 599. The disease spread rapidly, in spite of the measures taken to check it. September 20, the epidemic reached its maximum intensity. On that day, 1,543 new cases were reported as having been admitted to sick report. After reaching this high point the number of new cases rapidly became less, so that by the end of the month there were less than 100 new cases reported per day.

Meanwhile, pneumonia had become a frequent and fatal accompaniment of the influenza. Fifty new cases were reported, September 19, less than two weeks after the influenza had been discovered. The number rapidly increased: September 24, there were 342 new cases. The number each day remained at about 200 for four days; then there was a decrease, until, before the end of the month, forty per day had been reached. By October 6, the incidence had fallen to ten or so a day. Through the rest of the month the number varied between three and seventeen per day. There was no return to large numbers.

TABLE 1.—THE RISE AND FALL OF THE EPIDEMIC AT CAMP DEVENS

	Duration, Days	Cases Influenza	Cases Pneumonia	Deaths
Rise (Sept. 12-19).....	8	3,283	43	16
Peak (Sept. 20-21).....	2	2,722	205	43
Rapid decline (Sept. 22-29)...	8	3,141	1,495	298
Slow decline (Sept. 30-Oct. 18)	19	571	571	310

The Devens outbreak, so far as may be understood from the records at hand, could be divided into four parts, as shown in Table 1. It will be observed that the rise of the epidemic covered a period of about eight days; the peak, two days; the rapid decline, eight days, and the slow decline, nineteen days. Half of the deaths and nearly three quarters of the pneumonia occurred within less than three weeks of the outbreak.

CHARACTERISTICS OF THE TYPICAL EPIDEMIC

The characteristics of the Devens epidemic have been described here because they represent what has occurred at many camps. The earliest cases have often escaped identification. They have sometimes been taken for cases of food poisoning, meningitis or one of the common exanthems. The disease, which is epidemic, bears little resemblance to the coryzas and other respiratory affections to which the term "influenza" has been generally applied for the last twenty years. The so-called influenzas of ordinary times seem to have no relation to the present disease. Yet with a knowledge that influenza was in the vicinity, it is probable that cases of other respiratory infections have often been reported as that disease.

1. A report by the present writer, on an influenza epidemic in the Oglethorpe camps in March, 1918, may be found in abstract in the *Journal of Laboratory and Clinical Medicine*, 1918, 3, 46-49, and the *Military Surgeon*, 1918, 43, 398-402.

The leading symptoms in the typical cases are severe headache, chills or chilliness, pains in the back and legs, temperature sometimes as high as 104, great prostration, and drowsiness. Occasionally there are nervous symptoms; sometimes, but not always, the eyes and the air passages of the nose and throat are affected; there may be gastro-intestinal disturbances. The onset is sudden.

The fact that an epidemic existed in a camp has been generally recognized when the number of new cases has amounted to 100 or more per day. The incidence then increases rapidly. Sometimes the records show a great number of cases at the start, and there are marked fluctuations in the daily incidence as the epidemic continues. Striking irregularities do not represent the way in which the disease occurs, but are to be accounted for by the stress and difficulty with which the returns are collected. The greatest number of new cases reported for any day has often considerably exceeded 1,500 in a camp. At Devens the maximum was 1,543, at Grant, 1,810 and at Custer, 1,764.

The principal camp outbreaks have been charted as a part of the routine work of watching the progress of the pandemic. In making these charts it has been found convenient to mark time along the horizontal scale, giving one fifth of an inch to a day. The daily incidence of cases has been measured vertically, one inch representing 100 cases. So charted there is much similarity among the curves and especially among the members of certain groups of them. The explosive character of the infection is plainly indicated in most, as are the short duration, large incidence and rapid subsidence. The curves are never smooth, but are jagged, with steep inclines between the peaks and valleys. Only in exceptional camps has there been a recurring wave of noteworthy proportions during the course of the epidemic.

In another series of charts, the daily incidence has been plotted as rates, thus making allowance for the changes in strength that have occurred. In a third series the jagged irregularities have been eliminated, and smooth curves trace the rise and fall of the influenza and pneumonia. This has been done for all the camps. These pictures show as clearly as the data permit how the disease has come and gone. The charts furnish an invaluable means of analyzing and comparing the several camp epidemics. They have made plain many instructive facts and have pointed the way to many useful conclusions. But to go into the question of the individual epidemics would carry this discussion beyond its proper bounds.

Stated briefly, the influenza has usually occurred as an explosion, probably showing irregularities in the reporting. The maximum effect has been felt within about ten days after the onset. The duration has been less than three weeks, in most instances.

Within about a week after the outbreak of the influenza, there occurs an ominous prevalence of pneumonia. The pneumonia does not exist as a separate epidemic, but is always a follower of influenza. How the two diseases are related is not positively known. It is clear that the influenza paves the way for the pneumonia, if it does not actually produce it. Most of the pneumonia is of the lobular type and presents various unusual aspects. The time of greatest incidence is usually about a week after the greatest incidence of influenza.

The curves of incidence of pneumonia, when plotted as has already been explained, and with the same scales, are wholly different from those of influenza. The pneumonia does not appear to have been explosive. The curves for the different camps do not resemble one another. One would say that the pneumonia bore no fixed relation whatever to the influenza, except that it followed it.

SPREAD OF THE PANDEMIC

The second camp to report an epidemic, following Devens, was Upton, on Long Island, N. Y.; the third was Lee, in Virginia. Dix, in New Jersey, and Jackson, in South Carolina, followed immediately. Hoboken, N. J., Syracuse, N. Y., Gordon, in Georgia, and Humphreys, in Virginia, all reported on the same day. Within a week from the start, nine large camps in widely separated parts of the country were attacked. Others followed in rapid succession. Table 2 gives the order, according to the daily telegraphic reports.

TABLE 2.—ORDER IN WHICH THE CAMPS WERE ATTACKED

Order	Date	Order	Date
1 Devens.....	Sept. 12	11 Greene.....	Sept. 26
2 Upton.....	Sept. 13	11 McClellan.....	Sept. 26
3 Lee.....	Sept. 17	12 Kearney.....	Sept. 27
4 Dix.....	Sept. 18	12 Bowie.....	Sept. 27
4 Jackson.....	Sept. 18	13 Johnston.....	Sept. 28
5 Hoboken.....	Sept. 19	13 Sheridan.....	Sept. 28
5 Syracuse.....	Sept. 19	13 Sherman.....	Sept. 28
5 Gordon.....	Sept. 19	14 Dodge.....	Sept. 29
5 Humphreys.....	Sept. 19	14 Shelby.....	Sept. 29
6 Logan.....	Sept. 20	15 Custer.....	Sept. 30
6 Funston.....	Sept. 20	15 Hancock.....	Sept. 30
6 Meade.....	Sept. 20	16 Travis.....	Oct. 1
7 Grant.....	Sept. 22	17 Cody.....	Oct. 3
7 Taylor.....	Sept. 22	18 Forrest.....	Oct. 6
8 Sevier.....	Sept. 23	19 MacArthur.....	Oct. 7
8 Lewis.....	Sept. 23	20 Fremont.....	Oct. 8
8 Newport News.....	Sept. 23	21 Wadsworth.....	Oct. 11
9 Pike.....	Sept. 24	21 Wheeler.....	Oct. 11
10 Beauregard.....	Sept. 25	21 Greenleaf.....	Oct. 11
10 Eustis.....	Sept. 25	22 Las Casas.....	Oct. 29

During this period, September 12 to 30, inclusive, thirty-one camps were attacked. The epidemics in this group had subsided in every case by November 1, with the exception of Hoboken, Kearney, Bowie and Shelby. Ten additional camps were attacked during October, completing the list of all the important camps in the United States. Las Casas, Porto Rico, which is last on the list, did not report cases until October 29, on which date an accumulation of 562 admissions was reported.

COMPARISON BETWEEN TENT AND BARRACK CAMPS

Of the thirty camps attacked during September, eighteen were barrack camps, eleven were tent camps, and two were ports of embarkation. During October, eight tent camps and two cantonments were attacked.

The appearance of the pandemic in the tent camps was apparently delayed by reason of their position. They were mostly in Southern states and not immediately adjacent to the seaboard. Speaking generally, the pandemic spread from the sea inland, and from the Northeast to the West and South. There is no reason to suppose that it traveled more rapidly than persons could travel. In cases in which it has appeared to do so, it is probable that infection has been established from foci before the latter were recognized, or from earlier outbreaks.

The pandemic appeared seven days later in the camps than in the cantonments, when these two kinds

of camps are compared as groups. Taking the barrack camps as one group and the tent camps as another, it appears that the disease spread rapidly in both—as rapidly in the one as in the other. There was, in fact, a remarkable similarity in the rate of spread in each kind of camp.

Interest attaches to a comparison between the relative severity with which the tent and barrack camps were attacked for a number of reasons. Among these is the fact that the tents afforded a greater separation of the men than did the barracks. The distribution of cases in cantonments and tent camps during the period of September 12 to October 31, inclusive, is given in Table 3.

TABLE 3.—DISTRIBUTION OF CASES IN CANTONMENTS AND TENT CAMPS, SEPTEMBER 12-OCTOBER 31

	Cantonments	Tent Camps
Camps attacked	20	19
Combined strength	748,632	313,362
Influenza	158,104	69,761
Pneumonia	30,099	9,740
Deaths	10,533	2,923
Per cent. having influenza	21.1	22.2
Per cent. having influenza and pneumonia	19.0	13.9
Case fatality of pneumonia	35.0	30.1

It should be noted that while the number of cases of influenza reported by the cantonments has been more than twice the number reported by the tent camps, the strength of the cantonments has been greater in almost the same proportion. When rates instead of cases are considered there appears to have been less difference in the severity of the pandemic than might be expected. But if there has been as much influenza in the one kind of camp as in the other, there has been no similarity in respect to the incidence or fatality of pneumonia.

The percentage of cases of influenza that developed pneumonia has been greater in the cantonments than in the tent camps; 9 per cent. in the one, against 13.9 in the other. The case fatality of pneumonia has also been greater in the cantonments than in the tent camps.

In order to study more carefully the progress of the pandemic in the cantonments and tent camps, Table 4 was prepared. Here the camps and cantonments are divided into two groups; the figures given are in the form of rates (annual per thousand). These show the suddenness with which the epidemic appeared, how rapidly it rose to its point of greatest intensity, and how it subsided.

TABLE 4.—PROGRESS OF THE INFLUENZA IN THE TENT AND BARRACK CAMPS

Cantonments	Annual Rates	Date	Annual Rates	Tent Camps
First report.....	255	Sept. 12		
	965	Sept. 19	73	First report
Peak.....	5,765	Oct. 4	1,850	
	2,220	Oct. 10	3,865	Peak
	180	Oct. 26	605	Lowest since peak
Lowest since peak..	140	Oct. 28	1,610	
Daily incidence.....	330	Oct. 31	915	Daily incidence

According to this table, the tent camps followed the barrack camps in point of time but not in violence of outbreak.

COMPARISON BETWEEN CAMPS ACCORDING TO THE PROPORTION OF MEN ATTACKED

So much for the comparison between the tent and barrack camps. Turning attention now to the camps with regard solely to the proportion of men attacked,

the camps may be thus divided into groups according to the proportion of men attacked with influenza: Group 1, from 30 to 50 per cent.; Group 2, from 20 to 30 per cent.; Group 3, from 10 to 20 per cent.; Group 4, less than 10 per cent.

Cody leads the first group with 49.8 per cent. of all troops attacked. Other camps in this group are Beauregard, 39.6 per cent.; Wadsworth, 38.6; Bowie, 38.1; Hoboken, 30.5, and Devens, 30.1.

Dodge has had 20 per cent. of all troops attacked and is the first in Group 2. Fourteen other camps fall in this group: Custer, MacArthur, Meade, Pike, Grant, Greene, Funston, Forrest, Travis, Logan, Lee, Hancock, Sheridan and Greenleaf.

Fifteen camps are contained in the third group. Fremont, Lewis and Wheeler, standing at the foot of the list, made up Group 4.

As to fatalities from pneumonia, Grant and Sherman are foremost. At Sherman, 61.3 per cent. of all pneumonia patients have died, and at Grant 45.7 per cent. While the case fatality has been still higher at Wadsworth, Forrest and Greenleaf, the total deaths reported by these camps have been but little more than one third of the total reported by Sherman alone. Special mention should be made of the situation at Wheeler. Only seventy cases of influenza have been reported, but a total of 361 cases of pneumonia and sixty-one deaths have occurred.

SEVERITY IN RELATION TO THE ORDER OF ATTACK

Some interesting facts are revealed when the camps are arranged according to the dates of attack, as shown in Table 5. In this table the camps have been arranged

TABLE 5.—CASES AND DEATHS AMONG TWENTY CAMPS ARRANGED ACCORDING TO THE CHRONOLOGICAL ORDER OF THEIR ATTACK

	Total Cases Infl.	Total Cases Pneu.	Total Deaths Pneu.	Per Cent. Attacked Infl.	Per Cent. Pneu. to Infl.	Per Cent. Deaths Pneu.
Five camps* attacked (Sept. 12-18).....	46,569	7,968	3,000	21.4	17.1	37.6
Five camps† attacked (Sept. 22-24).....	44,000	8,011	2,867	21.4	18.2	35.7
Five camps‡ attacked (Sept. 29-Oct. 1).....	36,044	7,387	2,484	23.8	21.8	31.7
Five camps§ attacked (Oct. 3-11).....	20,096	1,837	579	28.9	8.9	31.5

* Devens, Upton, Lee, Dix, Jackson.

† Grant, Taylor, Sevier, Pike, Newport News.

‡ Sherman, Dodge, Shelby, Custer, Travis.

§ Cody, Forrest, MacArthur, Wadsworth, Greenleaf.

in groups of five wherein are indicated camps reporting epidemics at about the same general stage of the pandemic. This table merits careful study.

It shows that the percentage attacked by influenza increased as the pandemic continued. This is most strikingly indicated in the group which was attacked from October 3 to 11. This group reported 28.9 per cent. of all troops stricken, as compared with 21.4 per cent. in the first group. The inference is that the disease increased in infectivity, or that as the pandemic spread it reached troops who were more susceptible. The latter is probably the fact.

The percentage of influenza cases that developed pneumonia increased as the pandemic proceeded, until the last group (attacked from October 3 to 11) was reached, when a notable drop from 21.8 to 8.9 per cent.

is indicated. This decrease is remarkable. It is probably due to the fact that all the cases of pneumonia that would occur had not yet developed in these camps. Cody, Forrest, Wadsworth and Greenleaf, all in the fourth group, were just passing the peak of influenza during the last four days of October.

It is noteworthy that the case fatality in pneumonia decreased with each successive group.

SEVERITY IN RELATION TO GEOGRAPHIC LOCATION

On considering the camps when grouped according to their location, it has been found that those in the Southwestern section, including the states of Texas, Arkansas, Louisiana and New Mexico, had relatively more men attacked by influenza than did any others. No camp in this group had less than 20 per cent. attacked, and three (Cody, Beauregard and Bowie) exceeded 30 per cent. For the period of the epidemic including October 31, Cody was first on the list of all camps, with 49.8 per cent. of all troops attacked. No explanation is offered for this remarkable fact.

An arrangement of the camps both according to their percentages of men attacked and geographic location is given in Table 6.

TABLE 6.—RELATIVE INCIDENCE OF INFLUENZA IN THE CAMPS ARRANGED GEOGRAPHICALLY

Per Cent. Attacked	South-western	Atlantic Coast	South-eastern	North Central	Pacific Coast
30 to 50	Beauregard Cody Bowie	Hoboken Devens	Wadsworth		
20 to 30	Dodge MacArthur Pike Travis Logan	Meade Lee	Greenleaf Hancock Sheridan	Custer Grant Funston	
10 to 20		Newport News Dix Upton Syracuse Eustis	McClellan Johnston Jackson Shelby Sevier Gordon	Taylor Sherman	Kearney

In the Atlantic Coast group, two camps had more than 30 per cent. attacked; two camps had from 20 to 30 per cent., and five had from 10 to 20. This group is second on the list when the prevalence of influenza is considered.

TABLE 7.—RELATIVE INCIDENCE OF PNEUMONIA FOLLOWING INFLUENZA IN THE CAMPS ARRANGED GEOGRAPHICALLY

Per Cent. Attacked	South-western	Atlantic Coast	South-eastern	North Central	Pacific Coast
30 to 40				Sherman	Lewis
20 to 30	Travis	Meade	McClellan	Taylor Grant Custer	
10 to 20	Beauregard MacArthur Pike Logan	Upton Syracuse Hoboken Devens Lee Dix Newport News	Sevier Greene Hancock Sheridan Jackson Gordon Johnston	Dodge Funston	Fremont
Less than 10	Bowie	Eustis	Wadsworth Greenleaf Shelby		Kearney

One camp in the Southeastern group exceeded 30, and three had from 20 to 30 per cent. of all troops attacked. Of the seven remaining camps of the group, six had from 10 to 20 per cent., and one had less than

10 per cent. The North Central group had no camp that exceeded 30 per cent. Of the five camps considered in this group, three had from 20 to 30 per cent. of men attacked, and two had from 10 to 20. The influenza appears to have been least prevalent in the Pacific Coast group. One camp had from 10 to 20 per cent. attacked, and two had less than 10 per cent.

Pneumonia following influenza was most prevalent in the North Central group of camps. Sherman, in this group, had 35.7 per cent. of all influenza cases develop pneumonia. Three camps had from 20 to 30 per cent.

Lewis, the only other camp to exceed 30 per cent., is in the Pacific Coast group. Two other camps in this group reported less than 20 per cent. Lewis is farthest north. The Atlantic Coast group had one camp with from 20 to 30 per cent. of pneumonias; seven with from 10 to 20 per cent., and one with less than 10. Rates distributed in the Southeastern group were: from 20 to 30 per cent. at McClellan; from 10 to 20 per cent. in seven camps; less than 10 per cent. in three camps.

TABLE 8.—RELATIVE CASE FATALITY OF PNEUMONIA IN THE REGIONAL GROUPS OF CAMPS

Group	More than 60 Per Cent.	40 to 60 Per Cent.	30 to 40 Per Cent.	Less than 30 Per Cent.
Southeastern.....	1*	2	4	5
North Central.....	1	1	2	2
Southwestern.....	1	1	1	3
Atlantic Coast.....	...	2	5	2
Pacific Coast.....	...	...	1	2

* Figures indicate number of camps.

The case fatality of pneumonia has been greater in the Southeastern, North Central and Southwestern groups than in the Atlantic and Pacific Coast groups.

TABLE 9.—AVERAGE PERIOD OF INCREASE, CREST AND RAPID DECLINE OF THE EPIDEMICS IN THE REGIONAL GROUPS OF CAMPS*

Group	Increase (Days)	Crest (Days)	Decline (Days)
Atlantic Coast.....	7	7	5
Southeastern.....	6	9	6
North Central.....	6	4	6
Southwestern.....	5	6	6

* Twenty-five camps are considered in this tabulation.

The increase has been most rapid in the Southwestern group: five days. The Southeastern and North Central groups reached the high point in six days. The Atlantic Coast group was on the increase for seven days. The average period of very high incidence was nine days in the Southeastern group. The shortest period of high incidence was four days, in the North Central group. The period of rapid decline was five days in the Atlantic Coast group, and six days in all other groups.

CASES OF INFLUENZA AND PNEUMONIA AND DEATHS IN THIRTY-EIGHT OF THE LARGEST CAMPS

Table 10 gives for each large camp the number of cases of influenza and pneumonia, percentages of troops attacked, percentages of deaths and the length of the epidemic in days. The camps have been arranged in the order of the percentage of men attacked in each.

Next, in Table 11, is given the total number of cases of influenza and pneumonia and deaths week by week from the beginning of the pandemic to November 1,

for all troops in the United States. This table shows the remarkable activity of the influenza pandemic, considering the camps as a whole. Within these weeks it had reached its acme and had begun to decline. Its

TABLE 10.—INFLUENZA, PNEUMONIA AND DEATHS IN THE LARGE CAMPS, PERIOD SEPT. 12 TO OCT. 31, 1918, INCLUSIVE

	Cases Infl.	Cases Pneu.	Total Deaths	Per Cent. At-tacked Infl.	Per Cent. Pneu. to Infl.	Per Cent. Deaths Pneu.	Duration Epi-demic in Days
Group 1							
Cody.....	2,337	252	46	49.8	10.8	18.2	28
Beauregard.....	5,252	1,007	422	39.6	19.2	41.9	36
Wadsworth.....	5,505	357	60	38.6	6.5	16.9	20
Bowie.....	4,052	119	104	38.1	2.9	89.8	32
Hoboken.....	13,563	2,280	794	30.5	16.9	34.9	42
Devens.....	13,398	2,288	794	30.1	17.1	34.9	49
Group 2							
Dodge.....	9,398	1,847	570	29.0	19.7	30.8	32
Custer.....	11,626	2,437	669	29.0	20.9	27.8	31
MacArthur.....	6,010	852	188	28.1	14.1	20.4	24
Meade.....	11,449	2,013	796	27.8	25.3	27.4	41
Pike.....	13,273	1,379	455	26.7	10.3	32.7	37
Grant.....	10,717	2,335	1,068	25.8	21.8	45.7	39
Greene.....	4,200	626	258	25.8	14.9	41.8	30
Funston.....	13,526	2,328	888	24.9	17.2	37.4	41
Forrest.....	2,307	333	22	24.0	1.1	66.6	25
Travis.....	8,470	1,742	168	24.7	20.5	9.0	30
Logan.....	3,137	393	16	24.6	12.4	3.8	41
Lee.....	11,298	1,919	672	22.9	17.0	34.6	43
Hancock.....	7,715	1,209	462	22.2	15.8	37.4	31
Sheridan.....	4,758	521	132	20.4	10.3	24.6	28
Greenleaf.....	4,747	343	263	20.3	7.2	75.2	20
Group 3							
Taylor.....	11,587	2,800	830	19.2	24.2	29.0	39
Dix.....	9,283	1,673	829	19.0	17.9	49.8	44
Jackson.....	7,500	1,114	362	18.9	14.9	31.3	43
Shelby.....	1,761	94	19	18.8	5.3	24.8	28
Syracuse.....	2,031	401	164	18.4	19.8	40.6	42
Sevier.....	4,526	896	319	16.3	19.7	35.7	38
McClellan.....	4,718	993	218	16.1	20.2	21.3	28
Eustis.....	1,745	67	10	14.8	4.0	16.2	36
Newport News.....	3,897	601	195	14.2	15.5	30.5	34
Upton.....	5,090	974	343	13.9	19.2	34.9	48
Sherman.....	4,789	1,717	1,058	13.5	35.7	61.3	32
Kearney.....	2,450	186	37	13.5	7.9	19.8	34
Gordon.....	4,155	626	192	11.3	14.8	30.4	42
Johnston.....	2,117	383	161	11.1	18.2	40.9	30
Group 4							
Fremont.....	2,347	392	132	9.8	16.1	31.2	23
Lewis.....	3,141	994	148	9.7	31.6	14.8	33
Wheeler.....	70	361	61	0.8	516.0	17.8	

subsidence was as rapid as its rise. In the space of less than two months it had come and gone, leaving only a few cases in some of the camps in its trail.

TABLE 11.—INFLUENZA AND PNEUMONIA CASES AND DEATHS REPORTED AMONG ALL TROOPS IN THE UNITED STATES FOR EACH WEEK DURING THE PERIOD FROM SEPTEMBER 12 TO OCTOBER 31, INCLUSIVE

Week Ended	Cases		Deaths	
	Influ.	Pneu.	Influ.	Pneu.
September 20.....	10,094	758	2	96
27.....	37,493	4,313	19	953
October 4.....	88,478	8,655	13	2,431
11.....	90,393	17,882	165	6,055
18.....	43,779	11,013	270	5,289
25.....	19,996	2,454	256	2,757
November 1.....	16,516	3,004		1,173
Total.....	306,719	48,079	725	18,704

II. THE HANDLING OF THE EPIDEMIC

ACTION BY THE SURGEON-GENERAL'S OFFICE

The steps that were taken to combat the pandemic may be divided into those followed by the direction and guidance of the Surgeon-General's Office at Washington, and those that the camps themselves initiated. These steps may be collectively described as having fallen under three heads, (1) isolation, (2) sanitation and (3) education.

1. By isolation is meant any and all procedures by which infected could be separated from the suscep-

tible persons. Included in this list were steps for the prevention of crowding, quarantine, head to foot sleeping, the separation of heads at mess, and the use of cubicles and masks.

2. Under sanitation may be included the cleaning and airing of barracks and bedding, the oiling of floors to keep down the dust, the boiling of mess-kits and many other procedures.

3. Education, always a predominant motive in the Army, was applied as never before to the prevention of disease among troops. The medical officers were taught what to expect in the way of symptoms and what principles of prevention to put into effect. The men were taught something of the principles of disease transmission and how to carry out their part of the work of prevention. Orders were issued, lectures given, and examples shown. Throughout, the line gave cordial cooperation.

The following paragraphs give a summary of the action taken by the Surgeon-General's Office, and a brief account of what an inspector from that office saw when he visited one of the large camps.

Soon after the pandemic disclosed itself, a recommendation was made by the Acting Surgeon-General to the chief of staff that further increments of the draft be stopped and that all troop movements between infected and uninfected camps be checked. On recommendation, the Adjutant-General ordered the setting aside of buildings to supplement the base hospitals during the epidemic and the furnishing of all necessary unskilled labor for the hospital needs. On September 24, instructions were sent to all camp surgeons regarding the symptoms, epidemic aspects and methods of handling the disease. Large numbers of medical officers and nurses were hurried to the camps. Instructions were sent to buy locally any medical supplies not promptly obtainable through medical channels.

Special sanitary inspectors from the Surgeon-General's Office visited many of the camps with a view to investigating the situation and proposing any improvements that appeared possible in the methods of handling the situation. When thought necessary, camps were quarantined against the civil communities or vice versa, as appeared desirable to the Acting Surgeon-General in order to protect the troops or civilians or both. There was issued a set of a dozen rules on personal hygiene designed to aid the individual in maintaining health during the emergency.

REPORT OF AN INSPECTION OF A CAMP

Under date of September 30, there was sent to camp surgeons, surgeons of recruit depots and independent stations, department surgeons and officers in charge, Air Service Division, the report of a sanitary inspector from the Surgeon-General's Office to show the steps taken at one camp to handle the situation in a satisfactory manner and the recommendations for further improvement made by the inspector. The following abstract is taken from that report.

It should be stated that the policy of the Surgeon-General's Office has been to leave many of the details of camp management, so far as relates to sanitary and medical matters, to the camp authorities on the theory that they alone can be perfectly familiar with the local needs and resources, giving such aid in the way of recommendations as a general knowledge of the situation, and the correct principles to be observed, permit.

The inspection was made two weeks after the epidemic started. The epidemic was at its height: about one in ten of the command was sick with influenza. Of this number about a third were in hospital; the rest were in their quarters. There were 536 cases of pneumonia; there had been 253 deaths.

The report stated that there was no overcrowding. The barracks were marked with their maximum capacity on the basis of 45 square feet of floor space per man, and this was not exceeded. About one fourth of the entire number of troops was under canvas, five men to a tent. The men slept with head and feet alternating to increase the distance between heads as much as possible. In some of the barracks the "cubicle system" was contrived by means of shelter tents suspended between the beds in order to make the separation of the men more complete.

The heating arrangements were in operation in all the buildings, and the freest possible ventilation was enforced. Beds, bedding and clothing were kept outdoors all day, weather permitting. The tents were furled daily. All floors had been oiled once, and some twice, since the epidemic started. Overcoats and woolen underwear had been issued. There was ample bedding. The men who were not sick were kept outdoors all day. An officer was on duty in each barrack day and night.

The camp and barracks and messes were clean. Ample steps had been taken to eradicate flies. Dishes were boiled after each meal. When individual mess kits were used, they were washed in boiling water after each meal and actually boiled at intervals. There were no common drinking cups in use. Dust filled the air on the day of the inspector's visit and was very annoying.

Two thousand negroes who had arrived at camp from civil life between two and five days earlier had been put in an area by themselves and efficiently quarantined, the result being that no case of influenza had appeared among them.

THE HOSPITAL FACILITIES

The capacity of the base hospital had been increased by fitting up a group of eighteen company barracks, fortunately empty, and placing this annex under the base hospital administration. The annex was opened about ten days after the epidemic started, and had a capacity of 1,040 patients. It was described as in excellent condition. Each patient had 100 square feet of floor space; quartermaster cots were used with straw mattresses and the soldiers' own blankets. The Medical Department supplied sheets, pillows, pillow cases and pajamas. There were one female nurse and 186 enlisted men from the line on duty; 100 more nurses were expected. There were 977 patients; of these eighteen had pneumonia.

Ambulant cases, that is, influenza patients whose temperatures had been normal for from twenty-four to forty-eight hours, went out of their particular building to the nearest mess for meals. Men with abnormal temperatures were not allowed to go out to the lavatories. Influenza patients in hospital were being kept from seven to eight days; the rule was to keep them till the temperature had been normal four days.

The attendants wore masks, but cubicles had not been installed. The order and system were reported by the inspector as most commendable.

Further to supplement the base hospital, there was a second group of company barracks. These were run as field hospitals. Their capacity was 700 patients. The conditions here were not quite so good.

The base hospital was stated to be in excellent condition. There were 2,800 patients here, about 300 of whom were venereal and in tents; about 400 others were to be transferred to the annex three quarters of a mile away. There was only slight crowding. The porches were occupied by beds. All pneumonia patients in wards were cubicled, but not those on the porches, though the beds were close together. The influenza patients were not cubicled because of lack of sheets, but there was no serious shortage of supplies. At the outset the camp surgeon directed the camp supply officer to purchase whatever was necessary.

There was one ward full of sick nurses, of whom thirty were said to have pneumonia. One nurse, one dietitian, one medical officer and one dental officer had died. The patients were arranged heads and feet alternatively. The attendants were all masked. The ventilation was ample. The supply of medical officers, nurses, laboratory workers and enlisted men, Medical Department, was being increased. Numerous circulars relative to the prevention of influenza and its complications had been issued by the camp authorities, and these were very complete and satisfactory.

THE INSPECTOR'S RECOMMENDATIONS

In his reports to the Acting Surgeon-General and to the commanding general of the troops, the inspector described the whole situation as well handled. Nevertheless it was recommended among other things that no soldiers be returned to duty after influenza until at least ten days after their temperature had become normal; that all influenza patients should have at least 100 square feet of floor space and be screened from each other, the idea being to prevent pneumonia, and that no influenza patients be treated in barracks that were in part occupied by healthy soldiers.

It was recommended that both in barracks and tents the cubicle system be adopted for all healthy men by the use of the shelter tent supported at one corner of the bed by a stick or hung from the ceiling by wire, and that mess tables be so arranged that the men should sit either on one side of the table alone, or else that the occupants of the two sides of the table be separated by a screen of cheese cloth suspended above the middle of the table.

It was recommended that the quarantine of the camp against the surrounding country, and of the surrounding country against the camp, be continued and that no one be allowed to enter or leave the camp except the friends and relatives of the seriously ill; that as soon as the epidemic disappeared the population of the camp be reduced so as to be placed on the basis of 50 square feet per man; that improvised corridors covered with canvas be provided to connect the lavatories with the barracks which are used for hospital purposes; that steps be taken to prevent men from crowding together at post exchanges and about stoves in barracks; that the laying of dust in camps be expedited, and that great care be taken in disinfecting clinical thermometers.

Some of the recommendations made were already contemplated by the camp authorities or were under way, but not yet completed, the inspector stated, on account of lack of time or material.

III. THE CAUSE OF THE OUTBREAK

CHARACTER OF THE INFECTION

It is usually impossible accurately to trace the origin and paths of spread of a pandemic like the present one. The popular demand that this be done is natural but unreasonable. It rests on the supposition that disease can be run down like a criminal; that visible and recognizable traces are left by it. This is not always true. The pursuit is often so illusive that deductions stronger than inferences are not warranted regarding it. With a disease like influenza, the proof seldom amounts to complete demonstration. The routes of infection are generally hidden beneath a mass of imperfect observation, wrong theory and illogical deduction. Under the circumstances the best that can be done is to follow out the larger aspects of the question.

The present pandemic offers no exception to this rule. In the minds of thousands of persons the disease is hopelessly confused with minor respiratory diseases that have for years been loosely called influenza infections. Many do not know "Spanish influenza" when they see it, the symptoms being by no means easily recognized by those who have never had occasion to deal with it. And to the best informed there are many things about influenza that are shrouded in mystery. Consequently all that will be attempted here is the recording of some apparently well established facts and the drawing of some apparently safe inferences, leaving to others and especially to later investigators the task of interpreting the pandemic more fully.

These remarks seem to be called for by a demand for an explanation of the way in which the pandemic spread from one organization to another in a camp, from one camp to another, and from the camps to the towns and vice versa.

THE MEANS OF TRANSMISSION

First, it is necessary clearly to understand that the influenza to which attention is now called is a specific infective disease like cholera, typhoid, smallpox and others, and it has, at all times and in all places, shown a uniformity in its configuration and in its cause such as almost no other infectious disease has. These words were written nearly two generations ago by Hirsch, but they seem to have been forgotten. They were forgotten in England in 1889, when the pandemic of influenza broke out in that country and for some time failed of recognition. They were forgotten in the United States, where for nearly thirty years no sensational outbreak of the disease had occurred.

The disease is carried from place to place by persons, not things or by the general atmosphere, as was once supposed. Its rapidity of spread is due to its great infectivity, short period of incubation, missed cases, and absence of timely precautionary measures. An epidemic does not easily start, but once the flame is well kindled a conflagration occurs which cannot be stopped. The epidemics stop themselves. This they do either by an exhaustion of the susceptible material, by a reduction in the virulence of the causative agent, or both.

The disease has been taken into camps, from camp to camp, from camps to towns, and from towns to towns, by people. It cannot be said to have been proved that this is the only way in which it has been transmitted, but there seems no reason to look for

more probable ways—according to modern theories, all respiratory infections are conveyed by contact. The travel and mingling that occur under ordinary circumstances seem to be quite enough to account for the facts.

In theory and practice influenza is preventable, but it is very difficult to control under municipal and military conditions. It rarely happens that the necessary measures—chiefly based on the principle of isolation—are taken in time. In the present pandemic the disease, on more than one occasion, has been confined to certain wards of hospitals to the exclusion of others. It has visited some camps and some organizations more severely than others. It is not possible as yet to state to what extent it has been actually controlled in the camps.

The pandemic may truly be termed an epidemic of epidemics. Like all great outbreaks of this most infectious of communicable diseases, the epidemics have occurred with electric suddenness, and, acting like powerful, uncontrolled currents, have produced violent and eccentric effects. The disease never spreads slowly and insidiously. Wherever it occurs its presence is startling. A large proportion of the men in one company may be infected while those in the next company may entirely escape.

How far the pandemic will spread will apparently depend only on the epidemic vagaries of the disease and the amount of material on which it can feed. It is too early to foretell the end or to measure the damage that will be done before it disappears. Apparently it is destined to have a world wide distribution.

It is not improbable that the pandemic may disappear from the United States as rapidly as it came, although most persons hold the opinion that its final disappearance will be gradual, the extinction of the disease being postponed for many months. It is already practically gone in most army camps. It may or may not return. In the pandemics that sweep over the earth at long intervals, recurring waves of greater or less degree, separated by some months, commonly occur.

THE INFECTIVE AGENT

The causative agent is generally believed to be the bacillus of Pfeiffer; the means of transfer, droplets of moisture from the mouth and objects recently contaminated by the buccal and nasal secretions of those who harbor the virus. It is a fundamental assumption that influenza is produced when, and only when, material from the mouth or nose of infected persons gets into the mouth or nose of some one who is susceptible. As is plainly recognized in respect to intestinal infections, the hand probably plays an important part in the transmission of influenza. Coughing and sneezing help greatly to spread the infection.

It has long been known that interchanges of bacteria occur commonly from mouth to mouth under ordinary conditions of social intercourse. Most of these organisms are harmless under normal conditions of health. That their capacity for harm is sometimes increased, sometimes reduced, according to various circumstances, is highly probable. What these circumstances are is conjectural.

The Pfeiffer bacillus is no stranger to America; it was known to be rather commonly present in healthy persons before the pandemic. To account for the virulence that seems to be possessed by this organism now it has been suggested that (1) something must

have happened to increase its virulence, or (2) a new and more active strain of it appeared, or (3) the susceptibility of those attacked became reduced. Some persons hold that the causative agent has not been discovered and that the Pfeiffer bacillus is not really responsible for the disease. Some even go so far as to say that the disease is not influenza. With these disputed points it is not necessary to deal here. All three of the numbered propositions in this paragraph are accepted as probable by the present writer.

The conditions that govern susceptibility to influenza are not understood. Good general health, absence of fatigue and of cold and hunger are methods of prevention which have long been advocated by many and which in spite of scientific criticism still have much to recommend them. Whatever conduces to low bodily tone is believed by most persons to favor infection. Some, however, hold that specific immunity either does or does not exist and that wet feet, insufficient bedding, chill, hunger and fatigue have nothing to do with susceptibility.

Vaccination against pneumonia is practicable; but such preventive treatment is in the experimental stage as respects influenza. As to natural immunity, one attack is believed to protect against another, and some people seem to be immune without ever having experienced an attack.

The weather has always been supposed to exert an influence on influenza—the very name is derived from the effect which extraterrestrial conditions were supposed to exert on it. But although there has been a great deal of study of this matter from early times to the present, little is known concerning it. It is possible that the weather this fall has aggravated the influenza and contributed to the incidence of the pneumonia. The pandemic appeared after a very hot season had suddenly been succeeded by a very cold one. After the pandemic was well started, a particularly long period of fine weather occurred. More important, perhaps, the weather also precipitated a large amount of other respiratory infections, such as always become prevalent at this season. There is little doubt that many of these cases have been mistaken for influenza.

The epidemics which occurred in the spring of 1918 were like those that took place in the fall, except that the disease was milder and there was less pneumonia in the spring. Until the late summer the influenza reported from Europe was of a relatively mild type. It seems to have been as infectious there as it is here. Reports coming from all parts of Europe indicate that the percentage of persons attacked was about the same as in America.

IMPORTATIONS OF A VIRULENT STRAIN

The origin of the influenza pandemic now sweeping the world is usually attributed to Spain. It is doubtful, however, whether the disease was first seen in Spain or the United States. Some have held that it was brought to the Western hemisphere from China.

Early in the spring of 1918, a disease of unknown origin is reported to have appeared in Spain with the symptoms of sudden onset, severe headache, pain in the back, fatigue and perhaps nausea. After three or four days the patient recovered. Influenza appeared in America early in March, being reported from Fort Oglethorpe, March 18, and from some other parts of the United States much earlier.

In the latter part of March the disease appeared in the American Expeditionary Force, in the French and

British army, and in the civil population. About 25 per cent. of all who were exposed to the disease were attacked. The death rate was low. The first cases in Germany were among troops that fought with a (presumably infected) American regiment in July.

In September, the disease assumed a more serious character, being complicated with bronchopneumonia and septic pneumonia with a high mortality in some epidemics. The disease is said to have become more severe in Spain as the season advanced.

The first that was seen of the highly virulent infection in America was among shipping. During July and August a number of vessels plying between Europe and America experienced intense outbreaks of what was called "Spanish influenza." That patients with the disease were being brought into the country in this manner was stated in the daily press and in official reports in July.

Many patients were admitted to various ports of the United States from ships during June, July and August. The Navy has pointed out that the British steamship *Exeter*, which arrived from Liverpool, at Philadelphia, June 22, reported at quarantine that twenty-seven Lascars and an English quartermaster were ill with pneumonia. These patients, who were described as in a desperate condition, were removed to hospital. The opinion was reached that they had influenza.

July 7, the New York papers described an outbreak of "Spanish influenza" which had occurred on a troop ship returning to America. The vessel was compelled to put back into port, there not being enough of her crew left to handle her. This vessel was the Cunard Liner *Khiva*, which had left Liverpool, June 15. Six days later six of the crew were taken suddenly ill. Two days later the number was increased by fifteen. Next day there were fifteen more cases. The ship then, pursued by a submarine, put into Halifax. Ten days later it proceeded to New York. There were no deaths.

August 18 and subsequently, the New York papers described outbreaks of influenza on the *Rochambeau* and the *Nieuw Amsterdam*. On one of these vessels, twenty-one cases of influenza had occurred. Some of them developed pneumonia and were sent to St. Vincent's Hospital. According to the New York *Times* of July 18, 1918, influenza had been coming to New York on vessels since the early part of June.

According to a Navy bulletin, two steamships from Norway and one from another Scandinavian port arrived in New York, August 14 and 15, having a number of patients ill with influenza.

The Surgeon-General of the United States Public Health Service reported a sharp outbreak at Fort Morgan, near Mobile; Ala., during the month of August. At the same time a steamship arrived at Newport News with substantially the entire crew infected. The *Bergensford*, which arrived in New York, August 12, had during the voyage 200 cases of influenza.

The foregoing, which makes no claim to being a complete list, is sufficient to show that influenza was brought to the United States and to various ports on vessels from Europe through the whole summer. In some outbreaks there was a great deal of a fatal pneumonia.

The patients from the vessels were sent ashore and soon mingled with the civilian populations. There were thus scattered rather widely along the Atlantic seaboard sparks from which the pandemic not improb-

ably arose. How abrupt were the appearance and spread is not yet perfectly clear. In arriving at an understanding of this point it is necessary to distinguish between the sudden appearance of the disease and its sudden recognition.

There is another and less definite explanation of the pandemic offered. It is less an explanation than a guess. The same kind of guess probably has been made ever since epidemics first visited the earth. It is that the marked virulence noted in the recent pandemic is due to the weather. The supposition is that in some way not determined, the mild influenza of last spring became greatly intensified this fall.

That the disease appeared suddenly in the camps and ran a meteoric career is clear enough. Evidence presented in Table 12 indicates that its appearance and course in civil life was somewhat the same. This table, compiled partly from data supplied by the Census Bureau, shows the dates when marked increases occurred in the number of deaths from pneumonia in

It was two weeks later (week ended October 19) that the camps of the Pacific Coast group reported the first marked increase in deaths from pneumonia.

Table 13 gives the number of deaths from pneumonia reported in forty-six cities from September 14 to November 2. It shows the rise and progress of the pneumonia, and, so far as that disease is the shadow of the influenza, it affords evidence of the case incidence of the whole pandemic of the larger cities. No reliable evidence of the prevalence of influenza in the cities exists. This statistical table is based on figures contained in a special report of the Bureau of the Census, Washington, D. C., on deaths from pneumonia, and appended to the regular weekly report of that bureau for the week ended Nov. 2, 1918.

The special report covers the eight weeks period from September 8 to November 2, inclusive. Marked increases in the incidence of pneumonia deaths in the different cities are indicated by the totals for the week being printed in boldface. A weekly incidence which

TABLE 12.—MARKED INCREASES IN DEATHS FROM PNEUMONIA IN THE PRINCIPAL CITIES AND ARMY CAMPS IN THE UNITED STATES FOR THE PERIOD FROM SEPTEMBER 8 TO OCTOBER 25, INCLUSIVE

Date Week Ending	Atlantic Coast		North Central		Southeastern		Southwestern		Pacific Coast	
	Cities	Camps	Cities	Camps	Cities	Camps	Cities	Camps	Cities	Camps
Sept. 14	Boston									
Sept. 21	Lowell New York	Devens Dix								
Sept. 28	Cambridge Worcester Jersey City Philadelphia Providence Washington	Lee Upton Syracuse Humphreys	Pittsburgh Chicago Minneapolis							
Oct. 5	Baltimore Buffalo Fall River Newark Richmond Syracuse	Meade Hoboken Eustis Newport News	Cincinnati Indianapolis	Custer Funston Grant Sherman Taylor	Birmingham	Beauregard Sevier Logan Hancock Gordon	Denver	Pike	Los Angeles	
Oct. 12	Albany Rochester		Kansas City Louisville Milwaukee Omaha St. Paul St. Louis	Dodge	Atlanta New Orleans	Greene McClellan Wadsworth Jackson Johnston Greenleaf		Travis Bowie	Oakland Seattle	
Oct. 19	New Haven		Cleveland Columbus Dayton Toledo			Sheridan		MacArthur Cody	San Francisco Spokane Portland	Fremont Lewis

cities and camps in various parts of the country. From the table it would appear that the increase in the incidence of pneumonia among the principal cities of the United States preceded the increased incidence among the Army camps. If this is so, it is to be inferred that the camps derived their infection not from one another but from their immediate environment.

This table shows how the pandemic spread across the continent. Both cities and camps in the Atlantic Coast groups were included. The states of Massachusetts, New York, New Jersey, Maryland and Virginia were attacked first. The first record of the spread of the pandemic near this group was the increased pneumonia death rate at Pittsburgh, Chicago and Minneapolis, cities of the North Central group, during the week ended September 28.

During the following week (ended October 5) pronounced epidemics occurred in five of the Army camps of the North Central group (Custer, Funston, Grant, Sherman and Taylor). During this week epidemics were reported by both cities and camps in the Southeastern and Southwestern groups, and in the city of Los Angeles of the Pacific Coast group.

is more than double the average weekly figure for not less than two weeks preceding (when these figures are available) is considered a marked increase in this memorandum.

COMPARISON WITH OTHER PANDEMICS

It is interesting to compare the present pandemic with others, but the comparison cannot be close. It is impossible to say how severe were some of those which are recorded in history for the reason that statistical data concerning them are meager and imperfect. It is said that in 1889-1890 no less than 25 per cent. of the population was attacked in London; 33 per cent. in Antwerp; 39 in Massachusetts, and 50 in Paris. In 1823-1833 about 40 per cent. of the population of Paris is believed to have been affected. In 1872, three quarters of the population of London and some German cities were supposed to have suffered. The records of earlier visitations are more obscure.

In the 1918 pandemic, about 20 per cent. of the soldiers in the camps were attacked. This seems rather a large proportion for the whole country or, indeed, for the cities, and yet it may not be far amiss. By the

time the disease has finally disappeared, it will have spread somewhat beyond its present limit; and if the patients that recovered in the Army are added to those of the full pandemic, the proportion of the whole population affected during this year will be found to have been still greater.

The total number of deaths resulting from the present pandemic will never be known. Observers of pandemics in other years have pointed out that influenza is a more fatal disease than is commonly understood, the unsuspected excess being due chiefly to lung and heart complications that do not promptly manifest themselves. Thus, although the number of deaths directly attributed to influenza in England and Wales

TABLE 13.—DEATHS FROM PNEUMONIA REPORTED BY THE PRINCIPAL CITIES OF THE UNITED STATES FOR THE PERIOD SEPT. 8 TO NOV. 2, 1918, INCLUSIVE

City	Week Ended							
	September			October				Nov.
	14	21	28	5	12	19	26	
Albany.....	2	0	2	2	11	22	34	24
Atlanta.....	7		4	7	25	42	86	
Baltimore.....	7	5	19	87	371	793	600	204
Birmingham.....	2	2	4	11	9	12	19	15
Boston.....	27	93	175	225	177	126	52	38
Buffalo.....	8	12	16	41	98	125	134	90
Cambridge.....	7	4	23	27	15	13	4	6
Chicago.....	15	24	74	246	476	863	826	456
Cincinnati.....	5	4	6	15	20	21	19	14
Cleveland.....	5	8	10	15	22	64	106	153
Columbus.....	1	4	6	10	13	28	21	9
Dayton.....	1	1	2	5	11	38	43	31
Denver.....	3	3	8	19	25	19	34	19
Fall River.....	0	5	6	11	9	23	10	
Grand Rapids.....	2	1	1	3	7	6	15	
Indianapolis.....	3	6	10	24	16	110	97	
Jersey City.....	2	6	21	50	144	225		
Kansas City, Mo.....	5		10	11	21	55	75	55
Los Angeles.....	9		2	14	41	81	49	53
Louisville.....	3	9	4	14	77	154	157	60
Lowell.....	1	8	16	64	104	85	65	19
Memphis.....		4			67	152	127	60
Milwaukee.....	4	5	13	15	54	83	136	90
Minneapolis.....	2		11	13	11	10	22	
Nashville.....	2		3	5	29	47	27	13
Newark.....	0	6	6	22	70	126	137	138
New Haven.....	0	2	4	6	9	22	29	
New Orleans.....	5		3	17	49	177	233	117
New York.....	74	98	145	434	1,142	2,099	2,251	1,855
Oakland.....	4	0	2	3	11	21	27	
Omaha.....	1		2	7	22	22	7	
Philadelphia.....	20	32	76	307	938	1,644	1,132	467
Pittsburgh.....	12	17	33	57	102	161	457	537
Portland.....	4		3	8	5	25	11	16
Providence.....	4	10	2	38	61	62	48	32
Richmond.....	4	1	3	20	42	73	43	
Rochester.....	1	2	7	6	22	25	28	33
St. Louis.....	12	13	20	25	45	76	75	87
St. Paul.....	3		3	1	30	11	10	
San Francisco.....	6	14	15	15	19	42	58	59
Seattle.....	4	2	4	6	16	22	7	17
Spokane.....	2		1	1	2	5		
Syracuse.....	0			14	3	50	32	
Toledo.....	2	0	5	3	7	19	43	41
Washington, D.C.....	10		17	57	101	96	61	36
Worcester.....	7	9	38	68	53	38	22	14

in 1890 was reported as 4,523 per million, the registrar general, by analysis of the vital statistics for the period, stated that the number of deaths directly or indirectly attributable was 27,074 per million, or nearly seven times the apparent rate. In London the general death rate was increased by more than 20 per cent., in Berlin by more than 60 per cent., and in Paris and Brussels by more than 100 per cent.

No records now available show that there has ever been so much immediately fatal pneumonia as in the present pandemic. If to the known deaths are added an allowance based on the registrar general's estimate quoted above, the magnitude of the catastrophe appears truly appalling. Both in and out of the camps there will have been fewer cases of influenza and far more deaths than the records indicate. The cases will be overestimated—the deaths will not all be told.

BARRACK LIFE AND RESPIRATORY DISEASE

SOME EPIDEMIOLOGIC OBSERVATIONS ON THE RECENT OUTBREAK OF INFLUENZA *

VICTOR G. HEISER, M.D.

NEW YORK

The influenza epidemic through which we are passing has been responsible for 17,000 deaths among our troops in this country, and the toll among the civil population already exceeds 200,000 lives. This is a far greater disaster than has befallen the United States in many decades. We have come to look on the war in Europe as one of the greatest destroyers of human life in the history of mankind. Yet in the brief space of a few weeks more deaths have been caused by an acute disease than have occurred on all the battlefields in any similar period, notwithstanding that millions on millions of soldiers have been involved.

Among the civil population of New York, 50 per cent. of all deaths occurred between the ages of 25 and 45. From figures obtained from Boston, Philadelphia, Baltimore and other sources, it can be predicted that similar percentages will probably obtain in many other communities. The deaths in cities among the sexes appear to be about equal, although there was a slight excess among females in Boston. Thus, 100,000 persons between the ages of 25 and 45 have succumbed to respiratory disease. This means that we have already lost among civilians 50,000 men in the prime of life. This is a serious drain on our man power.

Viewed from another standpoint, it is apparent that it is more dangerous to be a soldier in peaceful United States than to have been on the firing line in France. From the statistics so far available, the death rate in the military camps is higher than among the civil population, even in similar age groups. The mortality in New York and Chicago, for instance, shows that the death rate in the Army is more than double that of the civil population of the same age group. There is also the possibility that when allowances are made for the fact that defectives have been eliminated from the Army, and that these poor "risks" swell the civil death list, the corrected margin will be still further increased in favor of the civil population.

A study of the Students' Army Training Corps and of vocational students shows that as living conditions approach those in the Army, so does the death rate approach that of the Army. Further observation shows that with few exceptions the mortality rate from influenza and pneumonia bears a very close relationship to the number of persons housed and living under barrack conditions. That is, when large numbers of persons live, sleep and mess together in one room, the death rate is almost invariably higher than among similar groups living under civilian conditions. In other words, the more nearly the housing, eating and congregating conditions in the Students' Army Training Corps resembled those which obtain normally at colleges or such as exist in civil communities, the death rate was correspondingly lower. The death rate among college students living in their customary way was about the same as that of the towns in which they were located.

* The data assembled in this paper are based on reports to Nov. 9, 1918, but subsequent deaths are more likely to emphasize the difference between civil populations and "barrack" populations than otherwise.