THE VELOCITY OF ACCOMMODATION.—Dr. Banet has made a series of experiments upon this point and arrived at the following conclusions: The accommodation-apparatus acts with varying degrees of rapidity according to different circumstances, the principal of which are age, lesions of refraction, practice (?), individual characteristics, time of day, and fatigue. Relaxation occurs with very much greater rapidity than active accommodation. That with two eyes of much the same refractive character the accommodation velocity of either, measured separately, is about the same as that of the two acting in concert. That in the present state of knowledge no absolute data can be given as to the time occupied by the action of the local mechanism for any given distance.—*Journal of Physiology*, vol. vi., Nos. 1 and 2.

THE SALIVARY SECRETION.—Dr. J. N. Langley has studied this subject by experiments upon animals. Bernard was the first to show that when the chorda tympani is cut in the dog, there follows a slow continuous secretion of saliva from the sub-maxillary gland. The cause of the secretion was sought for by Langley, and he arrived at the following results: When the chorda tympani is cut on one side, the central secretory centre gradually increases in irritability until the blood supplied to it serves as a stimulus. Nervous impulses are in consequence sent down the sympathetic fibres which are still connected with the sub-maxillary gland and the paralytic secretion begins. The increase of irritability of the central secretory centre is not confined to the side on which the chorda tympani is cut, but extends, although to a less degree, to the nerve cells of the opposite side. If all the nerve fibres running to the gland are severed, the paralytic secretion is caused by the peripheral centre only. The state of the gland suggests also that the growth of protoplasm is not brought about by the secretory nerve fibres.—*Journal of Physiology*, vol. vi., Nos. 1 and 2.

RESPIRATORY CENTRES.—Dr. Richard Nitschman has made a series of experiments upon chloralized rabbits, and arrived at the following conclusions about the centres of respiration. There are automatic respiratory centres in the cervical spinal cord. These cervical centres of respiration can be longitudinally divided without the respiration experiencing any change, so long as the division is in the median line. After longitudinal division of the cervical cord and the calamus scriptorius, the respiration remains alike on both sides of the body, but upon irritation of the vagi, trigemini, and brachial it is not synchronous, whilst irritation of the sciatic causes both halves of the diaphragm to act alike.—*Pflüger's Archiv*, Band xxxv., 10, 11, and 12 Heft.