Bell, A. S. CLINICAL METHOD OF DETERMINING TYPE OF MENINGOCOCCUS. [Lancet, May 24, 1919.]

Bell describes a simple and quite accurate method for typing the meningococci obtained by culture from the spinal fluid in acute cases. The results can be secured in twenty to twenty-four hours, as compared with the seventy-two hours or more usually required. The spinal fluid is cultivated and a very concentrated emulsion of the organism is prepared. One drop of each of the four type sera is placed on a glass and with each is mixed one drop of the culture. The mixture is examined with a small magnifying glass and the type is indicated by the serum with which agglutination first appears. If the result is not clear, or is negative after five minutes, pools of each serum are prepared containing two, three, and four drops each of serum. To each of these one drop of the bacterial emulsion is added and the results noted as before. If again the agglutination is not specific pools are prepared containing three, six, and twelve drops of the bacterial emulsion and the test is again applied by adding to each of these one drop of type serum, three tests being made with each serum. The results should always be confirmed later by the slow method usually employed.

Carter, A. H., and Boyes, J. T. CEREBROSPINAL FEVER. [Lancet, June 21, 1919, J. A. M. A.]

One hundred cases of epidemic meningitis chiefly due to the meningococcus were analyzed by Carter and Boyes. The average age of these patients was 23.05 years. Certain camps, without a necessarily larger body of troops present, furnished a larger number of them, and more fatal cases, spread over the whole period. No satisfactory reason for this is forthcoming. No seasonal or meteorological condition can be shown to have influenced the incidence or death rate, except that the larger number of cases occurred in the three winter months, December, January and February. No single symptom can be considered pathognomonic of the disease, and even a combination of the principal symptoms may not be positive. A positive bacteriologic finding alone is conclusive. No treatment appears to be so effectual as early continued lumbar puncture and injection of serum. Wherever possible a serum of the particular type affecting the patient is preferred. The most frequent and the most fatal type was Type 2.

Nelli, L. SYMPTOMS OF EPIDEMIC CEREBROSPINAL MENINGITIS. [Gazz. degl. Ospedale, 40, April 24, 1919.]

This observer has analyzed one hundred cases and states that headache, vomiting and herpes were seen in every case. Bradycardia, even down to 35 and 40 in some cases, was constant. The pulse becomes more rapid in the graver cases. Purpura with swelling of the joints was present in one patient. In two patients there was no pyrexia, yet these patients died.