№	Star	AR 1870	Decl. 1870	Period days	Change of Magnitude		Discoverer
					From	to	
116	Star in Aquarius	22 ^h 21 ^m 31 ^s	—10° 39' 0	..	8	0	<i>Rümker.</i>
117	δ Cephei	23 24 20	+57 45,0	5.3664	3,7	4,8	<i>Goodricke</i> 1784.
118	S Aquarii	22 50 8	—21 1,8	..	8	11 <	<i>Argelander</i> 1853.
119	β Pegasi	22 57 27	+27 22,6	31,5 or 43,4	2	2,5	<i>Schmidt</i> 1848.
120	R Pegasi	23 0 7	+ 9 49,1	378	8,5	13,5	<i>Hind</i> 1848.
121	T Cephei	23 14 43	+55 19,9	..	8,2	8,8	<i>Argelander</i> 1863.
122	R Aquarii	23 37 46	—15 59,7	354 or 388,5	7	10 <	<i>Harding</i> 1810.
123	R Cassiopeiæ	23 51 49	+50 39,9	434,81	6	14 <	<i>Pogson</i> 1853.

2, Palace Gardens Terrace, Kensington, June 1864.

George F. Chambers.

On a new variable Star (S Delphini). By *Joseph Baxendell*, Esq.

On the 24th of October last I observed a strange star of the 8,6 magnitude at a distance of about 25 minutes from the fine double star γ Delphini, in the north preceding quadrant. It remained for several days without any very decided change; but afterwards gradually diminished in brightness, and at the end of December had fallen below the 12. magn. It was frequently looked for after its conjunction with the sun, but not seen till the night of July 29, when it was found to be of the 13,2 magnitude. On August 21 it had risen to the 9,4 magnitude, thus showing a variation of nearly four magnitudes in 23 days. It has since continued to increase, though much less rapidly, and last night, Sept. 5, it was of the 8,4 magnitude. Its approximate place for 1860,0 is:

$$20^{\text{h}} 38^{\text{m}} 51^{\text{s}} 9, \quad +15^{\circ} 53' 5.$$

It does not occur in the „Bonner Sternverzeichniss“, nor in any other Catalogue I have yet met with. As this is the

second telescopic Variable discovered in Delphinus, it will, in conformity with Prof. *Argelander's* system of nomenclature, be denoted by the letter S . The first Variable discovered in this constellation was incorrectly designated S Delphini in my note in the „Monthly Notices“, Vol. XXI, pag. 68, as I had then an impression, that another star in Delphinus had been discovered to be variable by Dr. *Henke* of Driesen; but I found afterwards that Dr. *Henke's* star was in Aquila, its position for 1855,0 being: $20^{\text{h}} 7^{\text{m}} 55^{\text{s}} 1, \quad +8^{\circ} 38' 8$.

The new Variable affords another illustration of the apparent tendency of variable stars to occur in groups, as it is less than one degree distant from R Delphini, and there are the Variables R Sagittæ, T Aquilæ and S Aquilæ (*Henke's* star) in the immediate neighbourhood.

Manchester, 1864 September 6.

Joseph Baxendell.

Beobachtungen auf der Sternwarte zu Athen. Von Herrn Director *J. F. Julius Schmidt*.

Comet I. 1864. (Fortsetzung).

In meinem neulichen Schreiben habe ich Ortsbestimmungen des Cometen bis Sept. 4 mitgetheilt. Nach diesem Tage trat eine kurze Regenperiode ein, und erst Sept. 9 erhielt der Himmel wieder seine gewöhnliche Klarheit. Während des Vollmondes blieb der Comet zwar am Refractor sichtbar, wenn er auch nur 10° bis 5° hoch stand, doch unterliess ich die Beobachtung, weil ein schwacher Scirocco, und bei sehr hohen Temperaturen die Unruhe der Luft eine genaue Ortsbestimmung doch nicht zuließ. Nach dem Vollmonde hatte

der Comet noch über 2' Durchmesser, und einen Kern der 11^m. Am 20. Sept. sah ich den Cometen zuletzt in einer wahren Zenithdistanz von 88° 6'; September 22 habe ich ihn zwischen Wolken zuletzt in 87° 56' wahrer Zenithdistanz gesehen und beobachtet. Die folgenden Oerter können die Genauigkeit der früheren nicht beanspruchen, da die Höhe des Cometen stets nur (in den letzten 5 Tagen) zwischen 7° und 3° betrug, und es wird wohl besser sein, die Positionen von Sept. 19 bis Sept. 22 nicht weiter zu berücksichtigen.