The Journal of the American Medical Association

VOL. XXXV

CHICAGO, ILLINOIS, NOVEMBER 3, 1900.

No. 18

Original Articles.

ACUTE SENILE ENDOMETRITIS.

CASES AND PATHOLOGY.*

L. H. DUNNING, M.D.

INDIANAPOLIS, IND.

Professor Diseases of Women, Medical College of Indiana, University of Indianapolis.

Since the publication of Skene's notable¹ article in 1894 on "Senile Endometritis," the lesion has attracted considerable attention, and text-books now include this form in the classification of endometritis. The disease as described by Skene is essentially a chronic one, and while his exposition of the subject is true and graphic, it can not be made to include cases of the type of those, the histories of which I shall present to-day.

In his admirable² article entitled "Post-Climacteric Endometritis," Sexton treats of the same lesion described by Skene and calls it a chronic inflammation. His cases were in one particular different from those here presented, in that there was no hemorrhage; indeed, he emphasizes the fact that there is an absence of hemorrhage.

Mundé, in his article³ read before the American Gynecological Society in 1896, wrote very clearly on the clinical history of this class of cases, but he does not attempt to add to our knowledge of the pathology of the lesion.

The author believes the pathologic findings in his cases clearly indicate that those described by Mundé, as well as his own, are at the onset acute in character and must, if properly classified, be denominated acute senile endometritis. The sole object of his paper is to endeavor to demonstrate this fact.

It has seemed to the writer that the histologic study of endometrium in senile endometritis has been limited to microscopic examination of scrapings, and he deems himself fortunate in being able to present the results of a microscopic study of the endometrium in two cases in which the uterus has been removed by hysterectomy.

So important has the writer deemed this subject that he has been minute in his descriptions of the histories of his cases and the pathologic findings, at the risk of subjecting himself to the charge of being tedious.

CASE 1.—Mrs. M. M., aged 63 years, was referred to me by Dr. J. S. Sherfee, of Fairland, Ind., Jan. 22, 1900.

She presented the following history: She had never borne children, though married since early womanhood.

*Presented to the Section on Obstetrics and Diseases of Women, at the Fifty-first Annual Meeting of the American Medical Association, held at Atlantic City, N. J., June 5-8, 1900.

¹ Trans. American Association of Obs. and Gyn., Vol. viii, pp. 314-319.

² New York Gyn. and Obs., June, 1894, p. 641.

³ Gynecological Transactions, Vol. ii, p. 29.

Her health had always been good until the time of menopause, which occurred at 40 years of age. At that period she was confined to her bed for six months on account of illness, the nature of which I am unable to ascertain; but subsequently she enjoyed fairly good health until six months ago, when without apparent cause she began having a seropurulent discharge from the uterus.

In spite of general and local treatment this discharge continued, and finally became bloody and exceedingly offensive. She complained of backache, pain and soreness through the pelvic organs and inability to walk or stand many minutes at a time. She had lost several pounds in weight and was noticeably anemic.

The sanguineous discharge was not profuse, but was of dark color and exceedingly offensive. A physical examination showed a small, short vagina, here and there in patches, eroded and quite sensitive to the touch. The cervix uteri was atrophied and was but slightly invaginated. The uterus was small and retroverted, though not adherent. There was no difficulty in passing a uterine sound. The withdrawal of the sound was followed by the discharge of probably two drams of dark, offensive fluid. The left Fallopian tube and ovary could be easily palpated. They were thought to be enlarged. Their palpation caused a great deal of pain.

My diagnosis was senile endometritis and inflammatory lesions of the left tube and ovary. Hysterectomy was recommended. It was performed at the Deaconess Hospital, Jan. 23, 1900. The operation was an easy one and the patient made an excellent recovery.

The left tube was considerably thickened but was not distended, and the left ovary contained two or three small cysts. Unfortunately before I saw the importance of a microscopic examination of the tube, it was lost. There were found evidences of a slight, recent, pelvic peritoneal inflammation manifested by adhesion of the diseased tube and ovary.

With the aid of my assistant, Dr. R. L. Ritter, a critical and microscopic study of the uterus was made, with the following results: In general appearance, it exhibited the characteristics one would expect to find in the uterus of a woman at the age of the patient. The organ was pale and perceptibly smaller than a sexually active one. It measured $2\frac{1}{2}$ inches in length, $1\frac{5}{8}$ inches in width at its widest points, and about $\frac{3}{4}$ inch at its thickest portion anteroposteriorly. The cervix was short, quite dense; the external os was rather flaring, the cervical canal patulous. There was no evidence of laceration of the cervix. There was absolutely no sign of disease of the organ to be seen externally, with the exception of a thick, dark-colored, granular material which exuded from the external os on pressure. On splitting the organ longitudinally the walls were found to be firmer or denser than the muscular tissue of a functioning uterus, and the muscular tissue somewhat paler in color. The walls were about $\frac{3}{8}$ inch in thick-

ness. Covering the entire mucous surface and filling the uterine cavity was the same material which could be forced out of the cervical canal by pressure. Part of it was breaking-down, liquefying blood-clot, diluted with what seemed to be pus colored by liberated hemoglobin of the clot. This material was thick and sticky, but could be scraped off easily, leaving a roughened, reddened endometrium below. The cervical mucosa did not seem to be greatly affected. When this material was thoroughly washed off, the surface of the mucous membrane was seen to be studded apparently with little nodules resembling somewhat the skin of a frog's back.

Thin sections stained with hematoxylin and eosin, photomicrographs of which are herewith submitted, show an endometrium thickened, densely infiltrated with leucocytes and highly vascular. All the superficial epithelium is lost, the roughened surface being covered

The musculature of the organ shows evidences of atrophy such as normally takes place after menopause. The arteries found chiefly in the middle layer of muscle also show degenerative changes. Their tunica intima is markedly thickened and consists of a rather homogeneous structureless material showing here and there a few small, shrunken, free-lying nuclei evidently of degenerated subendothelial connective-tissue cells. All the changes I have described are found best shown in the body of the organ proper.

In conclusion I would say, we have a uterus showing decidedly the atrophic changes of senility, yet with a mucosa the seat of an active persistent inflammatory process.

CASE 2.—Mrs. V., aged 63 years, was referred to me by Dr. Theodore Potter, Dec. 5, 1899. She gave the following history: She had been married 42 years and was the mother of 8 children, the youngest 24 years

Figure 1.

CASE 1.—Section showing mucosa and a portion of the muscularis. 1. Endometrium. a. Free surface of mucosa. b. Degenerating and disintegrating glands of mucosa. 2. Upper layers of muscularis, showing round-celled infiltration. 3. muscularis.

with a layer of material composed of granular debris, epithelial cells and leucocytes of various kinds, chiefly of the small mononuclear variety, "lymphocytes," and polymorphonuclear or pus corpuscles.

The epithelial cells are found chiefly near the orifices of the glands. These latter are greatly diminished in number, their course is quite tortuous and their epithelium in many cases is desquamating, filling the lumen of the gland with a mass of granular debris and cells in all stages of degeneration. There is no sharp-dividing line between the intact mucous tissue and the material on the surface. The minute blood-vessels with which the endometrium is abundantly supplied often run to the surface and seem to end at the surface. From these changes I am led to believe that there exists a purulent inflammatory process in this endometrium, subacute in character, or else at the time the organ was removed there was an acute exacerbation of a more chronic process. I believe the hemorrhage which had evidently taken place from the mucosa was due to the erosion of the minute blood-vessels described and whose blind ends may be seen at the surface. It has been a direct hemorrhage rather than a diapedesis.

Figure 2.

CASE 2.—Endometrium. a. Free surface of mucosa. b. Remains of glands.

of age. She passed the menopause twenty years ago. Her health had always been good. She had the first leucorrheal discharge of her life four weeks ago.

One week later, viz., three weeks previous to my examination, a bloody discharge appeared. It was slight at first, but had gradually become more abundant. This discharge had an offensive odor from the first. The discharge was thick and dark, and seemed to me to be pus stained by the coloring matter of blood. On examination the vagina was found small but elastic. The upper portion was slightly reddened and showed slightly ulcerated patches. The uterus was in normal position, the cervix short, and gave evidence of having been lacerated bilaterally. The external os was of sufficient size to admit the end of the small finger. The internal os, while not patulous, permitted the easy passage of an ordinary uterine sound. There seemed to be a considerable cavity in the body of the uterus and the uterine walls were thin and elastic.

A considerable amount of dark-colored sanguinopurulent fluid flowed out of the os on withdrawal of the sound. A small, sharp curette was introduced into the uterine cavity and an effort made to remove portions of tissue for microscopic examination. Only small shreds

of tissue and dark-colored blood-clots and débris were obtained. I was inclined to believe that we had to deal with a case of incipient cancer. Dr. Potter assented to this view, and we agreed upon a hysterectomy.

The operation was done at St. Vincent's Hospital, Dec. 22, 1899. No neoplasm was found in the uterus. The appendages were removed. The right ovary and tube showed the atrophic changes incident to the age of the patient. The left ovary contained a small cyst not larger than a hazel-nut. There was no pelvic peritonitis.

On examination the uterus was found to closely resemble the preceding one in appearance. There was the same thickened endometrium densely infiltrated with small round cells. This infiltration involved also the upper layers of the muscularis. The superficial epithelium normally present was entirely gone, and the surface covered with an indefinite layer of a confused mass of small round cells, larger connective-tissue cells,

The writer greatly regrets not having procured a bacteriological examination of the contents of the uterine cavities in these cases and the loss of the diseased tube and ovary in the first case.

Fortunately in Case 2, one tube and ovary were still attached to the uterus and were examined microscopically. The endometrium surrounding the entrance of the tube into the uterine cavity showed marked inflammation. While it was impossible to pass even a bristle through the uterine portion of the tubal canal, yet on microscopic examination its lumen was shown to be pervious. For a short distance away from the horn of the uterus the mucous membrane of the tube showed round-cell infiltration and other slight evidences of inflammation. This the writer regards as important, as it demonstrates the spread of the inflammation by continuity and clearly indicates the necessity of the extirpation of the uterus, tubes and ovaries in cases of

Figure 3.
Case 2.—Endometrium.

polymorphonuclear leucocytes, commonly called pus-corpuscles, and epithelial cells, the latter more numerous in the vicinity of the mouths of the glands. The glands were greatly diminished in number and there were very few which do not show some pathologic changes.

In some, the epithelium was exfoliated and the lumen of the gland filled with a granular mass containing many of these epithelial cells, and often "pus corpuscles." Others appeared to be disintegrating and appeared merely as little islands or masses of epithelial cells buried in the mucosa. Still others deep in the mucosa showed a dilated lumen with an epithelium intact. The arteries of the muscularis show the same thickened intima of the first uterus and even more of the hyaline material in their walls.

There was evidently here a very acute inflammatory process or an acute exacerbation of a chronic inflammatory process. There could hardly be a chronic inflammatory process of any long standing without development of connective tissue unduly in the mucosa, though in this specimen the infiltration of the mucosa by small round cells is so dense as to obscure almost entirely the structure of the mucosa, yet any pronounced increase of connective tissue would be apparent as it would to a certain extent limit the infiltration.

Figure 4.
Case 2.—Opening of Fallopian Tube into uterine cavity. Thick section, to show merely patulous condition.

senile endometritis where there are evidences of involvement of the appendages.

The writer will not attempt to discuss the etiology of the lesion in these cases further than to say he believes that gonorrheal infection can be excluded in both. The histories of the cases, together with the pathologic findings, will, I think, justify me in making the following summary and conclusions:

1. The lesion found in both uteri was an acute inflammatory process. It may be properly denominated acute senile endometritis.

2. The characteristic pathologic features of the inflammation are: *a*, A thickened endometrium, the free surface of which is devoid of its epithelial layer; *b*, increased vascularity with peculiar arrangement of small blood-vessels; *c*, small round-celled infiltration; *d*, diminished glandular elements; while a few glands are to be distinctly seen in many of them, their epithelium is desquamating and their lumen filled with granular débris; they may be said to be functionless glands; *e*, degeneration of the coats of the arteries of the muscular layer of the organ; in one specimen (Fig. 2) this degenerative process is distinctly hyaline; *f*, in not one section examined from various parts of the organ could there be found any increase of connective tissue.

3. The mucosa of both the cervix and body is involved in the inflammation, but is most marked in both cases in the body of the uterus.

4. The small round-celled infiltration extends into the upper muscular tissue, though the inflammation is more marked in the mucosa.

5. In both cases one uterine appendage was diseased; in one the ovary was cystic; in the other, one ovary cystic and the Fallopian tube inflamed. In this case there were slight recent peritoneal adhesions.

6. The microscopic appearance in these cases bears but slight resemblance to that found in case of interstitial endometritis.

7. In one case the acute inflammation seems to have developed without any preceding chronic inflammation.

Figure 5.

Case 2.—Cross-section isthmus of Fallopian Tube, showing slight round-celled infiltration of mucosa and some desquamation. Remaining portion of tube perfectly normal.

In the other case the acute attack may have been an acute exacerbation of a chronic inflammation.

8. In one case there was marked retroversion of the uterus, in the other the uterus was in normal position, and in neither case was there marked stenosis of the internal os; yet there was a considerable accumulation of fluid within the uterine cavity.

9. The presence of diseased appendages in both cases, and of pelvic peritonitis (mild) in one, seems to indicate that the inflammation is prone to extend beyond the limits of the uterus, and if such extension is demonstrable by combined examination, an extirpation of the uterus and appendages is probably the best form of treatment.

Iodoform Intoxication.—A 20 per cent. emulsion of iodoform and glycerin was injected into a cold abscess below Poupert's ligament in a male patient 30 years of age. The amount injected was 80, 90 and 100 c.c. at intervals of two to four months. The temperature rose after the third injection, the patient became confused, comatose, and the nose was remarkably swollen and red, while acne pustules appeared on the skin of the face and trunk, and dry, brown scabs obstructed the mouth and nose. Death followed in ten days. The autopsy disclosed total tuberculous caseation of the suprarenals, although there had been no clinical manifestation of this lesion. The communication in *Beiträge z. Klin. Chir.* xxviii, 1, describing this case states that it is the first on record in which iodine acne has been observed from iodoform intoxication.

DISEASE OF THE FALLOPIAN TUBES.

WITH SPECIAL REFERENCE TO SPECIFIC INFECTION.*

J. R. GUTHRIE, M.D.

DURUQUE, IA.

No structure of equal pathological interest and of such vital clinical importance to the medical profession was so long and so persistently neglected as the Fallopian tube. In the older medical literature it received only the briefest mention, and then the reference was most superficial. Little attention was given its anatomy or physiology, while its histology was wholly neglected. Nothing was known of its pathology and no symptoms were attributed to diseases of this important structure. What a marvelous development in pathological interest and clinical importance during recent years! Careful research will disclose the fact that generations ago the clinicians described disease in the Fallopian tube, but through lack of both anatomical and pathological precision their teaching made practically no modification in either the medical thought or practice of the day.

Perhaps the first well-authenticated account of a discussion of the Fallopian tube and its diseases was in the year 1665, and may be well epitomized in this statement "that occlusion of the tube is one of the most fruitful causes of sterility." Nearly a century later, in 1742, Josias Weitbrecht, the great anatomist of St. Petersburg, described the obliteration of the tube in a woman with one child and who was subsequently sterile. In 1766 Astuc gave considerable attention to the Fallopian tube and afforded perhaps the first accurate description of its diseases. He clearly describes hydrosalpinx, pyosalpinx and tubal pregnancy. But, as frequently happens, his teachings seemed for a time to be forgotten, or at least neglected. Nothing new on this subject is found in medical literature for nearly a century. In the writings of that able French observer, Colombat, as translated by Charles D. Meigs in 1845, there is found only a brief mention of this organ, and he practically describes only one disorder of the tube, atresia. Investigation was imperfect, reports contradictory, and knowledge of slow growth.

In 1854, Charles D. Meigs published his brilliant work. He likewise refers to atresia of the tube and to its effect in producing sterility. He speaks of a most remarkable death from what would to-day be diagnosed as a case of pyosalpinx. Acute observer that he was, scholarly and progressive in his field, yet he devoted only five pages to the tube, including its anatomy, physiology, pathology, clinical history and methods of treatment. T. Gaillard Thomas published his work on diseases of women in 1869. He was a bright light in his special field for nearly a generation, and yet he gives only two pages to a discussion of the tube and its diseases. In brief, an examination of older medical works and journals of all kinds, home and foreign, will show absolute want of literature on this important subject.

The medical man of to-day is well informed on all phases of disease of the tube and of the different views of tubal disease, its origin, pathology and methods of treatment. The causes of inflammation are numerous. When the tube is considered as a continuation of the structure of the uterus, the cause of the inflammation is easily understood. Any cause which may produce inflammation of the uterine structure may, by simple extension, cause an inflamed tube. Disease of bacterial

*Presented to the Section on Obstetrics and Diseases of Women, at the Fifty-first Annual Meeting of the American Medical Association, held at Atlantic City, N. J., June 5-8, 1900.