fails to find any such relation, his results are interesting in connection with the "narrow aorta" of Rokitsky and Virchow. Suter has gone thoroughly into the question, and in his tables has taken into account age, sex, and size of the individuals. He has measured the aorta in 2719 cases and tested the elasticity of its wall in 105. He concludes in part as follows:

1. The size of the aorta varies with age and sex, and in those of the same sex with the size of the individual.
2. It is not true that those who die of puerperal fever, typhoid, carcinoma, or phthisis have narrow or wide aortas.
3. The power of contracting when the blood-pressure is removed—i. e., the elasticity of the organ—decreases with age.
4. There is no such thing as the narrow aorta in the sense of Rokitsky and Virchow. The size of the organ in the cadaver does not correspond to its size in life, and measurements from the autopsy chamber cannot be substituted for them.

Case of Fatal Potassium Chlorate Poisoning.—JACOB (*Berliner klinische Wochenschrift*, 1897, No. 27) reports the following case: A seamstress, aged thirty-nine years, was admitted in an almost comatose condition; face was ashy, lips, fingers, and ears cyanotic; tongue blue-black; blue and copper-colored spots on the temples; marked dyspnoea, thready pulse. The patellar tendon reflexes absent. During the four days of the patient's life there was almost complete anuria, from 25 to 50 c.c. being passed daily. The characteristic methæmoglobin bands were plainly seen. Many corpuscles were present at first; later they diminished, but at the end the urine contained methæmoglobin, hæmoglobin, and traces of hæmatin. On the third day there was pain in both hypochondria, without enlargement of the liver or spleen. The patient died suddenly on the fourth day. On entrance the blood-count showed 4,425,000 red corpuscles and 84,000 whites. The second day the reds had sunk to 1,825,000, and the whites sank steadily to about normal before death. On the first day no very marked changes could be seen in the erythrocytes. On the following day normally colored reds were scarcely to be found, and many shadows, in part arranged in rouleaux, were present. There were, too, many fragments of red corpuscles, partly free and partly contained in the corpuscles, in the form of spheres. These forms increased so that finally there were large masses of debris. The white cells were often filled with the disintegrated reds. The stained covers were not so striking; the whites, however, looked swollen and had pale nuclei. The stained reds were so irregularly stained that many seemed to be colorless and to contain granules. Toward the end the preparations looked like poorly fixed covers. The author's finding agrees generally with that of others. He differs, however, from Riess in thinking that the increase in the white cells may be a reaction-leucocytosis. The author claims that potassium chlorate is a dangerous drug and that its use should be restricted, especially in pediatric practice.

Relation of Albumosuria to Fever.—SCHULTESS (*Deutsches Archiv für klinische Medizin*, vol. lviii. Nos. 4 and 5) has endeavored to solve the problem of the relation of albumosuria to fever by finding out how often albumosuria occurs in the urine of patients with fever and how frequent is its occurrence

in those with no fever. He thinks that 75 per cent. of the cases of albumosuria in the literature had fever; and although such figures are only to be received with caution, they point to a connection between the two. In the author's own work urines containing albumin were carefully excluded. After the reaction was determined any mucin or nucleo-albumin was removed by acetic acid; the filtrate was then carefully tested for albumin. If none was found, 25 to 30 c.c. were mixed with five to six volumes of absolute alcohol, stirred, and allowed to stand for twenty-four hours, filtered, the precipitate dissolved in warm water, and the filtrate again tested for nucleo-albumin, and with the biuret reaction. Of fifty-six cases with non-febrile disease, forty-three, or 77 per cent., had no albumose. In the fifty-nine cases with febrile diseases, 90 per cent. had albumose. The acute infectious fevers, as scarlet, diphtheria, typhoid, were especially noteworthy. In these the albumose curve coincided very nearly with that of the fever.

The Excretion of Leprosy Bacilli.—WEBER (*Deutsches Archiv für klinische Medizin*, vol. lviii. Nos. 4 and 5).—The appearance of several sporadic cases of leprosy has lately given a new interest to the question of the manner of the spread of this disease. Experiments to prove the contagious nature of the disease have so far been contradictory. If leprosy is really contagious and caused by Hansen's bacillus, the bacillus must leave the body in such a way that it can be transferred to others. To find in what way the germ does leave the body the author has examined, by the Ziehl-Nielsen method, all the secretions, the blood and tissue-fluids, the skin, scales, and hair. Bacilli were found in blood from the diseased tissue, in vesicles from the diseased tissue, in scales from the skin, in lanugo hairs, in sweat, and in semen.

From the above it is clear that the germs are excreted in various ways. Whether they are in a condition to cause leprosy anew has not yet been definitely shown.

Experiments with the Modern Treatment of Prostatic Hypertrophy.—CASPAR (*Berliner klinische Wochenschrift*, 1897, No. 27) discusses the effects which modern methods of treatment—castration, tying the vasa, and feeding with the gland-substance—have upon prostatic hypertrophy, and has tried to throw some light upon the question by experimenting upon animals. We know nothing about enlarged prostates in animals. The important thing is to bring about a reduction in the size of the gland, whether at the expense of the normal or pathological tissue is of no consequence. For the reason that enlarged human prostates always contain normal tissue, experiments upon normal glands will be of value. One may assume that the prostate bears the same relation to the genitals in men as in the lower animals. Experiments done so far have for the most part shown that castration is followed by atrophy of the prostate gland. Animals for experimental purposes must be chosen with the greatest care, as the structure and size of the prostate vary with age. The author experimented upon twenty-one dogs and rabbits, and found that double castration always caused atrophy of the prostate. Bilateral excision of the vasa deferentia had once a similar, although more delayed, result. In two other cases, however, no atrophy was to be found. The tying of the vas had no effect upon the testis during the four months the animals were

observed. Unilateral castration, unilateral vasectomy, and feeding the gland-substance had no influence whatever upon the prostate. Caspar disagrees with Sakur, who claims that with the loss of the testis-function something is withdrawn from the system that is necessary to the maintenance of the prostate. As agreeing with experimental and clinical facts, he adopts the view of Pezewalski, who holds that the division of the nerves of Cowper which accompany the vasa brings about an atrophy of the prostate, but not of the testis. The author declines to draw any practical conclusions from his experiments, and thinks it is too early for any accurate deductions from the clinical material so far at hand.

Cholecystitis Typhosa.—VON DUNGERN (*Münchener medicinische Wochenschrift*, No. 26, 1897) has been able to find four cases of cholecystitis in which the Eberth bacillus could be shown to be the cause. To these he adds the following: A woman, aged forty-six years, had typhoid fever in 1882; in 1887 she began to have severe attacks of cardialgia with vomiting. The same trouble recurred again the next year, and again in 1889, this time with icterus. No symptoms in 1895, when there was a periostitis of the lower jaw with necrosis. In October, 1896, there was colic, and in the region of the gall-bladder a tumor the size of a child's head, which proved to be an abscess connecting with the gall-bladder. The pus contained typhoid bacilli in pure culture. The blood-serum of the patient was very active, causing a complete Widal reaction in a dilution of 1 to 80, although the patient had had no typhoid for fourteen years. The author thinks it most likely that the bacilli entered the gall-bladder during the typhoid fever fourteen and a half years ago, and remained entirely latent for five years, causing the various attacks since.

It is possible that the primary infection may have been gall-stones, and that the gall-bladder was secondarily infected, forming a "liver typhoid," which did not involve the intestine because the germs were of low virulence and because the patient had an increased resistance due to the previous typhoid.

Etiology of Multiple Sclerosis.—BLUMBEICH and JACOBY (*Deutsche medizinische Wochenschrift*, No. 28, 1897) have studied the histories of the cases of multiple sclerosis in Gerhard's clinic, with especial reference to trauma as an etiological factor in this affection. In all twenty-nine cases were analyzed; twenty-three were men and six women. The fact that several times the affection began in childhood serves to bear out the theory of Ziegler, that multiple sclerosis arises upon the basis of a congenital disposition. Many of the cases had had infectious diseases; in only one, however (influenza), could any connection be made out between it and the onset of the multiple sclerosis. There was but one undoubted syphilitic. Six times there was a possibility of an intoxication, and two were alcoholics. All were carefully questioned as to trauma; eleven had been subjects of injuries. Five times the disease followed promptly upon the injury, in a sixth the multiple sclerosis followed at once upon the trauma; but this latter was due to a sudden dizziness. In the other five cases the trauma occurred a long time before the onset. Although the etiology of multiple sclerosis is not yet clear, the authors

think there can be no doubt that a relation exists between acute infections, intoxications and trauma. The trauma may cause the anatomical changes upon which the disease depends, or it may only occasion the breaking out of an already existing disease; again, it is possible that the trauma only plants the germ of the affection which may require a second influence to give it typical characters. On the other hand, it may be that the relation is the reverse, and that the giddiness and uncertain gait of the disease cause the trauma. A knowledge of the causes of multiple sclerosis will be of value in the way of prophylaxis and for treatment.

Röntgen Rays and Acute Dilatation of the Heart.—SCHOTT (*Deutsche medicinische Wochenschrift*, No. 31, 1897) in 1890 published a series of experiments showing that an acute dilatation of the heart could be caused by severe exercise. He was at that time obliged to depend entirely upon methods of physical diagnosis. Now he publishes a series in which the results of physical examination are corroborated by means of skiagraphs and diagrams from screens. The results showed that after exercise the heart was enlarged in all diameters, in one case as much 3.5 cm. laterally. The right ventricle not only showed the greatest increase in size, but its enlargement came on first and disappeared before that of the left ventricle. A bicycle ride of 5½ km. in twenty-three minutes caused a dilatation of 2 cm. in the right ventricle, and the apex to move outward 1½ cm.; a walk of 1½ km. in thirteen minutes, on the other hand, while causing a sweating as marked as the riding, brought about no increase in the size of the heart.

Anatomy of Gout.—RIEHL (*Wiener klinische Wochenschrift*, No. 34, 1897), opposing the view of Ebslein, says that in gout urate crystals are deposited in unaltered living tissues, and that the assumption that necrosis of the tissues must precede the deposit of the salts must be given up. Whether the tissue increase about the urate deposits is due to mechanical or chemical irritation is not yet clear. Gouty nodules are probably formed about as follows: First, a deposit of crystals in the lymph-spaces and vessels and in the connective tissue itself, due to this inflammation of the surrounding tissue with granulation tissue and giant cells. The author thinks that the failure of others to find urates in the healthy tissues is due in part to their having examined tissues from the cadaver, and in part to faulty methods of hardening.

Stomach Tetany and Autointoxication.—This affection is tolerably rare, about forty cases having been reported, of which not more than eleven are considered pure. The cause of the spasm is in doubt. The view that it is an intoxication has the most adherents, and is the one entertained by the author. If this theory is correct one would expect to find the poison in the stomach-contents and in the urine. GUMPRECHT (*Centralblatt für innere Medizin*, No. 24, 1897) has studied these secretions in a nineteen-year-old workman, who had dilatation from pyloric stenosis. The patient had typical tetany spasm lasting several hours and slowly disappearing through several days. The author reviews the literature of the subject and gives in detail the methods used in his own work. In conclusion, he says: "Neither