Pneumothorax has occurred five times, on one occasion accompanied by pus in the pleural cavity, but this was complicated with chronic tuberculosis. Two cases in the service were diagnosed as pneumothorax and were demonstrated as such to various members of the medical staff, because of the remarkable coin sound that was present. Both of these cases came to necropsy, and it was shown that pneumothorax did not exist and that the coin sound was due to a distended colon. The coin sound in both of these cases was present up to and over the second interspace of the left side and throughout the whole of the axillary space.

Otitis media has occurred frequently, and in many cases has been followed or accompanied by mastoiditis. Strange as it may seem, mastoiditis has occurred and been found postmortem without having given any objective or subjective signs during life.

PERCENTAGE OF INCIDENCE OF PNEUMONIA TO THE TOTAL
NUMBER OF MEN IN THE CANTONMENT

Month	Per Cent.
October, 1917	0.31
November, 1917	0.78
December, 1917	1.10
January, 1918	0.73
February, 1918	0.34
March, 1918	0.52
Average	0.65

The percentage of the incidence of pneumonia to the total number of men in the cantonment for the months from October to March is rather interesting, and is given in the accompanying table. It is hard to say definitely what is the cause of the high incidence of pneumonia at this camp. For the first four months, the physical condition of the men was very poor; a tremendous amount of dust was present; many of the men had the habit of going to bed in their wet underclothes without properly covering themselves, and sudden changes of temperature took place. Undoubtedly by this exposure they were made more susceptible to respiratory infections. The large number of men suffering from measles predisposed them to infections of the respiratory tract. In addition, practically in every one of these men there were enlarged tonsils. The teeth and gums also were in a very bad condition. The factors mentioned above, namely, measles, the physical condition of the men, the large amount of dust, and carelessness in barracks, together with the exposure and fatigue, must all be considered as factors in the incidence of the different pneumonias.

SUMMARY

Probably the three most potent factors in the predisposition to pneumonia in this camp were:

1. The large number of men suffering from measles.
2. The large amount of dust.
3. Carelessness of the men in their barracks, giving rise to what were at first minor respiratory infections.

These three factors all give rise to coryza, laryngitis, tracheitis or bronchitis and so cause conditions predisposing to infections that ultimately may result in pneumonia.

Importance of a Sanitary Code.—As a basis for the activities of the health authorities, a sanitary code or set of rules and ordinances or regulations should be promulgated. These regulations should be prepared with extreme care and permit of intelligent application and enforcement.—P. A. Surg. Paul Preble, *Public Health Reports*, March 2, 1917.

EPIDEMIC OF STREPTOCOCCUS PNEUMONIA AND EMPYEMA AT CAMP DODGE, IOWA *

JOSEPH L. MILLER, M.D.

Major, M. R. C., U. S. Army

AND

FRANK B. LUSK, M.D.

Captain, M. R. C., U. S. Army

CAMP DODGE, IOWA

From Sept. 20, 1917, until May 10, 1918, 675 patients, whose cases were diagnosed as lobar pneumonia, have entered the base hospital at Camp Dodge, Iowa. This does not include a limited number of cases diagnosed clinically as bronchopneumonia. At necropsy, however, not infrequently it was found that, in reality, we were dealing with a confluent bronchopneumonia which was localized chiefly or entirely in a single lobe. Inasmuch, however, as these cases, if necropsy had not been performed, would have been placed on record as lobar pneumonia, they have been so considered in this discussion.

Although measles was quite prevalent throughout the entire winter, pneumonia as a complication was very infrequent; only since about May 1 has pneumonia as a complication of measles reached any considerable number, and this period is not connected with the epidemic of streptococcus pneumonia, which at the time mentioned was on the wane.

The ordinary clinical lobar pneumonia, due to the pneumococcus, prevailed until about March 20; then, abruptly the streptococcus type predominated, with a very great increase in the incidence of the disease. From September 20 until March 20, which marks the period of ordinary lobar pneumonia, 276 cases were treated; from March 20 to May 10, 400 patients with pneumonia entered the hospital.

A very mild type of pneumonia prevailed during the early autumn, with a mortality in the first hundred of 7 per cent. The onset was usually severe and evidence of intense toxemia prevailed, but the course of the disease was short. In the first eighty-eight cases terminating by crisis, the duration in seven was 2 days; eleven, 3 days; thirteen, 4 days; thirteen, 5 days; eighteen, 6 days; nine, 7 days; ten, 8 days; two, 9 days, and five, 10 days or more. In this series of 276 cases, empyema was present in thirty-one, or 11.2 per cent. Even in these early cases, however, there was a marked tendency to multiple pus foci and the mortality from empyema was high. In those of this series where the type of pneumococcus was determined, Type I was found in 22.8 per cent.; Type II, typical and atypical combined, 46.8 per cent.; Type III, 7.6 per cent., and Type IV, 22.8 per cent.

The colored troops were much more susceptible than the white, 116 being colored and 159 white, and one American Indian. The colored troops did not enter camp until we had received forty white soldiers with pneumonia. Allowing for this, it is seen that there was about an equal number of the two races, although the colored troops made up only about one sixth of the total strength of the camp. The mortality among the colored troops was also somewhat higher—12.9 per cent., with 9 per cent. among the white troops,

* Read before the Section on Practice of Medicine at the Sixty-Ninth Annual Session of the American Medical Association, Chicago, June, 1918.

the total average morality being 11.2 per cent. Empyema occurred in 8.6 per cent. of the cases of pneumonia in the colored troops; in the white troops, 13 per cent. It was much more fatal among the former, 70 per cent. dying as compared to 57 per cent. among the latter. This is in striking contrast to the results observed in the streptococcus empyemas. The mortality in the thirty-one empyemas was 61.3 per cent., which is accounted for by the marked tendency to multilocular pus collections. This feature, so characteristic of the streptococcus empyemas, also prevailed in our early cases of pneumococcus pneumonia, although in the majority of these, streptococci were found in the pleural exudate, either alone or in conjunction with the pneumococcus.

The epidemic of streptococcus pneumonia appeared suddenly between March 18 and 20, continued with great severity for six weeks, then gradually became less intense, still continuing, however, May 10 at the rate of four or five cases daily. The virulence of the epidemic, however, became less marked after the first three weeks, although the number of new cases remained high.

It was immediately recognized by the ward surgeons that we were dealing with a different type of pneumonia. Evidence of severe intoxication appeared very early; empyema became very frequent and developed extremely early, two patients entering the hospital with pleural exudate who had been drilling the day previous. While the involvement in the lung maintained a lobar type, clinical evidence of complete consolidation was far from constant. Dulness with suppressed breathing and subcrepitant râles, but inconstant or localized bronchial breathing were the usual findings; rusty sputum, provided there was expectoration, was the rule. Early roentgenoscopy showed that the infiltration was lobar in character. The development of an exudate was often exceedingly difficult to determine by the ordinary physical findings, aided by the roentgen ray, and it became necessary to resort to frequent exploratory aspirations. These were often repeated several times before the fluid could be located. Early this exudate was only moderately turbid, contained numerous polynuclear leukocytes and showed on smear short chain streptococci. Gradually the fluid became definitely purulent.

The bacteriologic findings in ninety-five of these exudates showed pure streptococci in eighty-eight, all being hemolytic. Pneumococci combined with streptococci were found in three: two of Type II and one of Type I. Pneumococci without streptococci, but often combined with other bacteria, were found in four cases: one of Type I, two of Type II and one of Type IV.

At the outbreak of the epidemic, a special effort was made in taking the history to determine predisposing causes or evidence of any infection antedating the onset. It might be stated that the month of March was exceedingly dry, and clouds of dust hung over the cantonment. Either as a result of this, or independently, a mild tracheitis was very prevalent. Wherever a large group of soldiers congregated, the prevalence of coughing was very noticeable. It was, therefore, not surprising that many patients gave a preliminary history of mild sore throat, often of several days' duration, but more frequently a preliminary history of cough. History of great fatigue, or chilling was noted occasionally, but not with significant

frequency. Not infrequently from apparently good health, the onset was ushered in with a chill. While our patients came from all parts of the camp, there were certain organizations where the disease was more prevalent. Of the first fifteen cases, ten came from a colored regiment, which had been in Camp several months, and throughout the course of the epidemic this organization furnished the highest percentage. No one company in any organization furnished an unusual number of cases. The depot brigade, which was made up largely of new arrivals, suffered the least. At the time the epidemic began, the camp was not crowded and the number in each barrack was much less than it had been in the winter. The ventilation in the barracks, on account of the mild weather during March, was probably much better than it had been in the two preceding months.

As an aid to early diagnosis and exact localization of the involvement, the roentgen ray was found to be of considerable value. It was of much less assistance in differentiating between consolidated lung and pleural exudate, although when repeated frequently and compared with the ward findings, it proved to be of great value. In the beginning, an order was issued to all ward surgeons in the Medical Service that every pneumonia suspect should be roentgenographed within twenty-four hours, and in addition, every case diagnosed as bronchitis, where the temperature remained above 101, for twenty-four hours. By this means all pneumonias were detected early, and it furthermore stimulated the ward surgeon in that a more careful physical examination be made, as he knew that his findings would be checked up with the roentgen ray. Furthermore the roentgenoscopy was repeated every fifth day until the patient's temperature had been normal for one week. This was of considerable assistance in detecting empyema. The roentgen ray also gave us accurate data regarding the incidence of the lobe first involved. In 596 cases of both types, the first involvement occurred in the lower left lobe in 253; lower right, 252; upper right, 45; upper left, 23; middle, 12; entire right lung, 5; entire left lung, 1, and in both lungs, 5.

Complication with pus formation was very frequent. While empyema developed in 11 per cent. of the pneumonias previous to the epidemic, it appeared in 34.8 per cent. of the cases during the epidemic. One case of suppurative peritonitis was found in the 276 cases of pneumonia preceding the epidemic and five cases in the first 364 streptococcus cases. In three of these it was associated with suppurative pericarditis and in all cases with empyema. It was interesting that four of these cases developed within a few days of each other. Suppurative pericarditis was found in fifteen of the 142 cases of empyema, or 10.5 per cent. It was only found in association with empyema, and in only one case was it found in a colored soldier. Rectus abscess, which was more or less frequent in similar epidemics in other camps was not present in the first 364 cases, but one abscess developed in a more recent case. The colored soldiers developed empyemas in 20 per cent. of the pneumonias and the white soldiers in 45 per cent. The empyema mortality was 44 per cent. in the colored and 64.8 per cent. in the white soldiers. This lessened tendency to empyema and suppurative pericarditis in the colored troops was striking. The deaths from uncomplicated streptococcus pneumonia in the colored was 19.97 per cent.,

and 10.7 per cent. in the white. These figures show that the colored soldiers show a much higher mortality from uncomplicated pneumonia than the white soldiers, but a lower mortality from empyema.

The mortality in 142 streptococcus empyemas was 60.4 per cent. This is considerably higher than the average in other camps. There is no apparent explanation of this unusually high mortality, as the method of treatment was similar to that carried out elsewhere. It is true that in only a moderate percentage of these cases, death was due not to empyema itself, but rather to severe infection, the majority of the patients dying with merely a turbid fluid, the time being too short for it to become thick and creamy. The mortality in 346 streptococcus pneumonias was 32.5 per cent. as compared with a mortality of 11 per cent. in the previous pneumococcus type.

Arthritis as a complication appeared in only six cases. Suppurative otitis media was quite common. Erysipelas developed as a complication in seven cases. These were facial except in one case, where it developed at the site of the exploratory puncture. It was also interesting that four out of six cases of scarlet fever in nurses during this period occurred in nurses working in the pneumonia wards, although all wore masks and gowns, and were instructed regarding the disinfection of the hands.

The treatment of the pneumonias consisted in the ordinary hygienic measures. All patients were placed in special pneumonia wards, each bed separated by a sheet suspended from a wire and 1,000 cubic feet of air space to a patient. No attempt was made to reduce temperature by bathing. Every patient received morphin for pain or to control restlessness. Tincture of digitalis in 1 c.c. doses was given every three hours, day and night, from the time the patient entered the hospital until the crisis, unless evidence of its toxic effects developed, and this appeared in only a very few cases. It was decided during the autumn that we would use this cantonment as a control and not use the serum. It is interesting to note that in the 276 cases of pneumococcus pneumonia, only two patients died with Type I infection. The largest number of fatalities occurred in the Type II group.

In the early days of the streptococcus pneumonia, where aspiration showed a turbid fluid containing streptococci, immediate drainage was resorted to. As the mortality was extremely high, this was discontinued and the fluid became definitely purulent in from seven to ten days. The patients were then operated on by rib resection and drainage. The operated empyemas were irrigated and dressed by the surgical service, but held on the medical service in a special ward, the medical department being on the lookout for new accumulations of pus or other complications. When early rib resection was performed, the patients frequently returned from the operating room in a state of shock, from which they did not recover. Sixty-five cases were treated by immediate operation, with a mortality of 53.9 per cent.; fifty-five were aspirated, the operation being postponed as has been stated; 32.7 per cent. of these terminated fatally; four recovered by aspiration alone. The remaining number of the 142 empyemas, namely, twenty-two, reached an early fatal termination so rapidly that either the pus was not detected until necropsy or the patient's condition was such that aspiration was considered inadvisable.

It would appear that the streptococcus empyemas might be divided into three groups: (1) those who die early, no matter what form of treatment is instituted, from acute toxemia; (2) those with multiple pus foci, difficult to detect, because of the inability to locate and drain all foci; these all die, and (3) those who usually recover, either from early operation or aspiration followed by operation; here are included those with moderate toxemia and those with localized pus accessible to drainage.

PATHOLOGY OF THE EPIDEMIC STREPTOCOCCAL BRONCHOPNEUMONIA IN THE ARMY CAMPS*

W. G. MacCALLUM, M.D.

Contract Surgeon, U. S. Army

BALTIMORE

In a previous paper¹ a peculiar form of bronchopneumonia, caused by a hemolytic streptococcus and prevalent in some of the Army camps, was described. This was based on observations made during February and March at Fort Sam Houston, Texas. Since that time I have been able to study during May a quite similar epidemic at Camp Dodge, Iowa,² where, however, the surrounding conditions, climate, housing of troops, and organization and construction of the base hospital were very different. Another striking difference lay in the great prevalence of measles in the Texas camp, whereas at Camp Dodge measles has not been very common. In the Texas camp we were greatly impressed with the importance of measles as a disease predisposing to secondary invasion of the streptococcus, although we realized the possibility that other diseases, such as scarlet fever or lobar pneumonia, might act as predisposing causes, and that the streptococcus might sometimes be the primary invader.

At Camp Dodge there were only sixty or seventy cases of measles in the wards, and the pneumonia appears to have arisen quite independently of it. In this camp, therefore, it was the prevailing opinion of the medical officers that measles played little part in the production of the epidemic. Of eighty cases studied clinically and bacteriologically by us at this camp, there were only fourteen in which there had been a recent attack of measles. Even this considerable percentage was the result of a small epidemic of postmeasles pneumonia which appeared during our stay. It was very severe among these men and several of them died, so that among the seventeen cases of interstitial bronchopneumonia that we studied at necropsy there were nine following measles. During the earlier part of the epidemic, however, there were, I am told, very few of such cases. In the Texas epidemic we found among twenty cases of bronchopneumonia at necropsy fifteen with a recent history of measles.

We received the general impression that in a community in which measles was very prevalent the hemolytic streptococcus was able to gain a foothold and to

* Read before the Section on Practice of Medicine at the Sixty-Ninth Annual Session of the American Medical Association, Chicago, June, 1918.

1. Cole, Rufus, and MacCallum, W. G.: Pneumonia at a Base Hospital, *THE JOURNAL A. M. A.*, April 20, 1918, p. 1146.

2. This study was made with the assistance of Capt. A. W. Sellards and Lieuts. H. M. Thomas, Jr., and A. B. Lyons.