inject 5 minims of this solution, increasing the dose minim by minim every second or third day until improvement or a constitutional effect of the mercury occurs.

SOME OBSERVATIONS ON THE TREATMENT OF TABES DORSALIS.

BY DANIEL R. BROWER, A.M., M.D.

Professor of Mental Diseases and Therapeutics, Rush Medical College;
Professor of Nervous Diseases Woman's Medical College (Northwestern University); Professor of Mental and Nervous Diseases, Post-Graduate Medical School.
CHICAGO, ILL.

The treatment of any disease, as a matter of course, is largely based upon its pathogenesis; and while we may safely assume that at least 90 per cent. of the cases of tabes dorsalis have an antecedent syphilitic history, yet it is not a true syphilitic disease, for in the remaining percentage of cases other etiologic factors are at work, among which are trauma, the acute infections, alcoholism, and the auto-intoxications, so that we may call the disease rather a degenerative sequela of these various processes, and we know not whether the disease begins in the neurons, in the connective tissue, or in the blood vessels. But let it begin where it may, the resultant is a sclerosis terminating in hypertrophy of the connective tissue and a destruction of the sensory neurons of the spinal cord. I am of the opinion that with many a too gloomy prognosis is made. In my experience the arrest of the disease is not infrequent, when the treatment is commenced in the pre-ataxic stage, and considerable improvement is possible even when the second and third stages of the disease have been reached prior to its commencement. The disease does not always interfere with the very successful use of the inherent powers of the patients. I have one patient who, since the pre-ataxic stage began about twenty years ago, has carried on a large manufacturing concern, employing at times several hundred men, and has accumulated a large fortune. I have another patient who, since the disease began about fifteen years ago, has attained eminence as a lawyer and as a politician, and has been the Governor of a great State, and is still in the active pursuit of his profession.

Treatment.—The first consideration in the treatment is the climate, and whenever it is practicable, as it often is in a great country like the United States, a warm, dry, equable climate of low level or moderate altitude should be selected as the place of permanent residence for the patient.

If the case is not advancing rapidly, an ocean voyage, giving the maximum amount of fresh air with a minimum amount of exertion, is often beneficial. I am in the habit of recommending to my Chicago patients, when possible, to spend two or three of the spring months in the Ozark Mountains of Missouri, in the mountains of North Carolina, or in a similar moderate altitude; or at Atlanta, Ga., or Los Angeles, Cal., or in similar low level climates, thus avoiding the cold and dampness and exceedingly variable climates of this season of the year in this locality.

The next important consideration is rest. When the disease is advancing rapidly, absolute rest in bed, with daily massage and the faradic exercise of the muscles is indicated, and this may often be continued with advantage for six or eight weeks, and then

patients gradually resume their ordinary avocations; and in every case I think a portion of each day should be spent in the recumbent position. I advise business men to provide themselves with a lounge in their office, and spend at least one hour in the recumbent position, and in this position they can transact much business, and at the same time give rest to the spinal cord. Excessive mental work and physical fatigue should be avoided; sexual excesses are especially harmful and seem to contribute to the production of optic nerve atrophy; whatever exercise the patient takes should always stop short of fatigue. Traumas being of serious consequence, especially concussion of the spine, should be avoided with great care. The training of the muscles in the work of co-ordination, as proposed by Fraenkel, I endorse, and advise various exercises, such as he proposed, for an hour or an hour and a half daily, and it is surprising what amount of power of co-ordination will be regained by the judicious following of this teaching.

The gastro-intestinal tract demands special attention; lavage and bowel irrigation are often indicated. Gastric crises are provoked by indigestion and colonic impaction. Food, therefore, should be easily digested, and constipation should be avoided. A change of life from one of activity to one of inactivity, forced upon the patient by the disease, tends to develop gout in those who are predisposed to it, and this should be constantly borne in mind and the diet regulated accordingly. Excesses in alcoholics and in smoking should be avoided. Electricity, especially the static form, from a machine of high potential, by insulation and by heavy sparks from the spine and lower extremities, is usually of very much service. Faradization of the skin by the wire brush electrode is also beneficial. These two forms of electricity, in my judgment, are very far superior to galvanization. When the bladder or its sphincter is weak faradization of this organ is often of service.

Suspension, first brought to our notice by the Russian physician, Motschutkowski, is of service when the disease is advancing after a more or less lengthened stationary period. I am sure that it is of service, although the practice is being abandoned by a great many. Stretching the sciatic nerves by manipulation, in the same class of cases gives results somewhat similar to suspension. Cold or tepid baths at a temperature not to exceed 96 degrees F., especially with jet or shower accompaniments, are useful. A hot bath, in my judgment, is often very injurious.

In those cases of tabes dorsalis in which the symptoms have developed very rapidly and the syphilitic infection is recent, vigorous antisiphilitic treatment is beneficial, and it should consist in the largest possible doses of the iodid of potassium together with the hypodermatic use of mercury. But in those cases of locomotor ataxia where the development has been slow, and where a long interval has elapsed since the primary symptoms occurred, I am sure that this form of medication often does great harm. An alternative on which I place the greatest reliance in the chlorid of gold and sodium. I think that this drug has some power in arresting the progression of connective tissue hypertrophy in the spinal cord, as well as in the liver and kidneys. I am in the habit of giving this drug in the tenth of a grain doses (.006 gram), three times a day, and usually combine it with the resin of guaiac (.18 grams). Chlorid of gold and sodium is a very unstable drug and great care should

be exercised in its preparation by the pharmacist and in its administration by the physician. It is my practice to have the drug rubbed up dry with the pulverized resin of guaiac placed in capsules, and given one hour before meals; or, in more urgent cases, I have it prepared freshly in solution and use it hypodermatically, thus hoping to avoid by the one method or the other the destruction of the drug in the stomach. This drug being a tonic, as well as an alterative, it is indicated in the several stages of the disease and may be continued with advantage for months at a time.

The next drug upon which I place reliance is a preparation of phosphorus, and for this purpose prefer the phosphid of zinc. This I give in the eighth (.008 gram) or tenth (.006 gram) of a grain doses, three times a day after meals. As a tonic and alterative, in alternation with the above I use arsenic, preferring the arsenate of sodium in about the twelfth of a grain (.0054 gram) doses, to any other of the preparations of arsenic.

When the disease has taken upon itself a rapid developmental state, full doses of ergot with the rest before indicated will sometimes stop the rapid progression of the disease. For the pains I have found that extract of cannabis indica, injections of cocain, and acetanilid alone, or in combination, are the most successful. I have seen no especially beneficial results from the use of nitrate of silver, aluminum chlorid or of mercury in ordinary cases, and my experience is emphatically against the use of strychnia in this disease. I am very confident that I have seen more than one case very greatly injured by the use of even ordinary doses of strychnia.

GUMMATA OF THE HEART IN A CASE OF CONGENITAL SYPHILIS.

FROM THE PATHOLOGICAL LABORATORY OF RUSH MEDICAL COLLEGE.

Read before the Chicago Pathological Society, Jan. 11. 1897.

BY E. R. LeCOUNT, M.D.
DEMONSTRATOR OF PATHOLOGY.
CHICAGO, ILL.

The following case is of interest as an instance of a lesion observed very infrequently, viz: gummata of the heart as a feature of congenital syphilis.

A child born at term, of young parents, died almost immediately following birth and was sent to the laboratory of Rush Medical College by Dr. A. B. Strong, of Chicago, in whose practice the case occurred and to whom I am indebted for the opportunity of making the examination.

Autopsy Abstract.—Weight 2,500 gm., length 50 cm.; the umbilical cord showed a recent ligature. The external surface of the body was the seat of a vesicular and pustular inflammation, most marked over the face, hands, feet and buttocks. The contents of these eruptions, although generally clear, were occasionally seropurulent and less often of soft, cheese-like necrotic material. The peritoneum was normal and its cavity empty. The thymus weighed 7.5 gm., and possessed several softened areas containing gray viscid pus.¹ The thyroid, submaxillary and parotid glands were normal in appearance except for an excess of blood. The pleural cavities were empty. The lungs contained numerous firm, white nodules, some of them

as large as cherries. On section these presented a smooth surface and appeared of equal consistency throughout. The lung tissue was non-crepitant or very slightly crepitant, fairly firm and oozed dark blood. The pleura over the left lower lobe was roughened by an easily removable fibrinous exudate. The liver was engorged with blood and extended to the level of the umbilicus. The spleen was slightly enlarged and congested; the pancreas was negative; the kidneys, stomach, intestines, lymphatic apparatus and the central nervous system all appeared unchanged.

The lines of ossification in both long and flat bones were irregular, wider than normal, and hemorrhagic.

The heart.—On the anterior surface of the heart midway between the base and the apex, involving that part of the ventricular wall which faces toward the left, and also overlapping the ventricular septum, was a white area, quite circular in form and measuring approximately 1 cm. in diameter. The edge of this white area fused gradually with the adjacent heart tissue, and on section appeared less firm than the unchanged myocardium (Fig. 1.) It was seen to involve the entire thickness of the ventricular wall, spreading out more as it approached the pericardium, which was unchanged except as to color. The change in the muscle also extended into the columnæ carneæ of the left ventricle. On the posterior or diaphragmatic surface, near the right border of the heart, were situated three smaller but similar areas.

FIG. 1.—Gumma near the interventricular septum.

The heart cavities contained considerable blood partially coagulated. The valves and orifices showed no changes.

Anatomic diagnosis.—Congenital syphilis, syphilitic interstitial pneumonia (nodular); syphilitic osteochondritis; gummata of the heart; pustular syphilides of the skin; multiple abscesses of the thymus; hyperemia of the liver and spleen.

Microscopic examination.—The presence of syphilis was confirmed by the examination of the viscera, characteristic changes being found in the liver, lungs, spleen and lines of epiphyseal ossification. In sections made through the large lesion on the anterior surface of the heart, the following changes were found: Between the heart muscle fibers throughout the area, which to the naked eye appeared white, there existed a marked infiltration of round or partially fibrillated cells, with faintly staining vesicular nuclei. Although the change prevailed throughout it was most marked around the blood vessels (small arterioles). The change here began immediately outside the media, where distorted cells and cells of peculiar shapes, with faintly stained nuclei occurred. These were evidently formative cells, and resulted from proliferative

¹ The mixed infection which had resulted in this case was investigated by Mr. H. G. Wells and reported by him to the Chicago Pathological Society (Transactions Chicago Path. Soc., Vol. II, 1897).