ance sense. The vasomotor derangement is supposed to be due to the loss of ovarian secretion during the menopause, this loss resulting in lowered metabolism or changes in the functions of the thyroid, adrenals, and pituitary body which, in combination, induce vasomotor disturbances. Vertigo at the menopause comes on at irregular intervals—attacks occurring sometimes days and weeks apart, or again, several times daily. In some, it comes on without apparent cause, in others only after unusual movements or postures. In most of the author's patients the vertigo was of the objective type. Nearly always there coexisted such disturbances as hot and cold flashes, cold perspiration, palpitation, blurred vision, flickering before the eyes, headache, nausea, tinnitus, etc. Before undertaking treatment, such other causes of vertigo as lesions of the internal ear or of any other part of the balance mechanism, cardiovascular, renal and ocular diseases, and especially muscular imbalance of the eyes, must be excluded. In the true climacteric vertigo, Sanes strives to favor general nutrition and elimination and administers an ovarian preparation each dose of which represents a grain of fresh ovarian substance. The dose used is five grains, two to four times a day. When this dose failed, better results were, in a few cases, obtained from larger amounts. The records showed about thirty-seven per cent. of cases improved, and about twenty-five per cent. completely controlled. The relief from vertigo either preceded, followed, or accompanied the relief from the other menopause symptoms. The drug was used for a month or for many months and in one case for three years before final cessation of the symptoms. In most of the prolonged cases, the vertigo would recur or become worse upon temporary discontinuance of the remedy. If it be true that, while the ovaries are still functioning somewhat, ovarian therapy not only supplies a substitute for lacking ovarian secretion but also stimulates the ovaries to increased secretion; the earlier in the menopause the ovarian substance is given, the better should be the results.

Jordan, H. E. A CASE OF FALSE HERMAPHRODITISM. [*Anat. Rec.*, Aug., 1918, 15, No. 1.]

Rudimentary phallus, vestigial scrotum, cryptorchism, prominent mammary glands, feminine hips, limbs and hands, polydactyly, left handedness, club-foot, adiposis and feeble-mindedness are reported in a male false hermaphrodite. Hyalinization of the walls of the convoluted tubules of the testicles was shown by microscopical examination, the patent tubules being lined with a variable epithelium, mainly of Sertoli-cell origin. Hyalinization and fibrosis of the intertubular stroma, with only scattered interstitial cells, which are about half normal size, with normally distributed content of lipoid granules collected into large groups. This case is marked by a great numerical decrease and change in the interstitial cells and a lack of sexual desire, agreeing with the

supposition that male erotic desire depends on the internal secretion of the interstitial testicular cells.

II. SENSORI-MOTOR NEUROLOGY

2. CRANIAL NERVES.

Dunn, J. SOME PHASES OF THE VESTIBULAR NERVE PROBLEM. [Arch. Ophthal., 1918, XLVII, 354.]

Holding head erect and turning it to the right causes the eyes to move to the left and an excess of movement in the endolymph with body and extremity adjustments. This triple coördination is mediated by the cerebellum, working so as to accomplish any voluntary movement, but not originating movement. The cerebellum both initiates movements and receives stimuli from muscles, joints, etc., involved in the movement, and enabling the movement to be continued or posture to be maintained. The coördination is independent of the will, and, after the movement is initiated, beyond its control. Connections for this purpose exist between the cristæ of the horizontal canals and both the cerebellar cortical centers and the vagic ocular centers. To permit other movements of the head a similar connection has arisen between other canals and the cerebellum and optic centers. Those centers alone whose hyperexcitation produce body falling and past-pointing of the type made manifest by irritation of the cristæ respond to vestibular peripheral irritation, others being directly associated with the cerebral motor cortical centers.

The primal eye being a part of the vegetative system and was under the control of the vagus centers, its development has also come under the control of the will, so that excitation of the vestibular terminals results in impulses to both the vegetative ocular nuclei and the voluntary nuclei. The nystagmus therefore has two elements. The one results from a hyperexcitation of the primary stem centers rendering possible a nystagmus, even in deep unconsciousness. The other results from overexcitation of the centers which have developed from voluntary control over the external ocular muscles. The same rules which determine the direction in past-pointing and body falling obtain here too. When the eyes are fixed and the head is moved the eyes move in the opposite direction with a slow, pendulum-like movement, but when the head is fixed and the eyes move, the movement is jerky. The former movement is under the control of the voluntary nuclei. In nystagmus the slow movement represents the response to irritation of the vagi nuclei. The rapid movement represents a past-pointing of the eye and necessitates an element of the will for its performance.

Movements of the eyeball, whether normal or pathologic—*e. g.*, albinism—do not give sensation of the movement of external objects, but vestibular nystagmus always does. Two cases of vertigo are described and analyzed, and lead to several questions: What are the functions of the