SURFACE COLOR.

BY R. W. WOOD.

SO many misleading statements appear in the text-books on optics regarding the phenomenon of surface color, exhibited by certain strongly absorbing media, that it may be well to point out certain facts which are not always brought out as clearly as they should be. We frequently find the statement that these substances reflect most strongly those colors which are absent in the transmitted light, or which, as we more frequently say, are most strongly absorbed. One of the more recent of the advanced text-books refers the phenomenon to total reflection, the explanation being that for waves of certain wave-lengths the refractive index is less than one, and that therefore at a certain angle there will be total reflection for these waves, and consequent surface color. As a matter of fact the dispersion curve runs below unity only in the case of a very few substances, consequently total reflection is wholly inadequate as an explanation. Moreover the selective reflection occurs at normal incidence.

In the first place the statement that the most strongly reflected light corresponds to the light absorbed is erroneous. Take cyanine as an example. The center of the absorption band is very near the D lines, but the surface color is not yellow, but a deep plum color, not so very different in hue from that of the transmitted light. If we examine the spectrum of the reflected light we find a very dark band in the green, the center being not far from wave-length .0005. The distribution of intensity in the rest of the spectrum is not very different from what it would be in the case of reflection from glass, which shows that the peculiar color of the dye is not so much due to a very powerful reflection of certain waves, as it is to its almost complete refusal to reflect a certain region of the spectrum. If we remember that the reflecting power depends on the refractive index

as well as on the coefficient of absorption, we can easily explain this behavior.

The intensity of the light reflected at normal incidence by a transparent medium is given by the formula

$$R = \frac{(n - 1)^2}{(n + 1)^2}$$

As n , the refractive index, varies with the wave-length, the reflecting power will be different in different parts of the spectrum. The formula calls for a stronger reflecting power in the violet than in the red, the variation being greatest for substances with high dispersion and low mean refractive index. An experiment has recently been described by B. Walter to illustrate this type of selective reflection. Using the method of repeated reflections originated by Rubens and Nichols for the isolation of long heat waves, a remnant of blue violet light is obtained. The medium used by Walter was oil of cassia, above the surface of which a silver mirror was mounted.

Fig. 1.

The sun's image after five reflections from the surface of the oil (the light being thrown back and forth between the surfaces as shown in Fig. 1), was found to be colored a deep blue. Though the formula calls for just such an effect, I have failed absolutely to obtain

a trace of it though I employed six reflections, or one more than Walter. The image was very faint and of a neutral gray color with no suggestion of blue.

In the case of absorbing media the refractive index varies over a much wider range than it does in case of transparent substances, consequently greater variation in reflecting power with changing wave-length is to be expected. Moreover the absorption influences the reflecting power directly, the formula for media of this nature being

$$R = \frac{n^2 + n^2x^2 + 1 - 2n}{n^2 + n^2x^2 + 1 + 2n}$$

In the case of a substance such as Hoffman's violet, the refractive index of which is unity for a certain wave-length, the amount of

light of this particular color reflected at normal incidence will be $\frac{x^2}{4 + x^2}$. It is clear that this will increase as x increases, and will equal zero if x is equal to zero, as it is in the case of transparent media.

Taking the values of n and x for cyanine obtained by Pflüger in the different parts of the spectrum, we can calculate the percentages reflected in the cases of the different colors. The curve shown in

Fig. 2.

Fig. 2 was obtained in this way; it shows very clearly the cause of the peculiar surface color of the dye. Starting at the red end of the spectrum we find that the intensity of the reflected light increases with decreasing wave-length, due to the joint increase in n and x . The increase continues for some distance after n begins to decrease, owing to the continued increase of x . After reaching a maximum at $\lambda = .00059$ it turns and descends rapidly attaining its minimum at $\lambda = .0005$, where we have very small values of both n and x , and consequently very small reflecting power. Beyond this point the rise is due almost entirely to the further increase in n , since x is practically equal to zero in this portion of the spectrum.

Scarcely two per cent. of the incident light is reflected at $\lambda = .0005$ causing the dark band seen in the spectrum of the reflected light, alluded to at the beginning of the paper.

An excellent way of showing the variable reflecting power of a film of cyanine is to compare it with glass, in different parts of the

spectrum. A little of the melted dye is pressed between two plates of hot glass, which are separated when cold. A spot is selected where the film has a good optical surface, and this spot is left on the glass the rest being cleaned off. By holding the plate in the spectrum formed by a prism or grating the reflecting power of the two surfaces can be studied. In some parts of the spectrum the cyanine reflects more strongly than glass, in other regions the reverse is true, while at wave-length .0005 the cyanine refuses to reflect to such a degree that the film appears as a black spot on the blue field reflected by the glass.

It is interesting to note the difference in the surface color of the dye when the reflection takes place at the surface in contact with the glass. A very convenient way of showing the yellowish-green color in this case is to press out a film of the molten dye on one surface of a prism of 8 or 10 degrees angle.¹ In this way the light reflected from the dye can be obtained uncontaminated with the light reflected from the first glass surface. The method is analogous to that employed by Lippmann in mounting his color photographs. The calculation of the curve of reflected intensities under these conditions makes a good exercise for the student.

¹ A suitable prism can be made in half an hour by grinding down a piece of thick plate window glass. A strip of thick glass cemented along one edge will be all that is necessary to make the glass take the required form. Grind on a piece of glass with very coarse emery at first, then use finer grades, polishing with rouge at the end. Small scratches do no harm, and a high polish is not necessary.