Epoch	Position	Nights	Dist.	Nights	Mean power	Magn.	Telescope in use.
1839,66	27°42	2	1"5±	1	320	2½, 8¾	Mr. Bishop's 7-inch refractor by Dollond.
40,67	25,19	2			370	3, 9	" " " " "
41,89	23,75	4			416	3, 9	" " " " "
41,96			1,667	1	420	3, 9	" " " " "
47,39	16,70	3			413	3, 8¼	My 6½ inch by Merz
48,75	14,53	3	1,767	3	417	3, 8	" " " " "
51,51	11,56	2	1,652	2	546	3, 8	" " " " "
52,74	10,78	2	1,683	2	375		" " " " "
53,73	7,33	3	1,768	3	467	3, 8	" " " " "
54,64	4,30	1	1,688	1	630		7½-inch by Alvan Clark.
59,58	357,72	3	1,675	3	532	3.4, 8	8¼-inch " " "
65,38	349,62	2	1,675	2	492	3, 8	8-inch by Cooke.

A fine state of atmosphere was usually present when these measures were obtained, and on several occasions the smaller star was so well seen that the observations were by no means difficult. An excellent filar micrometer by *Dollond* was used in every instance except in the year 1847.

The accuracy of the results for the present year is well supported by some measures which Mr. *G. Knott* has made with his excellent 7½-inch refractor by *Alvan Clark*; his results being: Pos. = 349°00; Dist. = 1"705; Epoch 1865,44.

It seems necessarily to follow from this, that the continued diminution of distance indicated by some of the later observations, employed by Herr *Behrmann* in his computations, probably arose from erroneous measurement. Certain

it is, that if the central distance of the small star were now what is indicated by the Ephemeris, it could scarcely be separated from the larger star even by the Poulkowa refractor; whereas, in fact, it may be well seen with an aperture of 6 or 6¼ inches (Engl.), and measured without difficulty with 7 inches on a suitable night.

Having always seen the smaller star with as much ease as might reasonably be expected under such circumstances as would induce me to measure it, I have never from my own observations suspected its brightness to be variable.

Hopefield Observatory,
Haddenham, near Thame. Sept. 1865.

W. R. Daves.

Observations de la planète (84) Clio à l'Observatoire Royal de Copenhague.

Mr. *d'Arrest* a obtenu les positions suivantes de la nouvelle planète :

1865	T. m. de Copenh.	Asc. dr.	Decl.	Etoiles de comp.
Sept. 14	8 ^h 27 ^m 51 ^s	21 ^h 20 ^m 15 ^s 12	—12° 21' 3"0	<i>a</i>
— 15	8 20 49	21 19 53,63	—12 14 32,9	<i>b</i>
Sept. 18	8 ^h 34 ^m 34 ^s	21 ^h 19 ^m 4 ^s 38	—11° 54' 36"1	<i>c</i>
19	8 16 36	21 18 52,94	—11 48 8,9	"
20	8 9 35	21 18 43,38	—11 41 31,6	"
21	8 7 47	21 18 37,04	—11 34 37,2	<i>d</i>
22	8 22 54	21 18 32,47	—11 27 ::	"
23	8 24 23	21 18 30,46	—11 20 55,1	"
25	8 46 20	21 18 34,42	—11 7 28,5	<i>e</i>
26	8 21 8	21 18 40,13	—11 0 39,0	"
27	9 16 31	21 18 47,60	—10 53 26,1	"

Sept. 18 la planète de 10^{me} grandeur,
 " 22 la planète très-faible, épais brouillard.
 " 27 éclat d'une étoile de 11^{me} grandeur.

Voici les étoiles de comparaison, déterminées récemment au cercle méridien par Mr. *Schjellerup*, qui y a joint quelques autres observations d'étoiles situées dans la route de la pla-

nète et employées par d'autres astronomes dans leurs premières observations après la découverte de Clio.

Positions moyennes pour 1865,0 :

Grandeur.			
<i>a</i>	8,5	21 ^h 17 ^m 21 ^s 23,	—12° 21' 24"3
<i>e</i>	9,7	21 18 22,87	—11 6 29,4
<i>d</i>	8,5	21 19 9,16	—11 29 51,3
<i>c</i>	8	21 20 11,52	—11 44 51,9
<i>b</i>	7,5	21 20 19,41	—12 14 55,6
	9	21 22 9,22	—12 39 55,7
	9,5	21 27 3,92	—12 55 44,6
	8,7	21 30 54,08	—14 8 19,6
	9	21 35 38,28	—14 17 4,7
	8,7	21 37 58,10	—14 17 40,9

Mr. *d'Arrest* a encore obtenu une position de la comète périodique de *Faye* :

1865	T. m. Cop.	Asc. dr. ☿	Décl. ☿
Sept. 14	8 ^h 53 ^m 2 ^s	21 ^h 46 ^m 6 ^s 73	+2° 52' 27"4

Aussi cette fois l'étoile de comparaison a été déterminée au cercle méridien. Cette position, comme toutes les précédentes, doit encore être corrigée par le moyen de la parallaxe, mais elles sont toutes corrigées de la réfraction.

Copenhague, 1865 Oct. 5.

Mynster-Fischer.