J'ajoute ici encore une éphéméride exacte pour l'intervalle du 27 Août au 12 Septembre 1894.

12^h temps moyen de Paris.

1894	α app.	δ app.	$\log A$	T. d'ab.	$1:r^2A^2$	1894	α app.	δ app.	$\log A$	T. d'ab.	$1:r^2A^2$
Août 27	3 ^h 53 ^m 39 ^s .6	+3° 32' 39"	0.16340	12 ^m 5 ^s	0.134	Sept. 4	3 ^h 59 ^m 49 ^s .6	+2° 55' 59"	0.15431	11 ^m 50 ^s	0.132
28	54 33.1	3 28 30	16228	3		5	4 0 26.4	2 50 53	15317	48	
29	55 24.5	3 24 13	16115	12 1		6	1 1.1	2 45 41	15203	46	
30	56 13.9	3 19 48	16002	11 59		7	1 33.6	2 40 23	15089	45	
31	57 1.2	3 15 16	15888	58	0.133	8	2 3.9	2 35 0	14976	43	0.132
Sept. 1	57 46.5	3 10 37	15774	56		9	2 32.1	2 29 31	14863	41	
2	58 29.6	3 5 51	15660	54		10	2 58.1	2 23 56	14751	39	
3	59 10.7	3 0 58	15545	52		11	3 21.8	2 18 16	14640	37	
4	3 59 49.6	+2 55 59	0.15431	11 50	0.132	12	4 3 43.3	+2 12 31	0.14530	11 36	0.131

Paris 1894 Juin 22.

L. Schulhof.

Elements of the Comet 1894 I (Denning)

by Dr. *J. R. Hind.*

Finding that my ellipse in A. N. 3237 differed sensibly from observations about April 25, I formed a normal place for that date, agreeing very closely with that of M. Schulhof as corrected in A. N. 3236, and combining this position, with the Hamburg observation on March 27, and that at Albany U. S., on May 9, I obtained the following orbit:

$$\begin{array}{ll}
 T = 1894 \text{ Febr. } 9.62424 \text{ G. M. T.} & \varphi = 44^\circ 50' 40''.7 \\
 \pi = 130^\circ 44' 55''.2 & \mu = 460''.9345 \\
 \Omega = 84 \quad 4 \quad 40.9 & \log a = 0.5909116 \\
 i = 5 \quad 33 \quad 19.2 & \text{Period} = 7.698 \text{ years}
 \end{array}
 \left. \vphantom{\begin{array}{l} T \\ \pi \\ \Omega \\ i \end{array}} \right\} 1894.0$$

For the Nice observation on May 22, $\Delta \cos \beta = -9''.6$, $\Delta \beta = +2''.4$ (C — O).

With these elements the nearest approach of the comet to the orbit of Jupiter is in longitude $282^\circ 30'$, distance = 0.15.

Twickenham 1894 July 9.

J. R. Hind.

Note on the group of Nebulae NGC. 3743-58.

In No. 3241 of this journal Dr. Kobold has pointed out an error affecting the places of the six nebulae NGC. 3745, 3746, 3748, 3750, 3751, 3753, 3754. As the positions of the neighbouring nebulae 3743 and 3758 are quite correct, it may not be out of place to explain how the errors arose.

Setting with Lord Rosse's six foot reflector for d'Arrest's missing nebula GC. 2464, Dr. Copeland found on Febr. 9, 1874, a group of six nebulae somewhere in the neighbourhood. These he observed again on April 5 and Lord Rosse and I observed them once in 1876. A »very red« star 8.9 mag. was micrometrically connected with them. On two other nights in 1874 Dr. Copeland observed two nebulae somewhere in the same neighbourhood and measured them from a star 8.5 mag. which on one of the two nights

was described as »reddish«. As this star had one of the 9th mag. 40^s p and 6.2 n of it, I identified it (quite correctly) as BD. +22°2380, and this star has accordingly been entered in Birmingham's and F. Krüger's catalogues as red. But the very natural mistake which I made was in assuming that the »very red« star was the same one, whereas Dr. Kobold now finds that it is BD. +22°2385, though he (curiously enough) repeats my mistake by saying »Copeland nennt ihn einmal very red, das zweite Mal reddish«.

The nebulae α and ι should therefore be removed from the diagram on p. 99 of the »Observations of Nebulae 1848-78«. Their positions in the NGC. are correct (3743 and 3758), the second one is No. 62 of Stephan's list in A. N. 2661.

Armagh Observatory, 1894 July.

J. L. E. Dreyer.

Wolsingham Observatory Circular No. 39.

The star BD. +50°2251 RA. = 16^h 1^m 5 Decl. = +50° 47' (1900) is variable.

Wolsingham Observatory, Tow Law, Darlington, 1894 Aug. 11.

T. E. Espin.