for only a single injection is required, and, when oil is used as the excipient, neither local discomfort nor febrile reaction occurs. Ten of the men were sent him daily for examination, and he found that by the sixth day their blood generally agglutinated each of the bacilli in a 1/10 dilution, in 1/100 on the eighth, and in 1/500 on the tenth day. It did not go higher for paratyphosus, but reached and remained at 1/1000 for typhosus for the next fortnight. At the end of the month agglutinating power for each infection fell off equally and rapidly. The power of deviating complement also began to appear about the sixth day, and remained in all during the latter half of the month, when it also began to disappear. As these men have been sent away on general duty we shall shortly see, he points out, whether the oily suspension of bacilli immunises them quite as well as the watery suspension more usually employed. The same author has a paper emphasising the importance of examining every chancre for spirochætæ before it has had any treatment except a boiled-water compress, and he details his rapid method of staining the smears. Sir Ronald Ross's paper on the Treatment of Dysentery is translated from THE LANCET into French.

New Inventions.

APPARATUS FOR THE TRANSPORT OF FRACTURED FEMURS.

THE mechanical contrivances for the treatment of fractured femur in this war are constantly undergoing modifications in the endeavour to deal with the varieties of conditions produced by gunshot wounds of the thigh. It would appear that in the hands of the expert the results obtained in a series of cases by any given method compare favourably with those obtained by any other. Success depends in the majority of cases upon the personal element involved, provided that the first principles in the treatment of fractures generally have not been ignored.

In the conflict of opinion a most important consideration is often ignored—namely, that of transport. Apart from active service conditions, the treatment of fractures of the thigh, whether simple or compound, from the mechanical aspect presented few difficulties since the question of transport was not involved. Herein lies the chief difficulty. Under active service conditions it cannot be guaranteed that any case of fractured femur will remain for any length of time in any one place until such a case has reached its ultimate destination in England.

During a recent period of six months at Netley it has been possible to observe the results of the overseas treatment of fractured femur from the mechanical aspect. One cannot say that the majority of cases admitted, whether of recent origin or not, were satisfactory. The almost universal method of treatment was by means of a Thomas splint with a "tie on" extension. The majority of cases, up to the time of admission there, had known no other, and these were almost without exception unsatisfactory as regards position and alignment of the bone. The few exceptions under this category were those where inquiry showed that constant daily readjustment of the extension had been carried out.

The general conclusion arrived at was that the "tie on" extension amounted to no extension. Certainly in no case admitted could the "tie on" extension be said to be doing its work. In most cases it was so loose that it did not even serve to steady the limb. In many cases which had been treated by a weight extension, the "tie on" extension being resorted to for transport, much of the good done as regards alignment and position had been nullified in the process of transport, the patient himself complaining bitterly of the pain and discomfort he had suffered since the weight extension had been removed.

In those cases where union had occurred overseas without a weight extension the results as regards position, alignment and amount of shortening may be said to have been, with very few exceptions, unsatisfactory. The same criticism applies to the majority of cases treated by the Wallace-Mawbray splint for transport.

In both cases the objections are two. 1. The amount of expert attention required to maintain extension at all, which of necessity cannot be guaranteed during transport. 2. The

fact that the amount of extension is unknown and subject to constant variation. With a weight extension it is possible to gauge the precise amount of extension, required for each case and to maintain that amount as a constant factor, or vary it precisely in accordance with the requirements of the case.

There is no doubt that the best splint for transport is the Thomas, with this essential reservation, that *there must be in addition a weight extension*. For the purposes of transport this weight extension must be self-contained, and for this purpose the following apparatus is suggested. A wooden

bracket (Fig. A), composed of an upper and under piece, is clamped to the end of the splint by a bolt and butterfly nut. The bolt lies within the curve of the end of the splint, and between the upper and under parts of the bracket, beyond the splint, rests a block to maintain rigidity. Attached to the upper surface of the bracket is a pulley wheel to carry the extension cord (Fig. A). For transport the splint is clamped to a wooden support (Fig. B) which rests on the stretcher.

In practice this apparatus has proved satisfactory. The advantage is that once the amount of extension has been applied it remains constant, requires no attention or adjustment, and nothing but deliberate interference can disturb it. In hospital the support may in some cases be employed, or the end of the splint may be rested on the foot-rail of the bed. When there is no rail, and generally, the best plan is to support the splint by a hook and cord passing over a pulley on an upright post. By this means not only is the precise angle comfortable to the patient assured, but the splint can also be raised to a steep angle and secured there if it is desired to get at the posterior surface of the limb for dressing purposes. If the post is so placed that the supporting cord to the splint pulls at an angle, an additional advantage is obtained since the tendency then is to pull the ring of the splint away from the perineum and thus avoid pressure and the possibility of pressure sores in this region.

For the efficient support of the limb in the splint strips of Sinclair's zinc should be used or a similar material giving a rigid trough in which the limb may lie at rest. Flannel bandages or slings of any such material should not be used; their tendency to stretch cannot ensure a uniform support to the limb. One has only to see the condition of affairs in such cases where strips of bandage have been used as slings to be convinced of their entire futility as an efficient positive support to the limb.

It is in the hope that the principle of weight extension in the treatment of fracture of the femur, both in hospital and continued during transport, will be adopted that these suggestions are made, and also in the earnest hope that the "tie on" extension be definitely condemned.

J. FRANKLAND WEST,
Captain, R.A.M.C., B.E.F.

N.B.—The pulley bracket is a modification of an iron pulley bracket designed by Lieutenant H. L. Wilson, R.A.M.C., and used by him at Netley, but in contrast gives perfect security, and is a universal fit to any Thomas splint.