Take at least two Röntgenographs, one in the ventral recumbent position with the anode centered over the sacrococcygeal articulation; this should be a standard exposure for the bladder, always preserving the same distance from anode to plate. Center as above and a Röntgenograph will result from which reliable comparison as to size, position and conformity can be made. The second exposure can be taken at any desired lateral or dorsal angle. The proper angle found, stereo-Röntgenographs would be of great value.

The bismuth solution is well borne by the bladder. In one patient with cystitis it was the only injection which he could retain with comfort. The solution can be removed by irrigation after the exposures have been made, or allowed to pass off naturally. The danger of bismuth poisoning from absorption is practically nil.

URETEROTUBAL ANASTOMOSIS

A PRELIMINARY REPORT

AXEL WERELIUS, M.D.
CHICAGO

Owing to the rather discouraging results following partial excision of bladder for malignant disease, the complete removal of the organ is, as a rule, the only procedure that holds out any hope of lasting cure. The all-important problem in such radical work is, of course, to establish an outlet for the urinary secretion, and one that combines the greatest insurance against ascending infection with the least annoyance to the patient.

Several more or less ingenious methods have been devised. Watson proposes preliminary nephrotomy and permanent drainage through the loin. This method may afford fairly good protection against kidney infection, but must be frightfully annoying to the patient.

Ureterotubal anastomosis.

Maydl's¹ plan of ureteral implantation into the intestine has been used. This method, or its modifications, originally devised for exstrophy of bladder, permit of control of urinary secretion, and would be excellent were it not for the great danger of ascending infection. The death-rate from this source has been 50 per cent. This is higher than when used for its original purpose, as in exstrophy of bladder, from long exposure to the exterior, ureters have acquired some degree of immunity to ordinary types of infection. The danger of infection in above operation is not only immediate, but threatened at any subsequent enteritis.

Pawlik² implanted the ureters into the vagina, Sonnenberg³ into the urethra and Rovsing⁴ with others,

1. Maydl: *Wien. med. Wchnschr.*, 1894, Nos. 25-29; 1896, No. 28.

2. Pawlik: *Wien. klin. Wchnschr.*, 1891, xli, 1814, 1815.

3. Sonnenberg: *Verhandl. d. Deutsch. Gesellsch. f. Chir.*, 1881, vol. x, No. 11.

4. Rovsing: *Hospitalstidende*, 1907, xv, 709.

left ureters in the skin opening, all with good immediate results.

Thus it seems that operations that aim to give the patient the greatest comfort are also associated with the greatest danger. The genital system, closely related as it is to the urinary apparatus, both in the early embryo and throughout life, should, it seems, be the ideal outlet for the urinary secretion. Through this route secondary infection ought to be reduced to a minimum. The only drawback is, of course, lack of control, but this is somewhat counteracted by the favorable location for wearing of a urinary receptacle.

Working along the above theory, I am now endeavoring to anastomose ureters with the tubes, as in the accompanying illustration and also to implant them into the uterus. The immediate results in dogs seem very promising. Besides a trial dog, thus far only two dogs have been used of a coming series.

Dog 1.—Brown and white, large bitch. Operation, January 4. Anesthesia, ether. The ureters were dissected up and cut off close to the bladder and distal end tied. Bladder was left intact. Both tubes were then cut off about 4 cm. from the uterus and proximal end tied. A through-and-through purse-string suture was then inserted in the distal end of the tubes, into which the ureters were drawn, as in the illustration, and the purse-string lightly tied around the tubes and reinforced by two interrupted sutures.

Dog 2.—Grayish-white bitch. Operation, January 9. Same technic as above. Both dogs are so far doing well, passing urine through the uterus and vulva.

The trial dog mentioned above was very small, with minute ureters, and consequently, very difficult to operate on, especially as the operation had not been thoroughly planned. Owing to recent pregnancy there was a great disproportion between the tubes and ureters, the caliber of tubes being many times larger than that of the ureter. In making the anastomosis, I foolishly used interrupted sutures, which made the union very imperfect. The dog died on the sixth day, urine having leaked at the anastomosis.

It is, of course, too early to report any results or give any conclusions, as the work has just begun, but the operation seems feasible, at least in dog.

1230 Sixty-Third Street.

THE FUNCTION OF THE CHORIOID PLEXUSES OF THE CEREBRAL VENTRICLES AND ITS RELATION TO THAT OF THE PITUITARY GLAND *

SIMON PENDLETON KRAMER, M.D.

Surgeon to the Cincinnati Hospital
CINCINNATI

The secretory organs for the cerebrospinal fluid are the chorioid plexuses of the third, fourth and lateral ventricles. These are highly vascular projections of the pia mater into the ventricles, covered with villous-like projections about 1 to 2 mm. in diameter. Under the microscope these villi are seen to be made up of a number of secondary villi about 0.25 mm. in diameter, which again show grape-like projections. Through the center of the villi run comparatively thick-walled blood-

*Read before the Cincinnati Neurological Society, and the Cincinnati Society for Medical Research, Jan. 5, 1911.

*For reasons of space, part of the illustrations are omitted. All appear in the author's reprints.

Fig. 1.—Dec. 21, 1910. Dog, 6 kilos; ether anesthesia. Jugular injection at I, of 1 c.c. extract of dog chorioid. Uppermost line, timer marking seconds; second line, blood-pressure; third line, respiration.

Fig. 4.—Dec. 28, 1910. Dog, 12 kilos; ether anesthesia. Jugular injection, at I, of 12 c.c. cerebrospinal fluid from a patient with cerebral edema of traumatic origin. Lines as before.

Fig. 6.—Dec. 31, 1910. Same dog as in Figure 5. Injection, at I, of 10 c.c. cerebrospinal fluid from same case (Fig. 5) of delirium tremens. Lines as before.

Fig. 7.—Jan. 3, 1911. Dog, 6 kilos; ether anesthesia. Jugular injection, at I, of 30 c.c. cerebrospinal fluid from an individual recovering from mild attack of delirium tremens. Lines as before.

Fig. 8.—Dec. 10, 1910. Dog, 11 kilos; ether anesthesia. Jugular injection, at I, of 0.75 c.c. pituitary extract. Lines as before.

Fig. 9.—Dec. 30, 1910. Dog, 12.5 kilos; ether anesthesia. Jugular injection, at I, of 1 c.c. extract of human chorioid plus 1 c.c. of pituitary extract. Lines as before.

vessels, giving the entire organ a highly vascular appearance. The connective tissue of the villi is that of the pia mater. Covering the villi of the plexuses is a layer of large spheroidal cells, in each of which may be seen, in addition to the nucleus, yellowish granules. These cells are probably the secretory cells of the organ. The appearance of the plexuses strongly suggests an inverted gland which, instead of pouring its secretion into a series of ducts, empties directly into the ventricular system of the brain.

If one strips the chorioid plexuses from both the lateral ventricles of a dog recently killed by bleeding, and rubs them up in 2 c.c. of normal saline solution and injects the filtrate into the jugular vein of another

a man 35 years of age, five and one-half hours after death from pneumonia.

Extracts made from the chorioid plexus of the fourth ventricle produce the same reaction.

The reaction exhibited in Figure 3 is independent of the action of the vagus nerves, since the same effect is obtained when both vagus nerves have been divided.

The thought suggested itself that possibly in certain cerebral affections, particularly in such as were associated with hypersecretion of cerebrospinal fluid, the fluid might contain an excess of this depressant evidently secreted by the chorioid plexuses. Accordingly experiments were made with cerebrospinal fluid obtained by lumbar puncture in three such cases.

Fig. 4, continued.

dog, it will be found that this extract causes a marked fall in the blood-pressure. This effect was found to be constant in ten experiments.

Figure 1 is from a tracing of such an experiment. The blood-pressure was obtained by connecting the carotid or femoral artery of the animal with a mercury manometer. The respirations were recorded by means of a Paul Bert pneumograph connected with a Marey tambour, so arranged that the up-stroke represents inspiration, the down-stroke expiration. The time of injection was recorded by an electric signal. The injection was made into the jugular vein. In all experiments the animals were under light ether anesthesia administered through a tracheal tube. This animal received 1 c.c., or half the total extract made from the chorioid plexuses of the lateral ventricles of another dog killed by bleeding. The drop in blood-pressure, which in this experiment was preceded by a slight momentary rise (not constant), began fifteen seconds after the beginning of the injection, reached its maximum (that is, 50 mm.) twenty-eight seconds after the beginning of the injection.

Figure 4 is a tracing obtained from a dog into whose jugular vein were injected 12 c.c. of cerebrospinal fluid obtained by lumbar puncture from a patient with edema of the brain following a blow to the head producing a subtentorial hemorrhage. It will be seen that the injection caused a marked (55 mm.) and prolonged fall in blood-pressure, followed by an exceedingly slow and incomplete recovery. The cerebrospinal fluid in this case evidently contained a great excess of the "chorioid depressant."

Figure 5 is a tracing obtained from a dog into whose jugular vein were injected 5 c.c. of cerebrospinal fluid drawn from a marked case of delirium tremens. This fluid was much more depressant than that from the previous case.

The injection of 10 c.c. of the same fluid fifteen minutes later into the same dog produced a most violent and prolonged depression, from which the animal incompletely recovered only after an interval of eight minutes. This is shown in Figure 6.

Fig. 8, continued.

tion, and returned to within 5 mm. of the original pressure in two minutes and twenty-five seconds. The rate of the heart-beat was apparently unchanged. The respiration was increased in rhythm from 40 to 60 per minute. Extracts made from the chorioid plexuses of the human brain show the same effects in even smaller doses.

Figure 2 is a tracing obtained by injecting into the jugular vein of a dog 1 c.c. of an extract made by rubbing up the plexus from one lateral ventricle in 10 c.c. of normal saline solution. The brain was removed from

Figure 7 is a tracing obtained by injecting 30 c.c. of cerebrospinal fluid drawn from a subject who had recovered from delirium tremens. There was but little depressant effect, a fall in blood-pressure of not more than 5 mm., which was quickly recovered. The individual from whom this cerebrospinal fluid was obtained had had a mild attack of delirium tremens which had lasted only forty-eight hours, and, at the time the fluid was withdrawn, the patient's pulse was normal again and the attack of delirium tremens had disappeared.

The patient died from whom the fluid used in Experiments 5 and 6 was obtained, and at autopsy a marked cerebral edema was found. The ventricles were filled with fluid which, when injected into dogs, was markedly depressant.

It also occurred to me that if such a depressant were secreted by the chorioid plexuses and poured out into the ventricular system it would meet there whatever was given off by the pituitary body, and it seemed wise to determine, if possible, what action these two principles might have on each other. As has been abundantly shown, the infundibular lobe of the pituitary gland contains a substance which causes a marked elevation in blood-pressure lasting for some time and accompanied by a marked slowing of the heart and an increase in the force of the cardiac systole.

Tracing 8 was obtained from a dog into whose jugular vein was injected 0.75 c.c. of an extract of the infundibular lobe of the sheep's pituitary gland. The well-known effect is here nicely shown.

When an injection was made of a mixture of the pituitary extract just mentioned and an extract of the human chorioid plexus it was found that these substances tend to counteract each other, though incompletely. Of course, it is difficult to determine the completeness or incompleteness of such a counter-action in the absence of known dosage.

If we examine a tracing (Fig. 9), of such an experiment we shall find it very interesting. The depressant effect of the chorioid is shown, but lessened in amount and only transitory in time. This is succeeded by a rise in pressure due to the pituitary extract but this also is lessened in amount (15 mm.) and greatly shortened in time of duration. At the end of two minutes and thirty seconds the blood-pressure was within 5 mm. of that which obtained before the injection. The slowing of the heart is present but also reduced. It is evident then that these two principles which, we presume are poured into the cerebrospinal fluid have a tendency to counteract each other in so far as their effect on the circulatory apparatus is concerned. It will be a matter for future investigation to determine, if possible, under what circumstances the one or the other gains the supremacy.

A further report on the nature of the chorioid "depressant" will also be made in the future.

4 West Seventh Street.

OPTIMISTIC THERAPEUTICS

GEORGE M. NILES, M.D.

Professor of Physiology, Southern College of Pharmacy; Associate Professor of Physiology, Atlanta School of Medicine; Gastro-Enterologist to the Tabernacle Infirmary
ATLANTA, GA.

Some months ago the Hon. Joseph Choate, at a celebration commemorating the landing of the Pilgrim Fathers, suggested that it might not be amiss to extol the Pilgrim Mothers also, who not only had to endure equally the hardships and privations undergone by the Pilgrim Fathers, but also had to endure the Pilgrim Fathers.

Along this line the thought has occurred to me that, while tomes have been written concerning the psychic attitude of our patients—how they should be lifted from the slough of disease and despond, and their feet planted on the solid ground of health and right thinking; how to combat the demon of introspection; how to

stem the increasing tide of "Americanitis," and so on *ad infinitum*—but little has been said about the psychic attitude of the physician who has these problems to shoulder as well as the mental foibles of the sick. He has to not only bear with the disease, but also bear with the changing, the complaining, and often repining personality of "experienced invalids," who know how to run up and down the gamut of symptomatology like an expert pianist, or who can argue from obscure cause to mysterious effect like a trained advocate.

We all have them: the surgeon hears of minor injuries entailing dreadful and far-reaching consequences; the gynecologist is regaled with lurid accounts of horrible ovarian and uterine pains; the ophthalmologist or aurist is told of ringing and singing sounds in the ears, or remarkable spots before the eyes assuming all manner of fantastic shapes; the internist hears of jumping and bumping hearts, of existences eked out with one lung completely gone, and the other one nearly out of commission, or of fierce pains in the back, denoting of course serious kidney disease; while the gastro-enterologist, I imagine, hears the most bizarre recitals of all. I am told sometimes of phenomena following the ingestion of simple articles of food which are neither explainable by physiology nor pathology, and which would make Sindbad the Sailor or Baron Munchausen turn green with envy. A prosaic piece of bread or an ordinary slice of roast beef, when deposited in the alimentary tract of some of these nervous dyspeptics, immediately becomes almost endowed with life, assuming a gastronomic importance anywhere from romantic to tragic. The wholesome meat of to-day becomes the poison of to-morrow, while some now despised viand may at the next visit be a longed-for delicacy.

Scientific diagnosis is being well taught at present by practically all the medical colleges. Each institution is vying with the other in furnishing thorough laboratory and clinical instruction, so that every medical graduate who lives up to his opportunities is amply equipped to diagnose disease as it comes before him. I am also glad to observe a healthy revival in therapeutic instruction. The belief in the efficacy of many forms of therapeutics—drug and otherwise—has been much undermined by the German school of therapeutic nihilists, aided and abetted, I regret to say, by one of our most eminent American physicians, who has recently taken up his abode in England.

Etiology, pathology, morbid anatomy, and symptomatology of disease have their essential place; but what the patient is interested in is the treatment and prognosis. When I read page after page of pathology, etc., and finally find the treatment briefly and inadequately expressed in a few lines, I feel that I have asked for bread and been given a stone.

Differential diagnosis appeals to the average lay mind, even though it be an intelligent one, about as much as the anatomic difference between a dinosaur and an anthropoid ape. The question "What is the matter with me, Doctor?" is overshadowed by the more pointed ones "How long shall I be sick?" or "Shall I get well?" and the doctor who gets busy at once doing something, whether it be an important therapeutic procedure or only a measure to relieve symptoms and render the patient more comfortable, until he can learn "where he is at," acquires an immediate influence not to be despised.

Far be it from me to decry any of the niceties of diagnosis, or the judicial poise of mind necessary to correct discrimination, where lines of diagnostic demarcation