The marked improvement in the objective findings in the spinal fluid following treatment, in practically all forms of syphilis thus treated, even in the absence of subjective improvement, lead one to hope that this form of treatment is along the right line for checking or, indeed, in some cases even curing central nervous involvement. I am led to believe that my earlier cases would have done better with injections of even smaller doses than those which were given. Subsequent injections will be given in small doses and diluted with a greater amount of the cerebrospinal fluid.

Cases to be thus treated should be carefully selected. For the present at least, this form of treatment should be restricted to those cases in which other forms of treatment have proved inefficient.

In all cases the danger of the treatment in its present unperfected state should be pointed out to the patient and his family, and the responsibility for its administration should be shared by them. It is to be hoped and expected that more careful study and selection of cases will prove the intradural injection of neosalvarsan a potent remedy in combating cases of cerebrospinal syphilis.

602 East Liberty Street.

HAY-FEVER

R. CLAUDE LOWDERMILK, M.D.
GALENA, KAN.

PROPHYLAXIS AND CURE WITH POLLEN TOXIN

During the season of 1913 I treated nineteen cases of hay-fever with successful results in sixteen. Of three patients who were treated before the onset of symptoms, not one had an attack throughout the season; and of sixteen in whom treatment was begun after the onset, thirteen were cured. Relief of symptoms was very prompt, usually within an hour or two and seldom later than twenty-four hours.

The method used is essentially a modification of the active immunization suggested by Allen and others, with the addition of a bacterial vaccine in cases complicated with infection.

THE TOXIN

The toxin was prepared as follows: One gram of mixed pollen (250 mg. *Ambrosia artemisiaefolia*, 250 mg. *Ambrosia trifida*, and 500 mg. different varieties of *Solidago* found in this locality) was mixed with about 10 gm. of sterilized fine sea-sand in a sterile mortar; 100 c.c. of physiologic salt solution with 0.5 per cent. phenol (carbolic acid) was measured out and the pollen moistened with a portion of it. The mass was then ground for several hours, with the addition of portions of the salt solution, until the sand was reduced to an impalpable powder. The remainder of the salt solution was then added, the whole transferred to a sterile flask and allowed to stand at room temperature for twenty-four hours with frequent shaking. The fluid was then pipetted off, centrifuged and sealed in glass ampules, each containing 1 c.c. This constituted the stock solution and was stored in a cool, dark place.

DOSAGE AND ADMINISTRATION

For convenience, the unit suggested by previous workers was used. This unit is the quantity of toxin extracted from 1 microgram (0.000,001 gm.) of pollen. As prepared, each cubic centimeter of stock solution

contained ten thousand units. From the stock solution two serial decimal dilutions were made containing one thousand and one hundred units to the cubic centimeter, respectively. By using a tuberculin syringe graduated in hundredths of a cubic centimeter it was possible to administer doses containing any desired number of units while the total volume of the dose never exceeded 1 c.c.

The site of injection was painted with tincture of iodine and the injection made subcutaneously in the usual manner. In no case was there any serious trouble at the site of injection.

The dosage in each case was determined entirely by the nature and extent of the reactions produced. Efforts to determine the proper dosage in advance by means of ophthalmic and cutaneous reactions were found to be unreliable. The interval between doses was from five to eight days at first, but this was reduced after a little experience to one or two days, depending on the time required for the reaction from the previous dose to subside.

TREATMENT AND REACTIONS

Three patients were given treatment beginning about a month before the time for the annual onset. All were adult men and had suffered annual attacks for eight, twenty and thirty-five years, respectively. The first received eight injections of from 25 to 1,000 units each; the second, six injections of from 25 to 700 units; the third, six injections of from 25 to 500 units. Doses were given at intervals of from five to eight days. Complete immunity resulted in each case.

In all true cases of hay-fever the administration of a proper dose of toxin is followed by a well-marked reaction. This begins in from half an hour to two hours and is characterized by an exaggeration of the usual symptoms of an attack; sneezing, lacrimation, itching of the eyes, cough, dyspnea in asthmatic subjects, and sometimes edema and urticaria. In the third case reported above, the third dose was increased from 100 to 300 units after an interval of seven days and was followed by a reaction which continued for about eighteen hours and caused the gravest alarm; there was a violent general urticaria, severe edema of the neck and throat, dyspnea and vertigo. The reaction terminated abruptly, the patient felt no ill effects from it, and consented to go on with the treatment. He had never been able to go near ragweed when flowering, but last season he handled it with impunity.

August 11 a patient came with a well-developed and rather severe case of three days' duration. For seven years he had suffered annual attacks of such severity as to compel a change of climate. The present attack had begun in the usual manner. I gave him 5 units; the reaction came on in half an hour and continued for two hours, followed by a sudden remission of all symptoms. I gave him eight injections of toxin at intervals of one and two days, the last being 1,000 units. He had no further trouble and continued at his work.

Fifteen others were treated after the onset of the annual attack; the number included both sexes and their ages ranged from 20 to 60 years. The initial dose varied from 5 to 50 units and the final dose from 100 to 1,000 units. The interval between doses was usually one day, sometimes two and seldom greater than two.

Of the sixteen patients treated after the onset of the annual attack thirteen were cured. The three not

benefited all began treatment more than a month after the onset of symptoms.

Vaccine as an Adjunct.—In all cases complicated by the presence of a bacterial infection injections of autogenous vaccine were given. Sometimes the vaccine was alternated with the toxin and in others it was given afterward. I consider this a very important element of the treatment which should never be neglected.

CONCLUSION

In spite of the conflicting theories concerning this form of treatment, the fact remains that the patients were cured—and “facts are stubborn things.” Unfortunately the method here described is not suitable for use by the general practitioner. To use it successfully requires an intimate acquaintance with the various phenomena of immunity. I offer it in the hope that it may be taken up by others and so perfected that all physicians may use it to the end that humanity may be relieved of this most distressing of all non-fatal diseases.

THE ISOLATION OF SPIROCHAETA PALLIDA FROM THE BLOOD IN SYPHILIS

HARRY F. HARTWELL, M.D.

From the Pathological Laboratory of the Massachusetts General Hospital
BOSTON

The rabbit's testicle provides a favorable environment for the growth of *Spirochaeta pallida*. It has been extensively used for the study of experimental syphilis and for the demonstration of spirochetes in syphilitic material. Uhlenhuth and Mulzer¹ were the first to show the infectiousness of the blood in syphilis by the injection of rabbits; Nichols and Hough² and others have had positive results from the injection of cerebrospinal fluid; Graves³ has reported spirochete-containing lesions with blood from tabetics and general paralytics.

In this study the blood from twenty-four untreated patients with early syphilis has been injected. It was withdrawn from a superficial vein of the forearm with aseptic precautions as in taking a blood-culture, and was defibrinated in a sterile test-tube. Two c.c. of the defibrinated blood were then slowly injected into the testicle. One animal was used for each case; in some cases one, in others both testicles were injected.

The latent interval after injection before lesions appeared varied from a little over one month in the earliest case to a little more than three months in the longest. In some the whole testicle was enlarged and quite hard; in others circumscribed nodules of cartilaginous hardness could be felt. The process sometimes showed marked thickening of both layers of the tunic. The nodules were usually situated in the upper pole of the gland, in or near the epididymis. In examining the testicle small lesions may be missed; the nodular feel of the normal epididymis and its apparent variation in size and consistency make the detection of these small areas of induration more difficult. In

one of our animals the epididymis was noted as hard two months after the injection of a suspected blood, and the testicle was aspirated on two successive days with negative results. It was excised eight months after injection, and motile spirochetes were found in the dark field, although no gross lesions were evident.

As soon as lesions were palpable, the animal was etherized and the testicle excised. The cut surface of a nodule yielded a clear sticky fluid, which contained large numbers of spirochetes. In microscopic sections this mucinous exudate was found in a loose fibrous reticulum, and in tissue stained by Levaditi's method the organisms were present in these areas in the largest numbers. Uhlenhuth and Mulzer⁴ and Noguchi⁵ consider this exudate peculiar to syphilitic orchitis of rabbits.

THE PROTOCOLS OF THE POSITIVE CASES ARE AS FOLLOWS:

CASE 1.—Man, aged 35. Primary lesion of penis, two weeks' duration. Glands in groins. No secondaries. Wassermann weakly positive. Feb. 15, 1913, blood injected into both testes, Rabbit 14. Spirochetes were found in the dark field and in Levaditi sections.

CASE 2.—Man, aged 29. Primary lesion of penis, eighteen days' duration. Multiple glands in groins. Primary side of tongue twenty-three days. No eruption. Wassermann strongly positive. Feb. 29, 1913, blood injected into left testis, Rabbit 15. April 20, 1913, diffuse orchitis and peri-orchitis with spirochetes in dark field and in Levaditi sections.

CASE 3.—Man, aged 28. Primary lesions of penis, four weeks' duration. Multiple glands in groins. No skin eruption. Wassermann strongly positive. March 29, 1913, blood injected into both testes, Rabbit 29. June 15, 1913, left circumscribed nodule in epididymis with spirochetes in dark field and India ink smear; none found in Levaditi sections.

CASE 4.—Man, aged 22. Primary lesion of penis, two months' duration. General papular eruption. Wassermann strongly positive. Oct. 21, 1912, blood injected into both testes, Rabbit 4. Jan. 1, 1913, left diffuse orchitis, spirochetes in dark field and Levaditi sections.

CASE 5.—Woman, aged 18. Papules about mouth, forehead and scalp. Fading macular eruption over body. General glandular enlargement. Condylomata about anus. Wassermann strongly positive. Jan. 7, 1913, blood injected into left testis, Rabbit 7. Feb. 23, 1913, diffuse orchitis with peri-orchitis. No spirochetes found in dark field. In the Levaditi sections they were numerous.

CASE 6.—Man, aged 30. Primary lesion of penis with general roseola. Wassermann strongly positive. Feb. 20, 1913, blood injected into both testes, Rabbit 16. April 18, 1913, left diffuse orchitis with spirochetes in dark field and Levaditi sections. In the right testicle nodules could be felt. It was excised Sept. 23, 1913, and spirochetes were found in the dark field.

CASE 7.—Woman, aged 26, pregnant five months. General roseola. Wassermann strongly positive. Feb. 21, 1913, blood injected into left testis, Rabbit 19. April 22, 1913, circumscribed nodule in epididymis with spirochetes in dark fields and Levaditi sections.

CASE 8.—Woman, pregnant six months. Husband treated in clinic for syphilis the previous summer. Brilliant maculopapular eruption over body. Wassermann strongly positive. March 22, 1913, blood injected into both testes, Rabbit 26. May 18, 1913, left diffuse orchitis, spirochetes in dark field and Levaditi sections. The right contained palpable nodules and was excised July 8, with spirochetes in dark field.

CASE 9.—Man, aged 29. Primary lesion of penis, two months' duration. General glandular enlargement. Was-

1. Uhlenhuth and Mulzer: Berl. klin. Wehnschr., 1912, No. 4, p. 152; Centralbl. f. Bakteriöl., Orig. Abt. 1, 1912, lxiv, 165.

2. Nichols, Henry J., and Hough, William H.: Demonstration of *Spirochaeta pallida* in the Cerebrospinal Fluid, THE JOURNAL A. M. A., Jan. 11, 1913, p. 108.

3. Graves, William W.: Can Rabbits Be Infected with Syphilis Directly from the Bloods of General Paretics? Observations on the Recognition of the Virus in the Later Periods of the Disease, THE JOURNAL A. M. A., Oct. 25, 1913, p. 1504.

4. Uhlenhuth and Mulzer: Arb. a. d. k. Gsndtsamte, 1913, xlv, 307.

5. Noguchi: Jour. Exper. Med., 1912, xv, 202.