physicians. Education alone will not bring about the contemplated result, for the simple reason that a man about to visit a place where gonorrhea is liable to be contracted is not bothering his mind about logic or pamphlets, but in most cases is under the influence of liquor, and does not remember any injunction he has read. For two years Dr. Robin has lectured to students of the state college on hygienic subjects, and also on venereal diseases. He found that the best results could be achieved through the personal relationship between the students and himself. He put himself in a position that caused him to be looked on by the students as their friend and adviser. One boy came to him and told him that he had contracted gonorrhea. Dr. Robin offered to examine the secretions for him, as often as necessary, explaining the seriousness of the disease, and volunteered such information as would be of benefit to him. He also pointed out the necessity of abstaining from intercourse or excitement. The result was very much better so far as that boy was concerned than if he had given him a single lecture and allowed him to go. Eight months after the attack he asked Dr. Robin if he would allow him to drink gin, he not having tasted liquor for the entire period, although greatly tempted. He was under treatment by another physician, and Dr. Robin referred him to his family physician. What is needed in the colleges is some physician to come in contact with the boys in the capacity of teacher or adviser, but not for the purpose of treatment. There should be someone to whom the boys can go and ask advice about themselves. Dr. Robin considers that education should be personal rather than through literature. Physicians know just about what effect literature has, at least those who have published bulletins for distribution among other physicians. Many people are in the habit of reading nothing unless it is spicy or sporty. In this connection there ought to be also some form of legislative enactment. While reglementation is decried by a great many people, yet it appears that when properly carried out it may be valuable. The control, however, should be placed in the hands of sanitary and not police officers. There should be plenty of dispensaries which would enable the poor young man to get away from the quack and the dispensing druggist, and these dispensaries should be under the control or supervision of the boards of health. An effort should be made to work on all sides rather than to confine the work to circulars alone.

DR. HENRY B. BAKER, Lansing, Mich., stated that in Michigan a law has been in force about nine years, which requires that there shall be taught every year, in every public school of the state, the means by which dangerous communicable diseases are spread, and the best measures for their restriction. There is no text-book on that subject, and if there were, it would probably be out of date in a year; therefore, the law of Michigan requires that the State Board of Health every year shall supply the data and statements which will enable the teachers to comply with the law. There are about twenty thousand teachers in the state, and this has been done for nine years. The diseases which have been dealt with in that manner are consumption, diphtheria, scarlet fever, typhoid fever, and so on, and the relative importance of these diseases has been set forth in leaflets supplied to the teachers. Syphilis and gonorrhea are not included in the instructions, notwithstanding the law as it stands would necessarily include them. A year or two ago the State Board of Health appointed Dr. Baker a committee to work out some plan whereby these diseases could be successfully dealt with. In each state the state board of health should be an educational force along this line. In Michigan, this law, for nine years, has worked wonderfully well in regard to the other diseases. Under that law there is supplementary action by the State Board of Health, which would be especially valuable for education along these lines.

DR. FRANK W. GOODELL, Effingham, Ill., said that education may be good or bad. A man may receive a moral education, and if he wants to become a first-class safe blower he receives an education in that line also. It is a question which has two sides of nearly equal strength and weight, as to whether or not men and women should be taught how to prevent the injurious effects of what would otherwise be injurious copulation. If

women are taught how to prevent venereal disease, and then taught how to prevent conception, prostitution will increase to an alarming extent. About the only way to prevent the contraction of these diseases is by impressing on the people the fact that a dangerous disease is apt to follow the committing of the indiscretion. Fear seems to be stronger than anything else to prevent crime. There is no one means which will alone prevent the contracting of these diseases, but the enemy should be attacked from every point. Another thing which has not been referred to is that of honor among men as well as honor among women. Nothing has been said about the seduction of men. Men are not prostitutes. So long as women continue to be more than ordinarily attractive to men just so long men will keep up the pursuit. Morality ought to be talked equally to both. In talking with a man who had been much among Indian women Dr. Goodell asked him in regard to the virtue of Indian women as compared with the virtue of our women, and he said it was astonishing, but true, that virtue among Indian women at the time he knew them was equal to virtue among American women. Dr. Goodell said that he knows of no particular means which he could recommend, but thinks that all means should be employed to eradicate the social evil.

DR. THOMAS GRANT YOUNG, Columbus, Ohio, stated that one of the most important present-day problems is the protection of the innocent from the latent gonococcus. Most men who have been once infected, and who contemplate matrimony, seek medical advice to determine whether or not they are free from infection. Many times this advice is worthless. Therefore, one of the most important things is to show the people, and especially the men who have charge of the medical laws, that the druggist is not a competent man to treat gonorrhea; for many of these cases are treated originally by the druggist. In the second place, the general practitioner should be impressed with the seriousness of the disease. He should be shown that it requires men of special training and equipment properly to treat gonorrhea.

DR. J. A. WORK, Elkhart, Ind., declared that early training in the schools is a grand work. Solomon, although unwise in some things, yet gave the best advice in regard to the training of children. His advice was to "train up a child in the way he should go, and when he is old he will not depart from it." John G. Holland teaches that a young man should keep himself pure, and Dr. Work believes that personal purity should be taught to the infant. He has seen children that were scarcely out of their short dresses who had impure thoughts.

DR. L. DUNCAN BULKLEY, New York City, said that education is well enough, but until these venereal diseases are put on the list of diseases to be cared for by the boards of health, physicians will never accomplish their purpose.

A STUDY OF METABOLISM IN A CASE CONJOINING MYXEDEMA AND DIABETES MELLITUS.*

AUGUST ADRIAN STRASSER, M.D.

ARLINGTON, N. J.

The metabolic changes in the adult have at sundry times and by many careful investigators received due attention, but it is very infrequent to see them investigated in childhood, whether in health or disease; and meeting a case where this research could be carried out and where the bodily health was away from the normal, I have herein set myself the task of studying the metabolism in a child suffering not only with an aggravated form of diabetes mellitus, but where such metabolism was further modified by the fact that he was a congenital cretin.

In the light of recent assertions as to the genesis

* Read in the Section on Diseases of Children of the American Medical Association, at the Fifty-fifth Annual Session, June, 1904.

of diabetes mellitus and the interrelation of it and other dyscrasias of similarly hitherto unknown etiology, the case assumes still further interest if we debate the question of the interrelation of the two dyscrasias found here. How far was the improvement in the child's mental status due to the small quantity of thyroid extract exhibited and the short time it was employed? How far was it influenced by the secondary disease, the diabetes mellitus? Was the chronic character of the disease at all influenced by the thyroid condition?

Of course these are purely speculative fields, but viewed from the standpoint of our present lack of positive knowledge, this contribution may give an impetus to further study along these lines to pediatricists, though I am well aware that two such formidable nutritive dyscrasias are very rarely found in conjunction in the one subject, and I consider my opportunity to study the case one of good fortune.

It may be best at this point to enter into the history of the case, to serve as a basis for the future deductions my experiments along the lines of metabolism have allowed me.

CASE 1.—On July 7, 1902, A. R. was brought to my office for diagnosis and treatment, if possible. From his mother I elicited the following history:

Family History.—In paternal line, father is of nervous temperament, but not pathologically. Grandparents normal. One aunt was very nervous and procreated an imbecile. In maternal line, no abnormality of any sort and grandmother still living. On both sides no history of venereal or neurotic affections with the single exception above.

Personal History.—The patient is the last of four healthy and normally procreated children. The mother lays stress on the fact of having suffered great anxiety because of financial reverses at the time of his gestation, but beyond that is at a loss to account for the child's condition. Symptoms of cretinism were visible since birth, but the diagnosis was not made, although the attending physicians seemed to feel that something was not in order. The child was breastfed and the characteristics of his disease, as is usual, did not develop until after he was weaned. He did not attempt to walk until after he was three years old. He suffered with frequent gastric and cardiac crises. After weaning, he became abnormally fat, developed a blank, mask-like face, with stubby nose and thick lips, and old photographs show the typical features of an early case of infantile myxedema. A year ago he was taken to the Postgraduate Hospital, sent there by his physician, where at last the diagnosis was made and suitable treatment by means of thyroid extract instituted.

Present Status.—Child is at present eight years old. He is heavy for his age, has a dull, uninterested face, mask-like, with depressed nose, open mouth, fat, everted lips, prognathic, thick, protruding tongue, lolling speech if a rare word escapes him, slow but not entirely absent mentality. Absolutely no thyroid gland tissue is palpable. Lumps of tissue over the clavicular spaces were present. The abdomen was large and the umbilicus protruded. The epiphyses were enlarged at the knees, ankles, elbows, and wrists. The muscular system was flabby and without tonicities. The skin was dry, scaly and boggy; no perspiration had ever been noticed. The hair was sparse, wiry and lusterless.

Treatment.—As the child at this visit had had no thyroid extract for over four weeks and had previously improved under its use, to judge from the mother's statement, its administration was resumed and in addition to aid his digestion and prevent the ever recurring gastric crises, a pepsin mixture was prescribed.

Keeping him under observation there was a distinct improvement, not only in the loss of the redundant tissue, but in increased intelligence. Although very weak at times, he

was much brighter and slightly more active. He had begun to combine a few words into phrases.

Occurrence of Diabetes.—With occasional short intermissions, the thyroid feeding was maintained until in September, 1902, his mother brought him to the office with the history of a new condition which had supervened in the last two weeks. After or perhaps because of over eating of grapes and ripe fruits of other sorts, the child had developed a glycosuria, which the attendant train of symptoms stamped as a diabetes. He had a constant violent thirst, was ravenously hungry and almost constantly micturating.

Urine.—Analysis of the urine showed 10 per cent. of sugar, and some days later even 12 to 15 per cent., passing four liters in twenty-four hours.

Treatment.—He was immediately placed on strict diabetic diet; Vichy given to allay the thirst and alkalize the urine, codein sulphate 0.006 gm. was given four times a day. For the obstinate constipation the fluid extract of cascara was used.

Knowing that occasionally thyroid feeding produced a temporary glycosuria, the tablets were stopped entirely to eliminate them as factors.

Two weeks of this treatment availed little, except that it caused less frequent micturition and less craving for food. The drowsy condition superinduced by the codein was objectionable, however, and the fact that the child was rapidly failing from the withdrawal of the carbohydrates and fats, caused the resumption of a less restricted diet. For a month I did not see the child again, and when he was brought it was found that although analysis showed only 8 per cent. of sugar by trituration and by Robert's fermentation tests, still the myxedematous condition was coming back. The child's intelligence was even less acute than before, mentality was painfully slow, his tongue was protruding worse than ever and the abdomen at umbilicus was larger than at any time I had ever seen him.

As evidently the exhibition or withholding of thyroid extract had made so very little difference on the course of the diabetes, I determined to resume the use of the tablets to at least keep up the good they had before accomplished. The parents stopped them, however, of their own accord, as the child was failing rapidly; and he never received any further thyroid extract in any form since that time. However, later developments showed that this was not essential in his case, as evidently the diabetes itself to a certain extent influenced the cretin condition, as witness the facts that at the time of making these experiments, although the child was very ill with the diabetes, his mental condition, though of course far in arrear of a nine year old boy's, was steadily improving; he was interested in his surroundings, combined words and phrases into sentences logically constructed, had the reasoning powers of a child of about five years of age, was crafty in attaining his objects, quicker in his movements, and, all in all, was making progress.

At the same time, however, the diabetes was far from showing a similar improvement; it was rather at a standstill; and although the child had grown taller, he was thinner also. That the condition was constant, frequent urine examinations have proven, for the percentage of sugar at no time fell below five. His diet was only nominally restricted. At rare intervals a very minute quantity of sugar would slip into the diet; but it was found that otherwise the craving for sweets was not very marked, provided some starches and fats were allowed, as in bread or potatoes. As giving them did not seem to aggravate the trouble and certainly did benefit the patient, they were carefully given, most of the food of course being still proteid in character. This is the reason for the apparent paradox in the diet lists in the body of the paper as viewed from the standpoint of severe diabetes mellitus. To investigate this seeming incongruity, the following experiments in metabolism, imperfect as they are, were undertaken.

They were imperfect because it was impossible without a specially constructed laboratory to determine the respiratory

TABLE 1.

Date	Food	Total Quantity	Water	Protein	Fat	Total Carbohydrate	Ash	Calories per lb.	Calories in Food
Nov. 4	Bread (gm. or cc.)	326.0	126.06	30.42	3.91	169.75	3.91		865.19
	Bread (per cent.)		35.6	9.3	1.2	52.7	1.2	1,205	
	Milk (gm. or cc.)	236.0	205.35	7.79	9.44	11.08	1.65		162.5
	Milk (per cent.)		87.	3.3	4.	5.	0.7	325	
	Egg (gm. or cc.)	56.07	36.45	6.21	8.24		0.56		95.63
	Egg (per cent.)		65.	11.7	14.7		1.	765	
	Butter (gm. or cc.)	42.53	4.68	0.43	36.15		1.28		337.96
	Butter (per cent.)		11.	1.	85.		3.	3,605	
	Oatmeal (gm. or cc.)	141.75	119.78	3.97	0.71	15.66	0.99		89.06
	Oatmeal (per cent.)		84.5	2.8	0.5	11.05	0.7	285	
	Banana (gm. or cc.)	85.05	64.03	1.10	0.51	18.71	0.68		86.25
	Banana (per cent.)		75.3	1.3	0.6	22.	0.8	460	
	Beef, chopped (gm. or cc.)	226.08	153.77	47.4	24.04		2.49		417.5
	Beef, chopped (per cent.)		67.8	20.9	10.6		1.1	835	
	Potatoes, boiled (gm. or cc.)	226.8	161.23	5.67	0.23	47.4	2.27		220.0
	Potatoes, boiled (per cent.)		75.5	2.5	0.1	20.9	1.	440	
	Corn, canned (gm. or cc.)	3.54	2.69	0.10	0.05	0.67	0.03		14.21
	Corn, canned (per cent.)		76.1	2.8	1.2	19.	0.9	455	
	Soda Crackers (gm. or cc.)	113.4	6.69	11.11	10.32	82.90	2.38		481.25
	Soda Crackers (per cent.)		5.9	9.8	9.1	73.1	2.1	1,925	
	Coffee (gm. or cc.)	66.70	66.05	0.027					0.68
	Coffee (per cent.)			0.04					
	Tea (gm. or cc.)	29.57							
	Water (gm. or cc.)	1065.0							
Nov. 5	Bread (gm. or cc.)	198.45	70.45	17.45	2.38	102.62	2.38		527.19
	Chopped Beef (gm. or cc.)	113.4	76.88	23.7	12.02		1.24		208.75
	Milk (gm. or cc.)	295.7	257.25	8.76	11.83	14.79	2.07		231.25
	Butter (gm. or cc.)	56.7	6.27	0.57	47.20		1.70		450.63
	Oatmeal (gm. or cc.)	113.4	95.82	3.18	0.06	12.53	0.79		71.25
	Oatmeal Gruel (per cent.)	59.2	54.23	0.71	0.24	3.75	0.30		19.37
	Oatmeal Gruel (per cent.)		91.6	1.2	0.4	6.3	0.5	155	
	Soda Crackers (gm. or cc.)	28.35	1.67	2.78	2.58	20.72	0.6		120.31
	Potatoes, cr'm'd. (gm. or cc.)	56.7	42.58	1.17	2.70	10.09	0.85		63.13
	Potatoes, cr'm'd. (per cent.)		75.1	2.06	3.	17.8	1.5	505	
	Potatoes, b'l'd. (gm. or cc.)	28.35	21.40	0.71	0.03	5.93	0.28		27.5
	Coffee (gm. or cc.)	59.20	59.15	0.023					0.68
	Coffee (per cent.)								
	Tea (gm. or cc.)	59.20							
	Water (gm. or cc.)	1065.0							
	Bread (gm. or cc.)	170.10	59.64	15.82	2.04	88.57	2.04		451.86
	Milk (gm. or cc.)	335.27	291.68	11.06	13.41	16.26	2.35		223.41
	Egg (gm. or cc.)	85.05	55.28	9.42	12.5		8.5		145.29
	Lamb (broiled) (gm. or cc.)	170.10	80.97	36.91	50.86		2.21		624.36
	Lamb (per cent.)		47.6	21.7	29.9		1.3	1,665	
	Potatoes (c.) (gm. or cc.)	42.53	24.04	1.10	1.28	6.87	0.64		47.34
	Oatmeal (gm. or cc.)	170.10	143.73	4.76	0.85	18.80	1.19		106.86
	Apple (stewed) (gm. or cc.)	42.53	25.98	0.085	0.34	15.12	0.23		22.81
	Apple (stewed) (per cent.)		61.1	0.2	0.8	37.2	0.7	285	
	Coffee (gm. or cc.)	59.20	59.15	0.03					0.58
	Coffee (per cent.)								
	Tea (gm. or cc.)	89.0							
	Water (gm. or cc.)	473.0							
Nov. 6	Bread (gm. or cc.)	198.45	70.65	17.47	2.38	103.58	2.38		517.17
	Milk (gm. or cc.)	414.0	360.18	13.66	16.56	20.70	2.90		81.24
	Beef stew (gm. or cc.)	226.8	191.65	10.43	9.75	12.47	2.90		185.0
	Beef stew (per cent.)		84.5	4.6	4.3	5.5	1.1	370	
	Lamb chop (gm. or cc.)	141.75	67.47	30.66	42.38		1.84		520.3
	Oatmeal (gm. or cc.)	170.10	143.73	4.76	0.85	18.80	1.19		106.86
	Butter (gm. or cc.)	56.70	6.24	0.57	48.2		1.70		450.62
	Coffee-cake (gm. or cc.)	56.70	12.08	4.03	4.25	35.83	0.51		203.13
	Coffee-cake (per cent.)		21.3	7.1	7.5	63.2	0.9	1,625	
	Molasses (gm. or cc.)	28.35	1.76	1.94	2.47	21.46	0.62		119.38
	Molasses cake (per cent.)		6.2	7.2	8.7	75.7	2.2	1,910	
	Apple (raw) (gm. or cc.)	14.18	12.07	0.064	0.07	2.09	0.04		9.06
	Apple (raw) (per cent.)		84.6	0.4	0.5	14.2	0.3	290	
	Coffee (gm. or cc.)	118.40	118.35	0.05					1.16
	Coffee (per cent.)								
	Tea (gm. or cc.)	29.57							
	Water (gm. or cc.)	1063.0							
	Bread (gm. or cc.)	141.75	50.46	13.18	1.70	74.70	1.70		376.55
	Milk (gm. or cc.)	414.0	360.18	13.66	16.56	20.70	2.90		284.35
	Roast Beef (gm. or cc.)	170.10	81.99	37.93	48.65		2.21		607.50
	Roast Beef (per cent.)		48.2	22.3	28.6		1.3	1,620	
	Oatmeal (gm. or cc.)	170.10	143.70	4.76	0.85	18.80	1.19		106.86
	Butter (gm. or cc.)	56.70	6.24	0.57	48.20		1.70		450.62
	Potatoes (gm. or cc.)	56.70	42.81	1.42	0.06	11.85	0.57		55.0
	Cauliflower (gm. or cc.)	113.40	104.67	2.04	0.57	5.33	0.79		35.0
	Cauliflower (per cent.)		92.3	1.8	0.5	4.7	0.7	140	
	C-starch pud'g (gm. or cc.)	56.70	7.94		0.1	85.9			200.0
	C-starch pud'g (per cent.)		14.		0.23	10.55	0.20	1,600	
	Stewed apple (gm. or cc.)	28.35	17.32	0.06	0.23	10.55	1.28		45.62
	C-cake (gm. or cc.)	141.18	3.02	1.01	1.06	8.96			50.78
	Coffee (gm. or cc.)	59.20	59.15	0.03					0.58
	Coffee (per cent.)								
	Tea (gm. or cc.)	29.57							
	Water (gm. or cc.)	917.0							
Nov. 7	Bread (gm. or cc.)	113.40	40.37	10.55	1.36	59.76	1.36		301.24
	Milk (gm. or cc.)	443.55	358.89	14.64	17.74	22.18	3.10		101.55
	Egg (gm. or cc.)	42.53	27.64	4.98	6.25		0.43		72.65
	R. Beef (gm. or cc.)	86.05	41.48	19.19	24.61		1.12		303.75
	Potatoes (gm. or cc.)	86.05	64.62	2.24	2.58	15.32	1.29		94.68
	Oatmeal (gm. or cc.)	226.80	191.64	6.35	0.112	25.06	1.60		142.50
	Ch. Beef (gm. or cc.)	56.70	38.45	11.80	6.01		0.62		105.06
	Coffee (gm. or cc.)	59.20	59.15	0.03					0.58
	Coffee (per cent.)								
	Tea (gm. or cc.)	29.57							
	Water (gm. or cc.)	798.0							
	Bread (gm. or cc.)	212.63	75.71	19.77	2.55	112.06	2.55		564.83
	Milk (gm. or cc.)	414.0	360.18	13.66	16.56	20.70	2.90		81.24
	Egg (gm. or cc.)	42.53	27.64	4.98	6.25		0.43		72.65
	Beef (boiled) (gm. or cc.)	198.45	75.61	51.99	69.36		1.79		1228.19
	Beef (boiled) (per cent.)		38.1	26.2	34.9		0.9	2,805	
	Oatmeal (gm. or cc.)	113.40	95.81	3.18	0.06	12.53	0.80		71.25
	Coffee cake (gm. or cc.)	14.18	3.02	1.01	1.06	8.96	1.28		50.78
	Stewed apple (gm. or cc.)	56.70	34.64	0.11	0.45	21.10	0.40		91.24
	Potatoes (gm. or cc.)	56.70	42.81	1.42	0.06	11.85	0.57		55.0
	Coffee (gm. or cc.)	89.0	88.95	0.04					0.97
	Coffee (per cent.)								
	Tea (gm. or cc.)	29.57							
	Water (gm. or cc.)	1123.5							

output of the subject. With this exception, however, the results are all vouched for. The mother of the patient, an unusually intelligent woman, was easily instructed and faithfully carried out the weighing and measuring of the food and drink and keeping an accurate record of the items of the diet; she collected and measured and sent for analysis the urine passed, keeping a record of the time and quantity of the micturition. A two-ounce specimen of every urination was saved in a clean bottle, labeled with time and quantity. The rest was poured into a separate vessel and a liter of this twenty-four hour mixture sent at the same time each morning for the determination of general constituents.

Examination of Excreta.—

The stools were collected and weighed and then sent to the office, the particular feces being marked off by the ingestion of lamp black. It was necessary to give patient an aloin pill every night, as otherwise I would have gotten no stool for a week or more. As it was, stools were only passed four times during the week of experimentation. As said above, for the figures of the tests of urine and feces, I can vouch, many of them being repetitions.

Another possible source of error, which, however, is minimal, is the fact that because of lack of laboratory facilities I was unable at first hand to make the chemical analyses of the foodstuff ingested. As representative, however, of the food value and percentage of chemical ingredients of American foodstuffs I could not go very far wrong in making use of Professor Atwater's figures, and these were used to serve as a basis for my work. The averages he gives were used as being most nearly correct for the food given the patient.

In the analysis of urine, the quantitative Fehling test was used for sugar; urea was determined by the hypobromid method; the total nitrogen by the Kjeldahl method as given by Neubauer and Vogel; acetone was tested for only qualitatively by Lieben's and Legal's

1. Methods and Results of Investigations on the Chemistry and Economy of Food, Bull. 21, U. S. Dept. Agriculture.

2. Neubauer and Vogel: Analyse d. Harns.

methods; diacetic acid by Gerhardt's method; albumin, indoxyl and chlorids by the usual methods.

The tests with the feces were as follows: After weighing, they were dried with a small quantity of phosphoric acid to prevent loss of nitrogen by evaporation; of the dried feces part was used to determine the total nitrogen in a given quantity by the Kjeldahl method; fats were determined by ether extraction and carbohydrates by difference.

As is usual, the factors of Rübner³ for caloric value (4.1 for protein (6.25xN) and carbohydrates, and 9.3 for fats) were used both in the determination of income and outgo of foodstuffs and metabolic products.

In Table 1 is recorded the daily food income of the patient, the chemical ingredients of which were based, as was stated above, on the figures taken from Professor Atwater's "Bulletin on the Chemical Composition of American Food Materials." It will be noticed that the diet is now anything but strictly antidiabetic; but I insisted on the same regimen being maintained as that on which he seemed to be doing fairly thus far.

In Table 2 I have gathered the totals of Table 1 to serve as a basis for comparison with the outgo in urine and feces in Table 4.

In Table 3 are combined the results of the urinalyses. I thought it would be of interest to see if there was any cycle in the excretion of sugar and how much such depended on the ingesta. In this, however, it failed, as although the record revealed that the bladder acted at almost regular intervals and excreted in narrow limits almost the same quantity at each micturition, yet the percentage of sugar remained fairly constant.

The twenty-four hour period went from 7 a. m. to 7 a. m. and in that period there were on an average five acts of urination. The average daily amount of sugar passed was about 150 gms. and the amount of urea varied between 4.5 to 10.0 gms. daily, which falls slightly below the normal for a child of his age and weight (14.5 kgm., 32 lbs.).

The presence of acetone varied, but was fairly constant so far as symptoms went, for on the days when it was present the child acted less well than when it was absent. Diacetic acid was never present in quantity large enough to be detected; albumin was not found at any time. In other respects the urine did not deviate from the normal.

The examination of the feces gave these results. One stool was examined and the results obtained used to compute the figures for the total excreta of the week.

311.84 gm. (11 oz.) feces yielded 91.93 gm. of water-free substance. From this water-free substance, divided into weighed parts of 6.25 gm. each, there were obtained.

Total N.	Fat (ether extract)	Carbohydrates.
1.065 gm.	23.50 gm.	61.77 gm.

Based on this, the following figures were obtained as results for the week:

Protein (Nx6.25)	Fat.	Carbohydrates.
19.96 gm.	70.50 gm.	185.31 gm.
Calories—81.87	655.56	759.77

3. Rübner: *Calorimetrische Methodik*, 1891.

In Table 4 we find the metabolic balance sheet. From it will be seen the income and outgo of the protein and carbohydrate constituents of food and excreta, and also their gain and loss to the body in terms of energy—calories. It will be readily seen that the child was more than holding his own, and as Klemperer⁴ says is the case in the convalescence from acute diseases, in simple inanition, and in some forms of obesity and in myxedema, he presented a diminished outgo of protein.

Termination of Case.—On Feb. 7, 1904, I was summoned to his bedside, because of what his mother called a spasm. This yielded, however, without any treatment; but there was weakness and trembling of the lower extremities at any and all times thereafter. Twenty days later, although rational and apparently as well as ever in the morning, he went into

TABLE 2.

Food	Quantity Gm. or Cc.	Water	Protein	Fat	Carbohydrate	Ash	Calories
Bread (per cent.).....	1361.0	484.52	126.87	16.33	717.25	16.33	3,605
Butter (per cent.).....	213.0	23.43	2.13	85.		3.	1,690
Egg (per cent.).....	227.0	65.	11.7	14.7		6.39	383
Oatmeal (per cent.).....	1106.0	147.55	26.56	33.37	11.05	0.7	695
Banana (per cent.).....	85.0	84.5	2.8	0.5	22.21	0.8	86
Chopped beef (per cent.).....	397.0	934.57	30.97	5.53	18.71	0.68	750
Milk (per cent.).....	2543.0	75.3	1.3	0.6	5.	1.1	1,500
Potatoes (b.) (per cent.).....	397.0	64.03	1.10	0.51	4.37	0.7	358
Corn (canned) (per cent.).....	3.54	269.16	82.67	42.08	17.80	3.97	14
Soda crackers (per cent.).....	142.0	87.	3.3	4.	0.67	0.03	600
Oatmeal gruel (per cent.).....	59.2	2212.41	83.92	101.72	127.15	2.98	20
Potatoes (c.) (per cent.).....	184.0	75.5	2.5	0.1	3.75	0.5	205
Lamb chops (per cent.).....	312.0	299.74	9.93	0.40	19.	0.9	1,150
Stewed apple (per cent.).....	99.0	76.1	2.8	1.2	73.1	1.3	145
Beef stew (per cent.).....	227.0	2.69	0.10	0.05	0.67	0.03	185
Coffee cake (per cent.).....	85.0	5.9	9.8	9.1	73.1	2.1	306
Molasses cake (per cent.).....	28.35	8.38	13.92	12.92	103.80	2.98	120
Raw apple (per cent.).....	14.18	91.6	1.2	0.4	6.3	0.5	9
Roast beef (per cent.).....	256.0	54.23	0.71	0.24	3.75	0.30	702
Cauliflower (per cent.).....	113.0	75.1	2.6	3.	17.8	1.5	35
Corn-starch pudding (per cent.).....	56.7	138.18	3.68	5.52	32.75	2.75	200 ²⁶
Bolled beef (per cent.).....	198.0	47.6	21.7	29.9		3.06	1,228
		148.51	67.70	93.29			
		61.1	0.2	0.8	37.2	0.7	
		60.49	0.20	0.8	36.83	0.7	
		85.4	4.6	4.3	5.5	1.1	
		191.65	10.43	9.75	12.47	2.99	
		21.3	7.1	7.5	63.2	0.9	
		18.10	6.04	6.38	53.72	0.77	
		6.2	7.2	8.7	75.7	2.2	
		1.76	1.94	2.47	21.46	0.62	
		84.6	0.4	0.5	14.2	0.3	
		12.07	0.06	0.07	2.09	0.04	
		48.2	22.3	28.6		1.3	
		123.39	57.09	73.22		3.33	
		92.3	1.8	0.5	4.7	0.7	
		104.67	2.04	0.57	5.33	0.79	
		14.		0.1	85.9		
		7.94		0.06	48.71		
		38.1	26.2	34.9		0.9	
		75.61	51.99	69.36		1.79	
							13,986 ²⁷

26. Figures from Jordan's Dietary Studies in Maine State College in 1895, Bull, 37, U. S. Dept. of Agriculture.

27. When computed according to Rübner's factors, total equals 14,163 calories.

coma at 5 p. m. with almost constant alternating tonic and clonic convulsions; death occurred sixteen hours later. Necropsy was refused by his relatives.

The urine passed thirty hours before death contained 4.8 per cent. of sugar, specific gravity, 1032, slight traces of albumin, a few granular casts, some acetone and gave a very faint reaction to v. Jaksch's test for diacetic acid. Before death, in fact during coma, there were many involuntary dejections and micturitions.

In the literature of pediatrics I have been unable to find such a combination as detailed above in the case under discussion, and I look on it as unique.* For this

4. Klemperer: *Klinische Diagnostik*.

5. v. Jaksch: *Klinische Diagnostik*.

* Since writing the above, I find that Dr. Alfred Gordon, in *American Medicine*, Feb. 6, 1904, presents in a paper two cases of myxedema with genuine diabetes mellitus in brothers aged 3 and 4½ years respectively, who are the offspring of neuropathic parents with degenerative tendencies. He reaches conclusions concerning

reason it becomes more than difficult to adequately determine any answer to the questions propounded in the beginning of the paper. There is on record, however, one case, reported by C. A. Ewald,⁶ in which a woman fifty-two years old, who had developed a typical myxedema, was cured by the exhibition of thyroid gland extract, and where diabetes of a much milder form than the one I detailed above, developed afterwards. This also became a permanent affection. There was in this case also no doubt as to the affection, the patient having been presented to the Berlin Medical Society, and others, as Ewald remarks, had independently arrived at the same diagnosis.

authenticated case of Ewald which furnished an exception to his theory. Of course that it is not an every day occurrence is freely admitted; yet the possibility of its occurrence must be likewise asseverated, for what has happened once or twice may happen again.

In this case it is only logical to look at both diseases as factors in the condition, but which was the more potent is purely speculative. So far as our knowledge, much enriched in the last few years, of cretinism and analogous conditions goes, the nutritive dyscrasia caused by the lack of thyroid substance in the blood is a profound one. This, in this patient, was ameliorated and decidedly improved by the oral ad-

TABLE 3.

Date	Time	Quant- ity Cm. ³	Sugar		Quan- tity in 24 Hrs. Cm. ³	Sugar		Urea Gm.	Total N. Gm.	Acetone	Diacetic Acid	Albumin	Indoxyl	Chlorids	Specific Gravity	Calories Wasted as Ni- trogen (a) and Carbohydrate (b)	
			%	Gm.		%	Gm.										
Nov. 4	10.30 a.m.	355.0	7.8	27.69											1033	a) 251.74	902.65
	1.30 p.m.	414.0	7.0	28.98											1031	b) 650.91	
	4.00	295.7	7.81	23.09											1034		
	6.35	236.0	8.5	20.06											1035		
	8.30	236.0	9.25	21.83											1040		
	4.10 a.m.	679.5	7.24	49.20											1030		
					2128.18	7.46	158.76	6.39	9.83	Trace...	Negative	Negative	Normal.	Normal..	1035		
Nov. 5	8.10	236.0	6.85	20.06											1036	a) 211	917.
	11.00	532.0	6.66	35.43											1031	b) 706	
	1.30 p.m.	295.7	7.46	22.05											1032		
	4.20	473.1	8.19	38.75											1034		
	8.30	295.7	7.25	21.44											1032		
	11.05	295.7	7.35	21.73											1032		
					2513.0	6.85	172.14	5.026	8.25	Strong...	Negative	Negative	Normal.	Normal..	1031		
Nov. 6	9.50 a.m.	503.0	6.94	34.91											1032	a) 363	902.
	12.05	207.0	7.67	15.88											1035	b) 539	
	3.25 p.m.	266.0	7.67	20.34											1038		
	8.05	295.7	7.93	23.44											1042		
	4.35 a.m.	414.0	8.25	34.16											1038		
					1684.57	7.81	131.56	10.12	14.15	Strong...	Negative	Negative	Normal.	Normal..	1038		
																	
Nov. 7	8.15	207.0	6.94	14.37											1035	a) 190	870.
	2.00 p.m.	591.5	6.75	39.92											1035	b) 680	
	3.55	325.3	6.56	21.14											1030		
	11.55	503.0	8.06	40.54											1039		
	6.30 a.m.	561.6	8.82	49.53											1040		
					2187.20	7.57	165.50	4.38	7.42	Negative	Negative	Negative	Normal.	Normal..	1036		
																	
Nov. 8	10.00	74.2	7.35	5.45											1037	a) 209	711.
	2.45 p.m.	473.1	6.49	30.60											1030	b) 502	
	5.30	325.3	6.57	21.36											1030		
	9.15	591.5	7.04	41.64											1035		
	3.20 a.m.	295.7	6.66	19.69											1030		
					1789.50	6.85	122.58	5.37	8.14	Negative	Negative	Negative	Normal.	Normal..	1032		
																	
Nov. 9	10.15	295.7	X	X											X	a) 273	924.
	12.00 m.	325.3	6.09	19.81											1032	b) 651	
	2.30 p.m.	440.0	7.14	31.42											1031		
	4.45	414.0	6.57	27.20											1034		
	8.45	473.1	6.00	28.39											1030		
	3.45 a.m.	591.5	7.04	41.64											1034		
					2539.58	6.25	158.72	7.78	10.67	Strong...	Negative	Negative	Normal.	Normal..	1032		
Nov. 10	11.45	591.5	6.41	37.90											1029	a) 230	956.
	3.00 p.m.	493.11	7.35	34.77											1032	b) 726	
	5.30	385.0	7.57	29.14											1033		
	8.30	236.0	X	X											X		
	11.05	591.5	6.66	39.39											1032		
	10.30	503.0	5.88	29.58											1030		
					2780.11	6.37	177.09	5.56	8.96	Strong...	Negative	Negative	Normal.	Normal..	1031		
Totals...																	6182.

Under these circumstances it seems strange that Lorand, who in bibliography quotes Ewald, and Bettman,⁷ who cites Ewald's case, should make the statement that in true myxedema, diabetes was a rare if ever-noticed occurrence; in other words, casting a doubt on the well

the etiology of both diseases very similar to mine, but I can not share his optimistic prognosis even for the complication of both diseases. Thyroid extract in both of his patients showed a very favorable influence on the conjoined morbid conditions, and even earlier on the diabetes than on the myxedema, thus explaining with a great degree of certainty the rôle of the thyroid gland in the as yet unknown pathologic metabolism, which is manifested in a group of symptoms called diabetes mellitus. He thinks it highly probable that independently of the digestive functions there is a special internal secretion of the pancreas which is at fault.

6. Ewald: Nothnagel's Handbuch, vol. xxii, 1896; Berl. klin. Woch., 1895, No. 2.

7. Bettman: Berliner klin. Woch., June, 1897.

ministration of thyroid extract tablets. As pointed out before when the additional abnormal state of diabetes mellitus supervened, the child, who had because of progressive weakness not received any further thyroid treatment, nevertheless immediately not only regained all the territory lost, but even gained new so far as disappearance of symptoms and increase of intellect were concerned.

How far, then, does the occurrence of diabetes mellitus and its train of causes and effects in this cretin condition explain the working of thyroid extract in the uncomplicated myxedematous state? Is it not a fact that to a certain extent the improvement in that interesting condition following therapeutics is due to the absorption and excretion of the mucin deposited in the

skin and the various connective tissue recesses by the use of the thyroid gland's secretion? Is that therefore its real function when we administer it therapeutically? Was not nature in this case perhaps performing the same duty that we with thyroid juice strive to do, namely, removing the superfluous mucin material that the body usually uses up in other ways, by superinducing a diabetes mellitus, one of whose main effects on the system is the increased loss of all sorts of tissue and tissue food by excretion?

In his recent monograph Lorand⁸ of Karlsbad seeks to expound this interrelation of diabetes and lesions and diseases of the ductless glands (Blutgefäßdrüsen) and quite ably shows that so far as symptoms, findings at necropsy, etc., go, that diabetes mellitus, exophthalmic goiter and acromegaly are dyscrasias based on the same causation, one often running into the other or exhibiting similar clinical features in the abstract. His argument is powerful and well backed up by many able authorities, but although no doubt true in the main has this case as its exception.

TABLE 4.

	INCOME	OUTGO		GAIN
		In Urine	In Feces	
Carbohydrates.....	1389.50 gm. 5697 Cal.	1066.35 gm. 4372 Cal.	185.31 gm. 760 Cal.	137.84 gm. 565 Cal.
Protein.....	581.93 gm. 2385 Cal.	421.37 gm. 1728 Cal.	124.75 gm. 512 Cal.	35.81 gm. 145 Cal.

He says: "It is a fact that sometimes in the course of diabetes mellitus, Basedow's disease has been observed as a hyperthyroiditis (Hartman, Grube, etc.). Now the question arises, are the lesions in the thyroids or adrenals degenerative, i. e., a decreased or rather an increased activity? We would not hesitate to decide for the latter, since in atrophic conditions of these glands (as in myxedema and Addison's disease) a glycosuria is extraordinarily rare, if it has ever been observed. On the contrary glycosuria has been frequently observed in diametrically the opposite condition to myxedema, Basedow's disease, and besides very frequently subsequent to thyroid feeding."

Now there never was the least shadow of a doubt as to the diagnosis of the condition preceding the diabetes mellitus in this case. If ever cretinism was typical, in symptoms and response to treatment, this was and that the diabetes was not due to an induced hyperthyroiditis is to my mind equally established. First because the child had preliminary to the supervision of the diabetes mellitus received no thyroid extract for the period of four weeks and secondly, the fact that the ingestion of thyroid was at all times very small in quantity, and thirdly, resumption of the thyroid feeding did not increase excretion of sugar, but served only to run the child further down in weight. That there was complete athyria of course only necropsy, which was refused, would have proven; but palpation of the throat was, because he was so thin, feasible to such an extent that the cartilages of the larynx and each individual tracheal ring, the separate muscles, felt as cords, could be mapped out down to the suprasternal notch and at no place was there any thyroid or any gland tissue palpable. Under these circumstances Lorand's theory as to the genesis of diabetes needs modification. So that although the facts he adduces to support his theory are all true and powerful arguments, yet we must sug-

gest that the last word as to the exact interrelation of the various dyscrasias is even now unspoken, and we still stand rather on the knowledge adduced by Pineles⁹ "that acromegaly, myxedema, cretinism, Basedow's disease and in some cases diabetes mellitus represent diseases of glands (Blutgefäßdrüsen) in the course of which anatomic or functional disturbances in other glands supervene. There appears accordingly to be some sort of anatomic and physiologic interrelation between these glands, so that clinically when one gland is diseased, certain vital symptoms referable to disturbances of function in another gland are observable." In other words, the exact conditions are still unknown.

As a condition without such a complication, diabetes mellitus in a child is not frequent. In the literature, so far as I could find, the number of cases does not exceed six hundred; Külz reporting the collection of 111 cases; Redon,¹⁰ 32; Stern,¹¹ 117, including 21 already discussed by Külz;¹² Wegeli,¹³ 108 cases; and Saundby, 159 cases.

Various other writers, Senator,¹⁴ Baginsky,¹⁵ Niedergesäss¹⁶ and Günther¹⁷ deal with various phases of diabetes mellitus in children, but as was pointed out, nowhere are the two conditions detailed above conjoined in one patient. In fact, Scholz¹⁸ in experiments was unable by the administration of thyroid preparations to produce a single case of even alimentary glycosuria in 100 cases of cretinism. That the use of thyroid preparations in other conditions have caused both alimentary and even permanent glycosuria, many careful investigators have affirmed and reported. Bettman⁷ reported the results of experimental administration of it to 20 people who were under observation for various ailments, and in 11 (or 55 per cent.) he was enabled to produce a temporary glycosuria, which, however, totally disappeared on ceasing the exhibition of the remedy. So that he too points out its rarity in myxedema following thyroid feeding, while it is comparatively frequent as a follower of exophthalmic goiter, in both temporary and permanent form.

So, if we study the literature for this case from the viewpoint of diabetes alone, it is simply for the reason that I have been unable to find that anything else bearing on the coincidence has thus far been written. But even so, the results of the study of such literature is very interesting and instructive.

Although diabetes mellitus in a child is infrequent, it is by no means so rare as was formerly thought and taught. Sex makes but slight difference in the occurrence of the disease and although Günther¹⁷ points out the greater frequency of diabetes in female children, Saundby in his collection found the sexes about equal in numbers.

PATHOLOGIC ANATOMY.

This field also offers nothing conclusive. Great as are the strides that have been made in unearthing the truths about such nutritive disturbances as myxedema,

9. Pineles: Volkmann's klin. Vorträge, 1899, p. 242.

10. Redon: Comptes rendus Soc. de Biologie, 1877, Paris, 1879, 6me, vol. IV, p. 175; Du Diabète mell. chez l'enfant, Thèse de Paris, 1878.

11. Stern: Arch. f. Kinderheilk., vol. II, 1890, p. 81.

12. Külz: Gerhardt's Handbuch d. Kinderkrankheiten, vol. III, Tübingen, 1878.

13. Wegeli: Arch. f. Kinderheilk., vol. xix, 1896, pp. 1-65.

14. Senator: Ziemssen's Cyclopaedia, vol. xvi, p. 898.

15. Baginsky: Arch. f. Kinderheilk., vol. xxii, pp. 257-265.

16. Niedergesäss: Diabetes Mellitus Infantum, Diss., Berlin, 1873.

17. Günther: Inaugural Dissertation, Marburg, 1874.

18. Scholz: Verhandl. d. inn. Med., 1902, p. 20.

8. Die Entstehung d. Zuckerkrankheit u. ihre Beziehungen z. d. Veränderungen d. Blutgefäßdrüsen.

athyria, infantilism, acromegaly, still the last words are far from spoken. And this is most true with the condition known as diabetes mellitus. Apart from the fact of sugar being found in the urine, together with the increase of other excretory products, no one symptom or group of symptoms points out to us where to hunt exclusively for the causative lesion. Theorizing does not help us to any extent, and postmortems themselves are unsatisfactory, for many times we can find no assignable cause whatever. That head injuries, operative procedures and traumata to other parts of the body may cause glycosuria is a well-known fact.¹⁹ So, too, it can be of very small value to know that there were found at various necropsies of children dead with diabetes mellitus—free fat globules in the blood, fatty degeneration of the kidney epithelium, parenchymatous nephritis, all, it seems to me, effects rather than causes of diabetes; there were also diminution in size of the pancreas, an atrophy, or again no changes at all in the pancreas.

Such being the state of our knowledge, it makes but slight difference what theory we assume for explaining the condition we know as diabetes mellitus. Even its physiology is not settled as yet, but its pathology is only empiric. Its etiology under these circumstances is not by any means proven. Stern¹¹ in his article lays it to heredity of a diabetic or a neuropathic condition, trauma, and too free ingestion of saccharin foodstuffs; while Bergesio²⁰ ascribes it to hydrocephalus, head traumata, heredity and dysentery. In my case the too free ingestion of grapes certainly seemed the immediate cause of the condition. Of course, to a certain extent, the neuropathic tendency is also assignable from the tendency on the paternal side.

PROGNOSIS.

The prognosis in any event is very grave always, and as Senator¹⁴ puts it, "the younger the patients are the more malignant and the more rapid in general is the course;" and as Stern¹¹ has pointed out even if for many years sugar has disappeared from the urine, most of the patients die from recurrence of the malady. So far as the duration of the disease is concerned, the prognosis is far from favorable. In 32 diabetic children Redon¹⁰ found two and one-half years the longest duration of the disease; and Külz¹² found that of 46 diabetic children, 16 died in less than one-quarter year; 8 in one-quarter year; 6 in one-half year; 5 in one year; 6 in two years; 4 in three years; and one in four years. So that more than half ran their course in three months or less. "It is probable," Senator says, "that in acute cases, where from the apparent beginning of the disease until death, but a few weeks or even days elapse, which for that matter occurs almost without exception in children or young persons, the excretion of sugar and perhaps the other derangements also have existed for a long time unobserved, until a sudden change in the patient's health arrested the attention, so that in these cases, as in most others, an accurate specification of the time can not be made. In a case of Becquerel,

the disease in a boy nine years old, would seem to have lasted but six days."

SYMPTOMATOLOGY.

It is not especially necessary to enter into the symptomatology of the disease in childhood, as it does not differ materially from those symptoms seen in adult life. There are a number of phases that merit consideration, however.

Sugar in the Urine.—This most important symptom of the disease may vary from a minute but constantly increasing amount to an almost incredible quantity. Even a small percentage, one or two, when measured by the energy wasted and figured in calories, is an enormous drain, so that when such loss as Senator speaks of in a child of twelve years, and weighing 17 kgm., who passed between 358 to 570 gm. of sugar daily, is taken into consideration we can see the disadvantage nature labors under in trying to restore to normal, functions so perverted. So, too, the amounts passed by my case, ranging from 120 to 180 gm. per day, and which in the beginning of the disease far exceeded this quantity, (at one time passing 4 liters of urine containing 12 per cent. of sugar, equal to 480 gm. in twenty-four hours) show an enormous loss of fuel value to the system.

Coincident with this, Senator points out the increased output of nitrogen, as urea. "The quantity voided in twenty-four hours may amount to twice or thrice the normal, so that in adults a discharge of 100 gm. or more (142.0 gm. in a case of Dickenson's) has been repeatedly ascertained." This I was unable to verify in my patient, for if anything, with a weight of 14 kgm., he was passing less urea than was normal for such weight and age.

So far as the bases in the urine are concerned, their increase or decrease are as yet not assignable any clinical value. It is only of interest that Neubauer²¹ "saw in the urine of a diabetic child six years old, as the mean of nine days, a daily excretion of 0.71 gm. of calcium phosphate (about double the normal quantity for an adult) and 0.388 gm. phosphate of magnesia, (the normal amount in the adult being about 0.6 gm.)"

The presence of acetone and diacetic acid are of more and graver prognostic importance. The former, acetone, is of less moment than the latter, for many if not most saccharin urines contain it. Its presence causes the peculiar aroma of diabetic urine, and in larger quantities the causation of the acetonemia, found at the time of diabetic coma, is sought. Petters²² first demonstrated its presence in diabetic urine. Rupstein and later Külz¹² likewise found alcohol in the urine of a diabetic, and that too in a specimen fresh from the bladder."

I can not accept that the presence of acetone in the urine is prognostically a rapidly fatal sign, for at all times it was possible to demonstrate acetone in my patient's urine, and instead of losing he really gained ground. As for diacetic acid, even very careful testing failed to give me the typical reaction at any time, except faintly just before death.

As for albumin of any sort, at no time from the beginning of the disease until death have I been able to demonstrate its presence to my satisfaction, and microscopic examination has thus far failed to show in standing or centrifuged urine any of the typical tubular casts described by Külz¹² and by Sandmeyer²³ and on

19. Reimer (Jahrb. f. Kinderheilk., X-1-4, p. 1 and p. 219, 1876) reports a case of diabetes mellitus in a 7-year-old boy, caused by a giant-celled glioma on the floor of the fourth ventricle. The symptoms were localized cephalalgia, violent vertigo, apathy, anorexia, but great increase in thirst, with the consequent excretion of strongly saccharin urine (6.8 pounds of specific gravity 1036). Only once before this had a swelling been found at this site (by Rilliet and Barthez, *Maladies des Enfants*, vol. iii, p. 555), but there was no mention of any glycosuria.

20. Bergesio: *Jahrbuch f. Kinderheilk.*, N. F., vol. xxiii, 1886, p. 217.

21. Jour. f. prakt. Chemie, vol. lxxvii, p. 65.

22. Prag. Vjhrsschr., 1885, vol. lxxi, p. 2, and 1857, vol. lv, p. 81.

23. Sandmeyer: *Verhandl. d. Xten. Congress f. inn. Med.*, Wiesbaden, 1891, p. 345.

whose occurrence during the prodromal phenomena of coma and during it, so much stress is laid. They are described as short, mainly hyaline or even granular casts, which appeared suddenly in up to that time cast-free urine in very great numbers. They were not missed in any single case of coma examined. No casts of any sort were demonstrable in my case, except a few granular ones just before death.

So much for the urine. Other symptoms are identical with those seen in the adult. The child was at all times so ravenously hungry that he would steal any and everything in the line of food, even a cake of compressed yeast.

As will be seen by Table 2 he consumed large quantities of fluids per day, but had learned to curb his thirst enough, so that when taken out he could go for many hours (six to ten) without a drop, but making up for it on his return. So, too, the bladder was under control. Where formerly he was almost constantly going to empty it, he latterly controlled it quite well for half a day at a time if away from home.

As in the adult diabetic the metabolic processes following a full meal, caused a cerebral ischemia and a great drowsiness supervened. Even at the end of a full meal his head would go down on his arms and he would fall into a sound sleep. Contrary to the findings of Blau,²⁴ who says that in the progress of the disease diabetics become more and more indolent and averse to mental employment, this case was if anything much brighter than ever before. Now when we consider the fact that cretinoid conditions are not conducive to mental activity and this same effect follows diabetes mellitus, again the question, how far the coincidence of the two in the same personage, the one tended to modify the other, assumes further interest and force.

Of interest in this same field is the condition of the mouth and teeth. The poor, uneven and delayed eruption of the teeth in cretins is a matter of common knowledge; this was well portrayed in the case discussed. When finally, however, thyroid extract was administered the teeth erupted very fast. However, the diabetes supervened, and immediately the acid condition of the mouth, and the occurrence of thrush tended to destroy the good done. Alkaline washes, however, removed this tendency and dental attention served to put the mouth in fair condition and remove the caries which caused frequent suffering to the uncomplaining little patient.

At no time had he been a sufferer from the pyodermatosis or of any trouble of an embolic nature which so frequently accompany diabetes of an aggravated form in the adult. So also no pruritus occurred; to be explained I suppose by the need of cleanliness inculcated on the child's attendants.

So far as treatment is concerned, he received scarcely any medicine. It was found that the action of codein was objectionable; bichlorid of mercury gave no results at all, even when exhibited for a month; cod-liver oil became distasteful, and gave no return commensurate with the difficulty of administration; so that the patient, besides an occasional cathartic, took nothing but one of the newer hemoglobin preparations, which, however, seemed to serve a very good purpose. Omitting it for only a few days, the loss was immediately discernible, and a few days elapsed after its resumption before the lost ground was regained. As he held his own with

it and the fairly modified diet, no change was instituted, except that as a result of the experiments the mother was advised to give him more freely of meats, as his protein income did not seem quite adequate to his needs.

In this connection it is interesting to note that, although the dietaries of Voit, Molescott, Wolff and Playfair call for from 3,032 to 3,160 calories daily for a man at moderate work, and which Atwater²⁵ places at 125 gm. of protein and 3,500 calories of energy for the American dietary, that this patient, even under the stress of these dyscrasias, consumed only the amount of food usually figured for a child of his age (83 gm. of protein and 2,000 calories of energy per day), although, of course, as seen by Table 4, he lost the greater part of it again.

SUMMARY.

Before closing this paper and formulating the conclusions warranted from the observation of this single case, it will be profitable to revert once more to the speculative question of the interrelation of these diseases.

After Ord, Gull, Murray²⁶ and Horsley had perfected their theories and technic in the production of athyria in animals, the knowledge of the physiologic functions of not only the thyroid but of the other glandular structures, the hypophysis cerebri, thymus, adrenals, and last, but not least, the pancreas, was quickly and thoroughly deepened. The French physicians have continued their studies on the lesions of the pituitary gland and the consequent acromegaly; English physicians have presented numberless cases of Basedow's or Grave's disease and myxedema, and justly insisted on the etiologic factors being sought respectively in hyperthyroidism and hypothyroidism, although their surgical aspects assumed greatest proportions in German hands; and it was left for American investigators to probe the pancreas and its lesions as causative of diabetes mellitus.

Recent work on the interrelation of these various functions and their disorders offers nothing conclusive as yet, so that one theory has at present as much or as little *raison d'être* as another, and we must only guard against drawing the outlines too sharply, for an exception here and a criticism there may change the face of the matter very shortly. Under these conditions it may, perhaps, be well to see just what the question covers in this case.

We have seen a case which, in its sixth year, presents the unmistakable symptoms of an infantile myxedema. There is no doubt as to the condition, for besides the typical general habits, the *sine qua non*, the absence of the thyroid gland was as plainly demonstrable as it could be *in vivo*. Of course, whether this was preceded by a fetal thyroiditis, as some writers have affirmed such cases are, must be purely problematical; all we know is that great nervousness and mental strain accompanied the gestation, and it is feasible to suppose that the maternal thyroid and its use, influenced somehow the thyroid growth and function in the fetus. This point is purely theoretical and not strongly defensible. In any event, when the child was born, it was puny, with ill-developed facial expression, and was pronounced by the attendant physician, idiotic.

25. Atwater and Bryant: The Chemical Composition of American Food Materials, Bull. 28, U. S. Dept. Agriculture.

26. Murray: Diseases of the Thyroid Gland, Twentieth Century Practice of Medicine, vol. iv.

Frequent gastric disturbances supervened, although as was said, the child was breastfed. After weaning, however, when the supply of thyroid extract he got from his mother's milk was cut off, the child immediately, without any intercurrent illness, infectious or referable to the thyroid, developed a typical sporadic cretin state.

After a number of years the appropriate treatment was instituted. Small, very gradually increased dosage was employed; the mother carefully instructed to watch the least symptom of overdosage. Frequent intermissions were ordered to prevent such occurrence. The child was several times presented to the classes at the Postgraduate, and the good effect of the thyroid gland extract emphasized. At this point the child was put into my care, inasmuch as the mother felt the trip to the city too irksome and embarrassing for herself and child, and she desired treatment at her own home. She had begun to let up on the administration of thyroid, alarmed by the child's progressive emaciation, although she had been informed that this tendency prevailed when thyroid was given. At that time, as I had charge of two other cases of a similar nature, one a typical cretin, in whose urine I had found an almost pure deposit of uric-acid crystals, in such numbers and such beautiful formation, as I had ever seen before, I obtained a specimen from this patient also to determine if the same condition prevailed here. The urine at that time was normal in every respect and contained no sugar. To continue the good accomplished, small doses of thyroid tablets (0.2 gm. pro die) were administered, with frequent but unnecessary intermissions, as at no time did any tachycardia, vomiting or other hyperthyroid symptom develop.

Now, because the supply had run out, the child received no thyroid for a period of four weeks, and at the end of that time is presented with a violent glycosuria. However, the interesting discovery is made that in contrast to the accepted pictures of both athyria and diabetes mellitus, the patient presented less hebetude, is brighter, more active, his growth was noticeable, and, in fact, the supervision of the one condition served to relieve the other. Whether or not we had a vicarious action of some other gland must, of course, be purely speculative; as *a priori* must also be the condition of the other glands. We might infer an increased adrenal activity from the fact of a very small loss of blood during extraction of carious teeth; certainly we know that both his testicles were atrophied a few months before death.

But it is not warranted to go so far as Lorand⁸ does and the broader statement of Pineles⁹ must still hold good, and besides, as Ewald⁶ and, I hope, I have shown, diabetes mellitus may follow myxedema, whether you ascribe it to thyroid feeding or not. Of course, the contention that without such feeding diabetes has not been seen after myxedema, must also be admitted, as I can find no such case in an exhaustive search of literature; while it is equally true that in the administration of thyroid extract in dermatologic and general practice in cases where no lesion of the gland was demonstrable, temporary glycosuria supervened.

115 Beech Street.

Administration of Digitalis.—There is no drug which requires a more intimate knowledge and appreciation of existing pathologic conditions for its successful administration than digitalis, writes William F. Clemsha. The more carefully and thoughtfully we give it, the more uniform and satisfactory will be our results.

THE SURGICAL ASPECTS OF MAJOR NEURALGIA OF THE TRIGEMINAL NERVE.

A REPORT OF TWENTY CASES OF OPERATION ON THE GASSERIAN GANGLION, WITH ANATOMIC AND PHYSIOLOGIC NOTES ON THE CONSEQUENCES OF ITS REMOVAL.*

HARVEY CUSHING, M.D.

Associate Professor of Surgery, the Johns Hopkins University.

BALTIMORE.

The contents of this paper may be divided as follows:

1. A consideration of the various operative methods of approaching and of dealing with the ganglion.
2. Further points referable to an operative procedure heretofore described.
3. Histories of twenty personal cases thus operated on.
4. A discussion of the surgical complications and sequelæ of these cases.
5. Notes on the distribution and function of the fifth nerve deduced from postoperative observations.
6. Afterword: pathology.

Neuralgia Quinti Major. (Tic douloureux. Epileptiform neuralgia.) This form of neuralgia has been placed in a group apart owing to the marked similarity that one case bears to another both in course and symptoms. In the earliest stages, each case appears to be one of simple neuralgia, and under this impression relief has been sought in removal of teeth or in other operative procedures. At last the full picture of this hideous disease unrolls, varying in unessential details from the classical type.

In a recent medical treatise¹ thus is introduced the section on that form of trigeminal neuralgia which, unlike the other varieties, the minor neuralgias of the fifth nerve, is an affliction demanding surgical intervention, and for the relief of which the removal of the semilunar ganglion (Gasser) must ultimately be contemplated. It is, perhaps, deserving of emphasis, therefore, at the start, that from this group of cases are excluded the comparatively simple neuralgias arising in the trigeminal territory; those which are incidental to some general state of bodily ill health; those reflexly occasioned by some local disease; those ordinarily confined to one or another of the peripheral branches of the trigeminal nerve, all of which are usually amenable to therapeutic measures² other than surgical or, at the outside, to minor operative procedures.

Barely thirteen years have elapsed since Horsley and Macewen,³ quite independently, and, so far as I am aware, for the first time, attempted through a high temporal opening to divide the trigeminal nerve in two cases of tic douloureux. Both resulted fatally, an

* Presented by invitation at a meeting of the Montreal Medical Society, 1904.

1. Head: Trigeminal Neuralgia. Allbutt's System of Medicine, 1899, vol. vi, p. 728.

2. At times very simple measures indeed, of which there is an interesting historical illustration. In one of the old numbers of the European Magazine are reprinted a series of urgent letters which John Locke, the philosopher-physician, wrote from Paris to Mapletoft and Sydenham asking for advice in the treatment of the distinguished patient, the Countess of Northumberland, whom he was attending at the French Court. Owing to "the inclemency of the weather," Locke had refrained from following his first inclination of opening her ladyship's bowels. With a moderation in the temperature and before Sydenham's reply could have been received a purge was administered and the patient's neuralgia disappeared. There is an amusing reference by Lord King in an early number of The Lancet to this correspondence, which comments not only on what must have been the inconveniences of Parisian life in the latter part of the seventeenth century, but also on Locke's clear statement of the case and his skillful treatment.

3. Victor Horsley: An address on the Surgical Treatment of Trigeminal Neuralgia, The Practitioner, 1900, vol. lxx, p. 251.