normal function, and the power of speech returned. Upon discontinuing the treatment nocturnal headache and delirium reappeared, whereupon three more baths were given, after which all symptoms of disease vanished. Aufrecht reports this single observation to encourage the further trial of the very favorable effects of hot water.

FREEMAN.

Treatment of Tubercular Meningitis.—By Dr. R. Hirschberg. (*Bull. Gèneral de Thérapeutique*, November, 15, 1894.) According to the author, death in cases of tubercular meningitis is caused not by the development of the tubercles but by intracranial compression, by cerebral asphyxia. The rational treatment of the disease lies, therefore, in trephining and drainage, procedures similar to those employed with good results in tubercular peritonitis. The operation should, however, be performed before coma sets in. Ord and Waterhouse report a case of tubercular meningitis in a child five years old, in which trepanation of the occiput was done six weeks after the onset of the disease. The dura and arachnoid were divided, and a probe and drainage tube introduced between the cerebellum and the medulla oblongata. The drainage was followed by an amelioration of the severe symptoms. In spite of the fact that the tube was kept in situ for seventeen days, and an intercurrent attack of measles developed, the child got well.

MEIROWITZ.

On the Analgesic Properties of Guayacol.—By Aporti. (*Gaz. d. asp. Deutsche Med. Zig.*) In a number of cases of pleuritis, pneumonia, gastralgia, acute articular rheumatism, myelitis, dysmenorrhœa, tubercular meningitis, neuralgia, neuritis, etc., the topical application of mixture of equal parts of glycerine and guayacol, two to three times daily, produced prompt and lasting effects. There are, however, contra indications for its use, as the presence of a high fever: nor is it always borne well, as cauterization and neurosis of the superficial structures of the skin may follow its application.

MEIROWITZ.

The Operative Treatment of Spasmodic Torticollis.—Richardson and Walton. (*American Jour. Med. Sci.*, January, 1895.) After reviewing the therapeutics of spasmodic torticollis the authors reach the following conclusions: 1. Palliative treatment, whether by drugs, apparatus or electricity, will rarely prove successful in well established spasmodic torticollis. 2. Massage may prove of value in comparatively recent cases. 3. Resection affords practically the only rational remedy. 4. Operation on the spinal accessory nerve may afford relief, even if other muscles than the sterno-cleido-mastoid are affected; on the other hand, the affection previously limited to the sterno-cleido-mastoid may spread to other muscles in spite of this operation. 5. No fear of disabling paralysis need deter us from recommending operation, as the head can be held erect even after the most extensive resection. The most common combination of spasm is that involving the sterno-mastoid on one side and the posterior rotators on the other, the head being held in the position of sterno mastoid spasm with the addition of retraction through the greater power of the posterior rotators. 7. It seems advisable in most cases to give preference to the resection of the spinal accessory as the preliminary procedure.

J. C.