the white corpuscles had been decreased to 93,750, or 1 to 23.

Today his hemoglobin equals 63 per cent., his red corpuscles 4,000,000 and his leucocytes only 19,000. The proportion therefore is 1 to 210, which approximates the normal proportion very much more closely than the earlier count. The blood also shows that the myelocytes are present in the proportion of from 8 to 13 per cent., and it is a noteworthy fact that his polynuclear leucocytes, which are usually decreased in this disease, amount to from 51 to 66 per cent., which is about the number usually found in normal blood. His spleen is found to be greatly enlarged. It extends almost to the anterior superior spine of the ilium and in the middle line as far as the umbilicus, but his color has greatly improved and there are many men who would appear far more anemic than he, although they would consider themselves as being in perfect health.

The second case is far more interesting because much more rare, representing as it does a case of possible lymphatic leukemia. It is that of a young man of 23, a barber by occupation, who is intensely pale and yellow looking. An examination of his chest and arteries shows loud anemic murmurs, his spleen is enlarged to even greater extent than the patient first brought before you and we have records of his blood over many months. October, 1895, an examination of his blood showed that his hemoglobin was 30 per cent., his red blood corpuscles 2,060,000 and his leucocytes 8000. He has been in hospital off and on since that time and we have a great many counts of his blood. His case is not a typical one, but I think we can say that it is one of lymphatic leukemia. At the present time his hemoglobin is 33 per cent., his red corpuscles the same as before and the leucocytes 6200. While the red corpuscles are somewhat decreased, as they frequently are in leukemia, the white blood corpuscles are not increased as is usual; indeed, this proportion between white and red is often seen in health. It is only when we consider his various corpuscles that we have much light thrown upon the diagnosis. We then find that his lymphocytes, which are ordinarily present in comparatively small numbers, amount to 84 per cent., that polynuclear leucocytes, which usually make up a large proportion of the white cells, only amount in this patient to 8 per cent., while the mononuclear leucocytes amount to 4 per cent. There is also absent in his case one other frequent sign of lymphatic leukemia, namely, very great enlargement of the lymphatic glands all over the body. This boy has no such enlargement, but his lymphocytosis, combined with his profound anemia, the fact that he has had hematuria, that he has hemorrhages in his retina and that he is intensely anemic and pallid without a corresponding loss of flesh, seem to make the diagnosis fairly clear.

In many of these cases of leukemia we find in addition to a decrease in the number of red corpuscles, an abnormal red blood corpuscle which possesses a nucleus, this corpuscle being called a normoblast and sometimes also possessing ameboid movement. In the presence of a patient suffering from multiple enlargement of the lymphatic glands, a symptom which this boy might present, it would be incorrect for you to jump to the conclusion that he was suffering from lymphatic leukemia, because a similar enlargement is seen in what is known as Hodgkin's disease or pseudo-leukemia and multiple tubercular infection.

The treatment given these patients has been a

nourishing diet, the meeting of current symptoms which have seemed of most importance, and ascending doses of arsenic until they reached the physiologic limit of the drug.

FIVE CASES OF CHOLELITHIASIS.

BY BAYARD HOLMES, M.D.,

PROFESSOR OF SURGERY IN THE COLLEGE OF PHYSICIANS AND SURGEONS OF CHICAGO, THE MEDICAL SCHOOL OF THE UNIVERSITY OF ILLINOIS.

The following cases are reported because each of them contains at least one point of real surgical interest:

Case 1.—*Indistinct pain in the abdomen, with sick headache; the morphin habit; empyema of the gall bladder; cholecystostomy; recovery.* The patient was a physician, 45 years of age, who came to Dr. Waugh's sanatorium for the treatment of the morphin habit. He was of very regular habits, used no tobacco or alcohol and had no significant previous history. He had indistinct pain in his back and abdomen for several years and was accustomed to take small doses of morphin to relieve this distress. These doses increased, however, in frequency and size until he found himself, under one pretext or another, taking larger and larger doses, up to 0.45 or 0.52 grams a day. He was bright and had been able to follow his practice, which was large and diversified. On coming to the sanatorium he stood the treatment well for a week or two, when without apparent cause and without any very great increase in this indistinct pain in his back he gradually began to have an evening rise of temperature, with some sweat. At length symptoms of peritonitis, distension of the abdomen and tympanites appeared, and it was discovered that his right abdominal region was filled by a tumor six inches long and three or four inches wide. It was so large a tumor that it was at first looked upon as a peri-appendicular abscess. In the course of twenty-four hours the symptoms were so grave that I was called in to open the abscess. This was Dec. 16, 1893. The patient at this time was only partially conscious; his abdomen was slightly tympanitic; the bowels had been freely moved by an enema; the patient's temperature was 102, his pulse 120. He had never had any paroxysmal pain, but complained of a deep pain in the back, which was relieved by the use of morphin. In examining the eyes the pupils did not show the effect of the morphin; the tongue was coated typhoid-like; the lips dry and parched; the skin hot and dry; the respirations rapid and superficial, as in peritonitis; the heart's action was perfect, but rapid; the abdomen was slightly distended, and in the right side was found a tumor, round, smooth and pressing up against the abdominal wall, which tumor I readily recognized as a greatly distended gall-bladder or possibly a pyonephritic tumor. His urine was examined and found to contain a trace of albumin and was otherwise highly concentrated. He urinated voluntarily. The case was considered one of great gravity, and an immediate operation was undertaken. The abdomen was opened by an incision along the outer border of the right rectus muscle two and one-half inches long. The tumor showed itself in the wound unattached to the viscera. The first effort, however, to examine it resulted in rupture of the sac and the discharge of a large quantity of thick pus. The walls of the tumor were instantly grasped in several pair of forceps and the tumor pulled out into the wound. By means of several silk sutures, it was attached to the margins of the wound and the edges were trimmed off, making a large opening into the gall-bladder. The finger was passed down to the cystic duct, but on account of the necrotic condition of the sac it was considered unadvisable to make any attempt to sound the cystic duct.

During the following twenty-four hours the temperature fell to 99 and rose again at the end of the second day to 101, after which it gradually declined until the twenty-fifth day. Upon the eighth day the stone appeared at the mouth of the wound and was with some difficulty removed. It was half an inch in diameter and about an inch long, being pear-shaped and having a facet at one end, showing that another stone had been in contact with it. This was dislodged a few days later. The patient made an uninterrupted recovery, with one exception. On the tenth day after the operation the usual irrigation of the wall of the gall-bladder was undertaken by the nurse, but the mouth of the fistulous opening into the gall-bladder had now become so small that the irrigator point did not allow a free return of the water from the cyst, and as a result the water probably flowed through the cystic into the hepatic duct and carried back with it the pus from the gall-bladder. The patient complained of a sudden dreadful pain, quickly went into a collapse

from which he was with difficulty revived, and in the course of a few hours he had a pronounced chill and a rise of temperature to 103 degrees, as represented in the chart. This was unaccompanied, however, by any symptom of peritonitis, and upon the following day the temperature fell to normal and never rose after that time. This is shown by the accompanying chart. In the course of a few weeks the fistula completely closed, but it required a secondary operation to remove one of the silk sutures, which had been too hastily placed during the emergency of the first operation in an inaccessible position.

This case illustrates the fact that a stone may give rise to severe disturbance and nervous unrest and distress which can not be characterized as biliary colic, and may even initiate the morphin habit in an otherwise healthy and well-balanced man. It also shows the danger of irrigating the gall-bladder after the obstruction in the cystic duct has passed away. In this case the irrigations were begun on account of gangrene of a large part of gall-bladder, but I thoughtlessly neglected to give directions that the irrigations be stopped as soon as the stone was dislodged and the bile began to pour out.

Case 2.—Impaction of a stone in the cystic duct for five years, with attacks of pain and vomiting at intervals of two or three months, each lasting a day or two. Relief from all symptoms for five years. Impaction of the stone in the middle of the common duct; rupture of the common duct, removal of the stone, suture, drainage and death from shock. Mr. P. when I first saw him was 51 years old. He was suffering from an attack of "gastritis," the symptoms of which were pain at the pit of the stomach, general abdominal distress, persistent vomiting and considerable prostration. This attack came on the morning after Thanksgiving, presumably after an unusually full meal. He said he had had such attacks repeatedly for three years. Usually it had been necessary to give morphin to remove the severe, rather constant pain, which was always felt slightly to the right of the middle line of the abdomen and just under the edge of the ribs. The morphin was given hypodermically and the patient carefully examined. A small round tumor the size of a hen's egg could readily be felt almost touching the abdominal wall in the region of the gall-bladder. The liver was not enlarged, the stools were normal colored, there was no jaundice. A diagnosis of a stone in the cystic duct was made and an operation for its removal was recommended. The next day the pain had disappeared, the patient recovered during two or three days on a light diet, and an operation was refused. During the following three years this happened at irregular intervals two or three months apart, one of which always occurred on the morning following Thanksgiving. I attended him through each of these attacks and always recommended cholecystostomy. The patient submitted to a rather restricted diet and during the five following years he had no attacks of pain and often spoke of his wisdom in refusing an operation. At last, upon the morning after the eighth Thanksgiving he sent an imperative message to bring him morphin as he was suffering from another attack of gall-

stones. This time the pain was much more severe than ever before. The patient showed jaundice six hours after the beginning of the attack, which jaundice deepened gradually during the twelve hours following while we were waiting for the return of his wife in order to operate upon him. It was almost impossible to remove the exacerbations of pain, which came on first at 4 A.M., again at 7 A.M., again at 7 P.M. and lasted from this time continuously until he was operated upon the next morning at 11; but vomiting occurred only two or three times, and nausea was not a prominent symptom. At first a hypodermic of 0.03 gram of morphin was given. Six hours later a teaspoonful of chloroform in olive oil was given by the mouth. After this the pain was controlled by inhalations of chloroform at each paroxysm. His wife returned twenty-nine hours after the beginning of the pain. At that time his pulse was 100, temperature 98 in the rectum; his abdomen considerably distended and his skin in the second stage of jaundice. The abdomen was opened by a small incision in the median line high up, and immediately about a quart of greenish fluid poured out. The finger discovered a completely atrophied gall-bladder, and a stone an inch long and half an inch in diameter in the common duct about its middle portion. This stone had the appearance of two truncated cones set with their bases together. Its surface was dark and nodular. There was evidence of the former contact of a small stone at one end. A long incision was then made on the outer border of the right rectus muscle, the abdominal cavity was carefully washed out with a large quantity of sterile water. The liver and intestines were retracted with the assistants' fingers and a row of twelve fine silk sutures were placed in the wall of the common duct over the perforation, which was found directly over the stone in the middle and beginning lower third of the common duct. The stone was used as a guide in inserting the sutures. When the sutures had been placed a forceps was carefully insinuated into the opening and the end of the stone was grasped, and by the aid of the finger behind it and the forceps it was slowly and carefully withdrawn without much enlargement of the rupture. The duct was explored in both directions without result. The sutures were then drawn together and tied and cut off short. There seemed to be no leaking, and yet fearing this possibility, the omentum was brought up and fastened so as to make a deep well leading down from the long abdominal incision to the rupture in the common duct. This well was filled with strips of iodoform gauze in Mickulicz's handkerchief, and in order to make the drainage more complete a large perforated drainage tube was passed down the center of the gauze for a distance of five inches. Everything was held in place by a few sutures in the omentum, and the abdominal wall was closed, allowing about two inches space for drainage. The operation lasted an hour and a quarter. The patient seemed to suffer only moderate shock, the pulse and temperature at the end of the operation were only a trifle different from what they were at the beginning. During the eleven hours that the patient lived the symptoms of shock increased and the patient slowly but surely, and in spite of every means of stimulation and care, declined. There was no hemorrhage. The dressings were saturated with a greenish fluid; the urine was high colored and the perspiration yellowish. The patient died eleven hours after the operation without reaction. Superficial post-mortem examination through the wound revealed nothing new.

This case illustrates the danger of neglecting to open the gall-bladder and remove the calculus as soon as the diagnosis has been made. It is also a warning against neglecting to operate in case of obstruction of the common duct at as early a moment as possible. Although many cases of traumatic rupture of the gall-bladder have been reported in which recovery took place; after rupture, especially when due to obstruction by a gallstone and presumably accompanied by infection of the biliary apparatus, death from peritonitis is the rule. Mayo of Rochester, Minn., reports a case of injury of the gall-bladder in a boy 12 years old, with recovery after repeated removal of a biliary ascites, and many similar cases are reported. Peritonitis is produced not by bile but by infected bile.

Case 3.—Obstruction of the common duct, absence of the gall-bladder; disappearance of the calculus under manipulation; drainage of the peritoneum; recovery. In August, 1894, Dr. Louis May called me to see a patient, 35 years old,

in whom he believed he had a case of obstruction of the common duct. The patient was a powerful, fat ice-man, of rather intemperate habits, who had suffered from frequent attacks of colic unaccompanied by jaundice. Three days before my observation he had been attacked with colic during his labor distributing ice. He administered to himself liberal doses of whisky, both hot and cold, and had one or two doses of morphin during the night hypodermically. The pain was very severe and the paroxysms were of rather long duration. In the morning he was completely jaundiced. The jaundice increased, the urine became black and the patient went into a comatose condition, with tympanites and occasional attacks of vomiting. This was the condition in which I found him. His jaundice was extreme; the clothing in which he lay was stained with bile. The urine removed from his bladder with a catheter was almost black. The respirations were shallow and the heart beats were faint. The pulse was 110 and the temperature 98. No tumor could be felt. The liver was considerably enlarged, extending downward about an inch and a half beyond the margin of the ribs. Operation for the removal of the obstruction was undertaken. The stools had been removed during the last two days by means of enemas and they were uncolored by bile. The patient was anesthetized with a small amount of chloroform and an incision made along the outer border of the right rectus muscle. The intestines were stained yellow, the liver itself looked almost black, so greatly was it engorged. The space in which the gall-bladder is usually found was occupied by a firm cicatricial mass, and the gall-bladder seemed to be entirely absent. Passing the finger down the common duct, at the first examination it seemed that a row of calculi could be felt, but a few minutes later and after a little further manipulation made to break up some light adhesions over the common duct, they entirely disappeared, nor could they, during the course of the operation, be felt again. At this time a greenish-yellow fluid was sponged out of the bottom of the abdomen and I was afraid, though I could not be certain, that I had ruptured the common duct. Pressure upon the liver caused distention of the common duct, and as soon as the finger was pressed upon it to milk it into the intestine the duct collapsed and the bile disappeared somewhere, presumably into the duodenum. I believed that the obstruction had in some manner been removed, and I feared that the common duct leaked into the peritoneal cavity. The stones could not again be felt; I did not think best to open the common duct and probe for them. Therefore, I built a sort of well leading down on the under surface of the liver between the right lobe and the quadrate lobe and paved the space with strips of iodoform gauze and sewed up the remainder of the wound, covering it all with an antiseptic dressing, held on with adhesive straps. The patient did not vomit after the anesthetic, and on the morning of the following day he was perfectly conscious, had a good appetite and several free bile-stained movements of the bowels. The wife made a superficial examination of the feces, but did not find any stones. This examination I consider valueless, however. The jaundice began to disappear. Upon the third day after the operation he resisted his wife's entreaties and the doctor's orders and dressed himself and went to his favorite saloons to see his friends. The doctor was called and induced a police officer to assist him in taking the patient home and putting him again to bed. The dressings were at first saturated with bile, but after a day or two the discharge was small, the packing was gradually removed and the wound closed without any event. The patient has not had an attack of biliary colic or jaundice since that time and has during the past four years been perfectly well.

This case illustrates the fact that after long continuance of calculi in the cystic duct the gall-bladder sometimes becomes atrophied, and it further illustrates the fact that the stones may be upon the verge of passing into the duodenum and need only slight assistance to dislodge them. It also shows that after forty-eight hours the adhesions about a Mikulicz's drain in the abdominal cavity are strong enough to withstand considerable assault. It seems to me now that in a similar case I should not be satisfied to leave a stone because I could not palpate it.

Case 4.—Repeated attacks of biliary colic for nine years, without jaundice; cholecystostomy and immediate removal of the calculus; recovery.—Mrs. P., 46 years old, the mother of five children, the youngest one 9 years old, had puerperal fever after the birth of this child. The symptoms were very grave. The patient had chills and fever and sweat for six or seven

weeks and recovered with broken health, very much emaciated. At the same time she had an abscess of one of her breasts, which required poulticing and afterward incision and drainage. When this child was 1½ years old she had an attack of colic, which lasted for three or four days, during which there was very severe vomiting and great prostration. From this attack she recovered promptly and since that time has had such attacks at frequent intervals, usually as often as two or three times a year. I was called to see her Feb. 13, 1896. She had been suffering for twenty-four hours with excruciating pain in the upper right side of her abdomen, extending into her back and right shoulder. A physician who had been treating her had made a diagnosis of gallstones and had attempted to administer remedies by the mouth, but on account of the frequent and persistent vomiting these remedies had had no effect. The pain was unabated and the vomiting occurred at intervals of ten to twenty minutes, with great retching. I found a large fat woman, with relaxed abdominal walls, moist skin, tender abdomen and a considerably contracted pupil, probably the effect of the morphin. There was no jaundice, and the liver was not recognizably enlarged. The urine was of normal color and contained only normal constituents. The temperature was between 100 and 101 degrees F. There had been no pronounced chill. The pulse was strong but rapid, 110. The patient looked anxious and begged for relief. I made a diagnosis of a single large stone in the cystic duct, and in trying to explain the condition to the husband I made a drawing of what I believed the stone to be, both in size and general appearance. This drawing he afterward compared with the stone which he saw removed. I administered a hypodermic of morphin, 0.03 gram, and urged an operation, basing my diagnosis wholly upon the symptoms. The abdomen was opened by a short incision in the median line and the finger pressed downward and to the right discovered the stone and a slightly distended gall bladder. A second incision was made along the border of the costal cartilage opposite the ninth rib, and the tumor was with some difficulty pressed up into the wound and fastened there with a couple of rows of strong silk sutures. The gall bladder was opened and the stone in the cystic duct was easily pressed forward into the gall bladder and outward through the wound. The stone corresponded almost exactly in size and shape with the sketch I had made of it before the operation. The wound was covered with antiseptic dressing, and during the following three weeks it continued to discharge more or less bile, when it closed suddenly and permanently. Since this time the patient has been wholly free from these attacks and has had no symptoms of sick headache.

This case shows the probable source of gallstones in the puerperal fever which the patient suffered a year and a half before any pain was felt. It also shows how long a patient is frequently neglected, suffering from stone in the cystic duct, a place so easily attacked and so amenable to simple surgical treatment.

Case 5.—Sudden impaction of stone in the cystic duct without previous symptoms of biliary colic or cholangitis, cholecystostomy and discharge of a large number of stones, with a single copious discharge of bile after ten days; closure of the fistula in two months, followed in the course of three weeks by renewed colic; second opening of the gall bladder; drainage for three months; recovery.—Mrs. E., aged 54, the mother of two healthy children, had never had any attacks of biliary colic or symptoms which could be referred to this condition. April 26, 1897, she was suddenly taken with a severe pain in the epigastrium, with vomiting and very slight indication of jaundice. The pain continued for twenty-four hours, with a chill at the end of this time and rise of temperature to 102. This was followed by repeated slight chills and a continuous high temperature for three days, when I was called to see the patient. She was a large woman of dusky skin and dark eyes. The abdomen at this time was excessively tender and in the region of the gall bladder especially sensitive, but by careful rolling of the tissues under the palm of the hand, a round smooth tumor could be felt, which was believed to be the distended gall bladder. There was nothing significant in the condition of the heart and lungs. The respirations and pulse corresponded with the temperature, which was now 102. The patient was much collapsed and begged to be relieved of the pain and vomiting. I recommended immediate operation, designed to relieve the gall bladder and remove the obstruction in the cystic duct.

An incision was made in the middle line large enough to admit the index finger, and the diagnosis of obstruction of the cystic duct and empyema of the gall bladder was confirmed. The gall bladder was surrounded on nearly all sides by light

adhesions, which did not, however, involve the anterior abdominal wall. An incision was made over the gall bladder and with some difficulty it was drawn up into the incision and fastened there by a row of silk sutures. The gall bladder was then opened and a cloudy whitish yellow fluid was discharged. The finger was passed into the gall bladder, but no stones could be found by a light examination. During the following ten days the patient rapidly improved, and a number of faceted gall stones were discharged. About this time the dressings were suddenly saturated for the first time with bile. A few more stones were then discharged into the dressings and the patient improved, but no more bile showed itself. July 1 the wound had completely closed, the patient was about and feeling well. July 26 the patient was again attacked with the same symptoms which she had suffered in April. I saw her at this time every day. On the third day I determined to reopen the gall bladder in the scar. The patient was as much prostrated as she had been before the first operation, and I determined now to keep the gall bladder open until I was certain that all stones were removed or dislodged. After this operation the gall bladder was drained until November 1, during all of which time there was a discharge of bile whenever the opening from the gall bladder was not plugged. After free drainage for two or three weeks without the removal of any stones, I introduced an olive-shaped urethral sound (about No. 15 American in size) into the gall bladder, and coiled up the silver handle to it so as to make a sort of spring resting against the abdominal wall, thus pulling the blunt side of the olive into the wound to act as a ball valve. This remained in the wound from the last of August until November 1, when it was removed. During all this time there was only a small and not very troublesome discharge of bile. The sound itself on removal was covered with the thinnest possible black film, which cracked, rolled up and fell off as soon as it had dried. I can not account for the obstruction which occurred in July. I was not able to find in the gall bladder, or in the cystic duct, or in the common duct any evidence of stone, although I was able without much difficulty to pass a sound into what I believed to be the intestine. During the past six months the patient has been very well and able to attend to her many duties.

Each of these cases has its own story to tell. In only one case did death result, and then because the warnings of biliary colic without jaundice were unheeded, and perhaps, too, because the necessities of the case almost demanded too great a delay after the last warning given by biliary colic with jaundice.

A CASE OF TUBERCULOSIS OF SPLEEN, WITH SURGICAL TREATMENT.

BY A. M. HAYDEN, M.D.
SURGEON TO ST. MARY'S HOSPITAL.
EVANSVILLE, IND.

A scarcity of literature on the subject of surgery of the spleen has induced me to report the following case, one that has been of unusual interest to me.

About Feb. 20, 1896, Mrs. Wm. F., age 24, first felt pain in the left hypochondriac region, which, slight at first, gradually grew more severe until she came under my observation, April 19, 1896. At that time the pain in the region of spleen was at times quite intense, while at others it was entirely absent. There was no rise of temperature, but her pulse was about 100. Her appetite was good, she had no cough or bowel trouble, and attended to her household duties as usual. I prescribed counter irritants externally and quinia and codein internally. Her condition remained about stationary until May 10, when her temperature was 101 degrees and pulse 120. The pain in the spleen was quite severe and on palpation I found quite an enlargement of that organ. She had no chills preceding the rise of temperature, nor other symptoms of malaria. She was well nourished, had rosy cheeks and weighed 108 pounds, which was her average weight.

Notwithstanding she had no symptoms of malaria, I put her on quinin, Fowler's solution and syrup of the

iodid of iron, with tincture of iodine externally, and kept her on a similar line of treatment for fifteen or twenty days, during which time her condition continued to grow steadily worse. I was still in doubt about my diagnosis and called council. After a thorough examination, which included the history of both her father's and mother's families, with negative results in regard to both tuberculosis and carcinoma, we were still in doubt about the nature of the trouble that was producing the tumor in the spleen, as it had now reached a size that would justify us in calling it a tumor. My council advised a trial of protonuclein, and it was given her for about two weeks. Meanwhile the tumor of the spleen continued to enlarge, forcing its way up under the short ribs and extending down into the left iliac region and over to the median line, reaching to the umbilicus.

An exploratory operation was done June 23, 1896. The incision was made over the most prominent part of the tumor. After reaching the peritoneal cavity we found the tumor free from adhesion to the abdominal walls.

It was softer than normal spleen structure and of a grayish color. I opened the splenic cyst or covering, which is composed of peritoneum and a fibrous coat intimately connected. Inside of the cyst we found a large mass of abnormal growth and broken down splenic tissue. After removing this soft tissue, which I did with my fingers and a spoon curette, I stitched the splenic cyst to the wall of the abdomen, leaving a cavity in the cyst as large as a baby's head, with no cystic wall separating it from the small portion of the splenic structure that was not removed.

I thoroughly packed the cavity with iodoform gauze and dressed the wound open to secure drainage, thinking that suppuration would take place and it would heal by granulations. A microscopic examination of the specimen removed blasted all such hopes, as it revealed pus corpuscles and tubercular bacilli.

The patient rallied well from the operation. Peritonitis developed on the second day, threatening to destroy the patient's life; after a few days she gradually improved.

Her pulse rate was unusually high compared with the temperature. A chart was kept for twenty-two days after the operation, from which I found an average morning pulse of 114.2, the highest rate being 150, which was noticed June 26, and the lowest 85, July 8. There was an average evening pulse of 119½. The highest reached was 144, June 27; the lowest, July 5, 98. During the twenty-two days there was a mean morning temperature of 99.5 degrees; the highest point being reached on the morning of June 26, 103 degrees; and the lowest July 8, 98 degrees. There was a mean evening temperature of 100.2 degrees; the highest on the evening of June 25, at which time it reached 104 degrees. The lowest evening temperature was 98.2 degrees, and occurred on the evening of July 14, 1896. The high pulse rate continued throughout the entire course of the disease. This high pulse rate has been a peculiarity of most cases of disease of both the spleen and pancreas that have come under my observation.

The patient gradually improved in her general condition as indicated by a good appetite, marked gain in flesh, weighing more three weeks prior to death than she had for several years previous. The opening into the spleen continued to discharge a small