The Journal of the American Medical Association

VOL. XXII.

CHICAGO, FEBRUARY 17, 1894.

No. 7.

ORIGINAL ARTICLES.

THE SCOPE OF DERMATOLOGY.

Chairman's Address read in Section of Dermatology and Syphilography,
at the Forty-fourth Annual Meeting of the American Medical
Association.

BY LOUIS A. DUHRING, M.D.

PROFESSOR OF DERMATOLOGY IN THE UNIVERSITY OF PENNSYLVANIA.
PHILADELPHIA.

On the occasion of the last annual meeting of this Section of the Association, the honor to preside over the deliberations of the present meeting was conferred upon me. It affords me pleasure to greet you as members of the Association, and especially as attendants upon the work of this Section, whose importance I believe must be more fully recognized by the profession with each succeeding year. Before taking up the original papers to be presented for reading and discussion I would ask your attention for a few moments to consider of what dermatology actually consists. The subject, I believe, will not prove uninteresting nor without profit practically. Let me propound the questions, What are so-called skin diseases, and what is their nature? Which diseases are entitled to be designated "skin diseases?" Is it possible to separate them from the many general diseases accompanied by cutaneous symptoms?

Answers to these questions may at first thought appear easy, and so they seemed to the writer many years ago, when thoroughly imbued with the spirit of local pathology, as taught by his former and respected master Hebra, it was deemed all-sufficient to study cutaneous diseases purely objectively. But the questions propounded upon investigation really became difficult problems, especially with the light that has been thrown upon dermatology by comparatively recent studies. The field, viewed in every direction, has become immensely enlarged, and the amount of work done, both clinical and pathological, has been remarkable. I believe, however, that a satisfactory solution of these problems may be found. The questions that immediately confront us are, shall the term, "skin disease," continue to be used in its ancient sense? Shall the affections of the skin be looked upon as morbid entities, as diseases whole and complete in themselves and confined to the skin? Or, taking a broader view of the subject, shall they be defined so as to include all manifestations that may occur upon the integument irrespective of the symptoms, cause or nature.

The first and restricted proposition is that according to which these diseases were studied and classified a century ago, by Plenck in Germany and by Willan in England, when the list of diseases was small and knowledge of them meager. I have no desire to belittle the work done by the fathers of dermatology. Willan, especially, was an eminent general physician as well as a distinguished dermatolo-

gist. His treatment for the commoner affections of the skin was both judicious and successful. But the important questions of etiology and pathology were then for most diseases not at all understood. Very little was known on those subjects. The origin of most disease was regarded as "obscure." The facts that a mite was the sole cause of scabies, and that ringworm was due to a fungus were still unknown, while the observation that almost all the drugs in the "Pharmacopœia" are capable of producing disorder of the skin was not even suspected. Affections of the skin were regarded mainly objectively, and were studied much as a model or a picture might be viewed. Beyond the actual expression on the integument they were not closely investigated; the causes were regarded as obscure; and this mode of studying them has held good through decades, up to almost the present date. This method is in no way to be criticized, so far as it goes, but by itself and without the assistance of general pathology, it is far too restricted to meet the requirements of existing knowledge. It fails to recognize the important fact that the integument is a part of the whole organism, and therefore is subject to the great laws of general pathology, and that diseases affecting the integument are not only to be studied as localized areas of disease, but also in their relation to the system at large.

For the majority of cases no laws can be laid down tending to separate the local from the so-called general diseases. This is particularly true of the inflammatory affections, and I believe in many cases it is wise not to press such questions too closely in our endeavor to determine these intricate problems. Local pathology, interesting and satisfactory as it is as a study, must never be permitted to outweigh the general process, to which in many instances we must look as the *fons et origo* of the disease. That some diseases are strictly local in all their aspects will be denied by no one, but it is practically often difficult to decide where to draw the line between such affections and those due to influences and causes remote from the skin, as, for example, in the case of the many and often obscure reflex affections.

On the other hand, it is not difficult with our knowledge of to-day, to give examples of some true skin diseases. Notably among these may be mentioned the local parasitic affections, the inflammations due to numerous external causes, as for example the rhus plant and certain of the hypertrophic, atrophic and neoplastic diseases, such as callositas, clavus, cornu, molluscum fibrosum and molluscum epitheliale. In these affections, and in some others that might be cited, as far as our knowledge extends, the skin is the only organ of the body invaded at any time in the course of the disease. But while this holds true for a considerable number, there are still many that are in reality widespread diseases,

the skin being only one of the organs involved by the process.

As instances of such may be cited the so-called exanthemata, as well as other eruptive fevers, the eruptions due to various poisoned states of the system, as in septicemia, glanders, leprosy and syphilis. There also exists another group of well-known cutaneous manifestations in which it is difficult to determine whether the process is really confined to the skin, and whether it does not also involve other structures of the body as, for example, the epithelium generally, as in pityriasis rubra and dermatitis exfoliativa. Another group would comprise herpes zoster, pemphigus and the like, where the nervous structures, central or peripheral, are in some way at fault.

Still another class consists of diseases which, while eminently skin diseases, are in some cases, at least, dependent upon certain peculiar states of the economy for their existence, as a type of which psoriasis may be given. As an example of the effect of the state of the system at large upon this disease, I may cite the case of an old gentleman, an experienced physician, whose chronic and inveterate psoriasis had resisted all manner of internal and local treatment, but who upon going to sea immediately experienced relief and a rapid diminution in all the symptoms, so that in less than a fortnight he had practically recovered from a disease which had for a long period proved entirely intractable to our best remedies.

Another well-defined disease attacking the skin is eczema. While this is often a local affection, it may also be started into existence and kept up by serious internal derangements of the economy. In the latter cases it must be regarded in part, at least, as a symptomatic affection, as in the case of other symptomatic disorders, as urticaria; for with improvement of the digestive and assimilative functions or nervous system the eruption improves or disappears. These observations are particularly striking in the case of infants and children.

But it is not so much my purpose to specify and to differentiate the local from the general diseases affecting the skin, as it is to direct attention to what I believe to be an important truth in medicine, namely, that many cutaneous lesions which have long been and still are regarded as local skin diseases are really cutaneous expressions of certain general pathologic processes, due to varied causes. If this proposition, which might be elaborated, be accepted as an accurate observation, it brings us to the point I am desirous of establishing, namely, that our ideas of dermatology should be greatly enlarged. I am firmly of the opinion that this branch of medicine should include all morbid manifestations that appear on the skin, whatever may be their cause, their nature and their character. Dermatology has properly to do with the integument and all that pertains to it, and moreover with all the varied causes that may disturb that organ. Thus the exanthematous and the numerous and diverse symptomatic eruptions, whether superficial or deep-seated, fugitive or persistent, are all entitled to a place in the group. One observation in support of this view is that practically it is often impossible to differentiate the several varieties of cutaneous inflammation which resemble one another, except through the causes which produce them. It is a notable fact that the same pathologic state of the skin and the same

lesions may be produced by several and even diverse causes, directly or remotely, entirely distinct from one another, as in the case of acne, acne rosacea and urticaria.

Thus we note in practice that the lines separating the so-called idiopathic from the symptomatic diseases are far from being so sharp as is generally supposed. Authors of text-books and systems of dermatology would have us believe that the various diseases may all be arranged and satisfactorily classified, and that they may in all cases be readily differentiated from one another. While this undoubtedly holds true for the majority of cases, ample allowance must be made for atypical, irregular and anomalous forms of disease, of which there occur, I am convinced, more examples than most authors are disposed to admit. Nor is sufficient latitude given in describing disease in general for complications which may occur, nor for the merging of pathologic processes into one another. On this latter point, in particular, I would lay stress, especially as expressing certain anomalous forms of diseases. Mr. Hutchinson, of London, was one of the first to direct attention to this point, of which we all I am sure have seen examples.

As our knowledge of disease grows, so does the list of diseases increase, new affections, new varieties and combinations appearing and being described. To illustrate this observation we have but to look over treatises on dermatology published twenty-five years ago and to compare them with the works of to-day. The growth in this direction has been remarkable. Not only has the list increased, but it is still growing from year to year. Observations of this kind go to show, I think, that it is only a matter of time when every lesion that may occur on the integument must be included in dermatology. There was a period, not many years ago, as most of us can remember, when erysipelas was regarded as an exanthema—an exanthematic fever, and as being therefore altogether beyond the pale of dermatology. In those days that disease, it was said, was not a skin disease but a constitutional or a general disease, and that its relation to the skin was only incidental. How fallacious was this view, how slight our knowledge about this disease then, I need not more than refer to. I believe that before long certain other diseases will be recognized as being properly entitled to a place in cutaneous medicine which now are relegated elsewhere. Such a grouping should obtain, not with the view of magnifying the subject, but that the various similar and dissimilar diseases and the expressions which they are capable of producing on the skin, may be brought together for purposes of study and comparison.

The time has arrived when we should endeavor to recognize not only the particular form of eruption, but what is more important, also, the cause which has produced it, upon which success in treatment may depend. Diseases of the skin must be studied from the standpoint of general medicine. It is not possible to comprehend the meaning of certain forms of inflammation of the skin without taking the broadest view of the subject. Thus if we are inclined to regard such diseases as dermatitis exfoliativa, pityriasis rubra, lichen ruber, dermatitis herpetiformis, and the like as mere local cutaneous inflammations we fail to understand the significance of the symptoms. Symptoms and causes must be studied together. The former are elementary, and constitute

the alphabet of dermatology, which it need not be stated must be learned. But eruptions in themselves, as mere forms of superficial inflammation, are by no means so important as the relation of the lesions to the causes. To express this idea more plainly, take the disease, eczema, as an instance. Here in some cases, as in neurotic eczemas, the discovery of the causes at work in producing the eruption is the key to the situation, without understanding which no success in the treatment will probably be obtained, and many other similar instances might be cited. Thus it happens that some diseases are practically uninfluenced by local treatment, and that not until we investigate their possible relations with the general economy do we appreciate their nature. The general practitioner, who has been trained in cutaneous diseases, has the opportunity of advancing dermatology quite as much as the specialist, and his services in aiding the good work can not be dispensed with. Some of the best work latterly has been done by men who are not only accomplished dermatologists but equally distinguished general practitioners of medicine and surgery. We owe a great deal to such writers and teachers as Jonathan Hutchinson, Paget, Stephen Mackenzie, Fagge, Pye-Smith, Robert Liveing, Duckworth and McCall Anderson.

The objective study of skin diseases is fascinating, and has without question its uses in more ways than one, but it is important that we take steps to advance beyond this elementary stage of knowledge, by attempting to recognize and understand the meaning of the local manifestation. It has been, and still is, too much the custom to study diseases of the skin in the light of pathologic pictures, to name the local manifestation and to so label it as a disease. It is much easier to give the disease a name and to label it than it is to comprehend the process at work. The former is comparatively unimportant for the patient, the latter a point upon which recovery may depend. The nature and meaning of the process in connection with the cutaneous symptoms has not received enough attention, and I believe this to be one reason why the treatment of many of these diseases in the past has been so notoriously unsatisfactory. At all events the relations of the cutaneous disturbance to other structures and to various states of the economy should be much more thoroughly investigated by dermatologists than is the present custom.

To recognize any one disease, say syphilis, in its varied manifestations on the skin, requires familiarity with all other diseases with which it is liable to be confounded. Therefore, to be an accomplished diagnostician, one must be conversant with every form of eruption to which the skin is liable, including the not rare atypical and aberrant forms. In regarding only the well-known and clearly defined obvious diseases of the skin as belonging to dermatology, this branch of medicine is not only belittled, but the true meaning of many lesions on the skin is not appreciated. I would insist, therefore, that the manifold and varied changes that take place in the skin due, as we now know, to such a multitude of diverse causes, should be viewed as phases of cutaneous medicine rather than as skin diseases. The idea of this vast array of diseases being morbid entities, for which the integument alone is accountable, must in many cases, at least, be abandoned in favor of the principles of general medicine.

Allusion has been made to the fact that the same

or similar lesions may be called forth by wholly different causes. This also is a point to which attention has not been sufficiently directed by writers. As illustrative of the observation, the similarity in the eruptions of eczema and scabies may be referred to. Here the causes at work are different but the result, as far as the skin is concerned, is much the same. On the other hand, the same cause may produce entirely different cutaneous lesions. A striking example illustrating this idea is found in dermatitis herpetiformis. Here the same cause at one period of the disease gives rise to an eruption resembling erythema multiforme, at another period to herpes, again to impetigo, and finally to pemphigus. We note thus that one process may occasion numerous distinct lesions, which individually may resemble what have been heretofore regarded as well-known distinct diseases.

If we would study dermatology with the view of learning all that it can teach us, not only of the skin but of general medicine, we must look in the majority of cases beyond the mere eruption, valuable and important as this is in all cases as a guide to the pathologic process at work. While, even in the case of local affections, it may not be necessary to look beyond the skin itself, we may nevertheless find interesting facts bearing on cause and pathology by searching within the economy. The cutaneous disturbance in many instances is distinctly indicative of some irritation or disorder in other regions, organs, structures or fluids of the body. That we are not able always to discover and to designate the causes, merely shows our ignorance. I am of the opinion that the relations of the skin to other parts and functions of the economy are at the present date only partially understood, and that there are many points which will sooner or later be elucidated which will bring cutaneous medicine still closer to general medicine. Much, however, has been accomplished in this direction during the past twenty-five years, and I am pleased to add that a fair proportion of this good work has come from the labor of our countrymen.

This brings me again to the question propounded at the beginning of these remarks, viz.: What are so-called skin diseases? The answer, I believe, has been given through the principles to which attention has been called. It is simply this, that our conception of the scope of dermatology must be so widened as to include every pathologic manifestation which occurs in the integument, irrespective of the cause or the nature, from a practical standpoint. The great value and importance of dermatology is that it should teach us to know the nature of various processes, as they affect not only the skin but the whole economy. Dermatology should be for the physician as a key with which the skin is made to reveal, in many instances at least, the nature of the process at work in the general system or in special organs, which without this aid might remain obscure. Striking examples supporting this view are noted in syphilis and in leprosy, where the cutaneous manifestations are sometimes the only indication of the presence of these diseases in the body. The recognition of the nature of the cutaneous lesions is often of the greatest value in the general diagnosis. This observation applies pointedly in the case of the erythemata and certain general infectious diseases.

While, therefore, not losing sight of the fact that

some cutaneous diseases are strictly confined to the structure involved and have no other than a local significance, there are many in which the skin lesions must be regarded as being merely one set of a series of symptoms due to some special or general cause having its seat in other structures as well as in the integument. The part that the nervous system plays in the production of varied diseases of the skin is, I am firmly convinced, immense. While its power and influence as a factor is well understood to-day, owing largely to the special writings of such observers as Eulenberg and Guttman, Bulkley, E. Long, Fox, Crocker, Schwimmer, Kopp and many others, I am of the opinion that its influence over the skin is at present far from being properly recognized. The nerves are avenues by which a multitude of diverse influences reach the skin. The future will, I feel sure, see this particular subject much more elaborated than it is to-day.

ERYTHEMA EXFOLIATIVUM RECURRENS.

Read in the Section on Dermatology and Syphilography, at the Forty-fourth Annual Meeting of the American Medical Association.

BY A. H. OHMANN-DUMESNIL, M.D.

ST. LOUIS, MO.

The erythemata have always possessed a large amount of interest for dermatologists, and the numerous apparently aberrant forms which have been observed in late years have invested the subject with an amount of importance which was not conceded to it heretofore. It is more particularly in regard to the etiology of the different forms that research has been made. In addition to this there have been presented certain forms which have excited much interest on account of their assumed rarity, but which are, possibly, not so infrequent either on account of mistakes in diagnosis, or ignorance of this pseudorarity, or an indifference to making any record. These circumstances, singly or combined, have conspired to produce an impression in many instances that a certain condition or group of symptoms was rare, when in reality it was common and had either escaped attention or failed to arouse any interest, under the mistaken idea that it was so well known as not to deserve even a passing mention. Under these circumstances, every new record helps to swell the list and without detracting from the credit of those who first noted the matter, it contributes to a vulgarization of a certain amount of knowledge which should come within the grasp of every one. It is for this reason that I wish to place upon record a condition which is an apparently unusual one, but which in my opinion is observed much more frequently than medical literature would lead us to suppose. I refer to intermittent scarlatiniform erythema or erythema exfoliativum recurrens. Medical literature, up to the present, seems to be rather meager, so far as the number of accounts of this trouble is concerned. A hasty search through current medical literature and works on dermatology has yielded but small results and larger returns will have to be left to the future.

Before entering into any farther considerations upon the subject, I will give the history of a case as furnished by Dr. Edward C. Bennett, under whose care it occurred, and to whom I am indebted, not only for the anamnesis but for the specimens which are figured, as well. I am desirous of returning my thanks to him for this, as he kindly conferred with me

regarding the case which he very justly considered an unusual one.

In order that a more complete record of the disease under consideration may be made, I will append condensed histories of other cases which have recently appeared in print, in order that a more easy reference may be made and the points of similarity, both from a clinical and etiological point of view may be made more apparent, and thus enable the reader to follow the reflections deduced, with greater facility.

The cases which I propose giving do not by any means include all those which have been reported, but they are sufficiently numerous to give a general idea of the characteristics of the disease, as well as to give a certain amount of information in regard to the clinical peculiarities of the process, and thus afford an opportunity of drawing some conclusions in regard to the etiology of the process. In addition to this, it affords us almost a certainty in deducing conclusions as to its comparative frequency. If we are to judge from the number of cases which have recently appeared in current medical literature, we would be led to the conclusion that the trouble while not frequent is far from being unique or even rare.

With these few prefatory remarks, I will proceed to give a brief résumé of a few cases which have recently been described and then make a brief analysis of them:

Case 1.—(J. Frank, M.D. and W. C. Sandford, M.D., in the *American Journal of Medical Sciences*, August, 1891.) John H. P., miner, 34 years of age, well-built and healthy. Skin is perfectly normal. His parents are living as also maternal grandmother; is the second of a family of thirteen, all of whom are living. On July 24, following his birth (Dec. 29, 1857) he was suddenly taken ill, vomited and in a few hours the entire surface of the body was scarlet red. The symptoms subsided in a few hours, but on the fourth or fifth day following the attack the entire cuticle was cast off, and a few days later the nails of his hands and feet were also shed. This was repeated every year on the same date. The patient first remembers the shedding in 1865 and he states that these attacks occur each year on July 24, usually at 3 p.m. and never later than 9 p.m. The paroxysm begins abruptly. Patient has a feeling of lassitude and weakness of fifteen to twenty minutes duration, followed by muscular tremors, nausea and vomiting, a rapid rise of temperature, skin and mucous membrane of tongue and mouth become red and inflamed, and are hot and dry. No perspiration appears after the paroxysm begins until the cuticle is cast off. The patient has been delirious three times during these attacks, once for nine days. In his early life the cuticle began to shed on the second or third day after symptoms appeared, and was complete by the fifth day; but each succeeding year it takes a little longer, until now it is ten or twelve days before shedding is complete. The cuticle can be detached in large sheets, and from the hands and feet in the form of gloves and moccasins. The nails are loosened and crowded off in about four weeks after the acute stage.

Here follows a detailed account of the attack:

Vomiting took place, the erythema extending visibly. Pulse 68; temperature 97 degrees. The highest pulse noted and temperature observed during the attack were 92 and 103 degrees respectively. July 26, two days after the beginning of the attack, the skin appeared normal, the temperature having returned to the normal. July 27 the epithelium of tongue and mouth came away. July 28, perspiration was free on forehead and under eyes. The cuticle on chest was raised in the form of blisters by the perspiration. Desquamation then set in and continued until August 11, when the left moccasin was removed. After the removal of the cuticle the skin was very soft and delicate; and where the former was normally thick, the new skin was very sensitive. August 26, the nails of the little finger and second finger of right hand were shed. September 2 those of the little finger, second and third fingers of the left hand were retracted. Both thumb nails removed September 5; and the nails from