BRACHIAL NEURITIS AND SCIATICA

HUGH T. PATRICK, M.D.

Clinical Professor of Nervous and Mental Diseases, Northwestern
University Medical School
CHICAGO

A misapprehension as to the frequency of brachial neuritis and of sciatic neuritis or sciatica I believe

Fig. 1 (Case 1).—Arthritic side, showing free flexion of thigh.

to be quite general. At any rate, I do know that in the profession at large, as I meet it, the diagnosis of these diseases is made altogether too frequently.

BRACHIAL NEURITIS AND SHOULDER
ARTHRITIS

A careful review of my records (private practice) for the last ten years shows that, of the patients referred to me supposedly with brachial neuritis, only about one in ten really had it. Approximately four fifths were cases of arthritis of the shoulder joint. The other one tenth was made up of cases of bursitis, syphilis, neoplasm, cervical rib, postherpetic pain, osteomyelitis, cervical caries and cervical arthritis. Indeed, when a patient states that he has for some time been troubled with neuritis of the shoulder or arm, one is pretty safe in assuming that it is something else, probably arthritis.

Why should so many physicians have made an erroneous diagnosis of brachial neuritis? Principally, I think, on account of the relative prominence of pain,

Fig. 2 (Case 1).—Arthritic side; no Lasègue sign.

because this pain was not distinctly localized in the shoulder joint and probably because they did not sufficiently examine the patient. A careful examination seldom leaves one in doubt.

Neuritis is rare; the pain is constant, is influenced not at all or but little by passive movement of the

shoulder, and is generally influenced not greatly by active movement, though the patient moves the arm carefully. Ordinarily, the suffering is intense, such as never occurs in arthritis except in the worst cases, when the patient cannot move the joint at all. Intermittent, grumbling pain is very, very rarely that of neuritis. There is distinct, generally exquisite, tenderness of the cords of the brachial plexus or of the nerve trunks or of both. In some cases there is distinct though slight tactile anesthesia and sometimes surface hyperesthesia—more properly speaking hyperalgesia. Tingling or subjective numbness of fingers or forearm is frequent. The deep reflexes of the arm are diminished or lost. The disease is nearly always self limited. It lasts a few weeks or a few months (in bad cases from six to nine months), there is definite recovery, and it seldom recurs.

Arthritis of the shoulder, on the other hand, is very common, and the pain fluctuates more or less, even when very bad. In the less severe cases it is intermittent. Not only is it intermittent within the twenty-four hours, but the patient has his good and his bad days. Often the pain seems to be influenced by the weather, and for some reason it is apt to be more annoying at night. As before stated, often the pain is

Fig. 3 (Case 1).—Normal side; leg almost horizontal; that is, no Fabere sign.

not distinctly localized in the shoulder joint. In a great many cases it is felt chiefly at about the insertion of the deltoid. For this I have no adequate explanation, but I know that it is very misleading. More than once I have known it to lead to a diagnosis of osteomyelitis. Sometimes the pain is rather diffuse, radiating up into the neck and down the arm. This pain in the neck and down the arm I am quite sure is often due to concomitant arthritis of the cervical spine, the nerve roots being irritated where they pass out through the intervertebral foramina. In some cases of shoulder arthritis alone I have no doubt that the nerve trunks may be involved where they pass close to the joint.

The striking feature, however, of the arthritis cases is that in whatever part the pain may be felt, there are certain movements of the shoulder joint, active or passive, which cause pain. These are by no means always the same; but by all odds the most frequent is adduction of the humerus with retraction and internal rotation, the movement involved in crossing the forearms behind the back. Consequently the woman has difficulty in fastening her skirts and her waist behind, and the man in putting his hand into his hip pocket. Another good test movement is to extend the arm straight upward and then rotate it as if operating a screw driver. In the examination of such a case,

movements in different directions, with the arms in different positions, should be carried out against resistance. For instance, with the arms extended horizontally and parallel in front of the upright trunk, the hands are pushed toward each other, away from each other, upward and downward *against resistance*. With the arms extended sidewise, making a cross with the body, resistance movements are made upward, downward, forward and back, etc. One or more of these movements will be pretty sure to be painful, which would not be the case in mild neuritis. But it must be emphasized that in many cases of subacute and chronic arthritis, many movements or most movements are entirely painless. I have had a number of patients who could play golf with perfect freedom; and such exercises as hammering, sawing, rowing and sweeping may have no immediate influence on the pain. Likewise, in the chronic or subacute cases, local tenderness may be but slight or even absent. Of course in acute arthritis practically all movements of the joint are painful, and there is limitation of passive motion, neither of which conditions obtains in neuritis.

Of very considerable assistance in the diagnosis is the state of the deep reflexes. These are uniformly

Fig. 4 (Case 1).—Arthritic side; knee cannot be normally depressed: fabere sign.

brisk, and sometimes considerably exaggerated. Even when the muscles are quite atrophic and the appearance of the arm would indicate loss of the deep reflexes, the wrist jerk and triceps jerk will be found to be quite lively, more so than those of the well side. The same, as a rule, applies to the scapular jerks.

Wasting of shoulder and arm muscles sometimes leads the observer astray. There seems to be a common feeling that atrophy means nerve trouble. Consequently wasted deltoid, biceps and triceps apparently lend color to the view that the shoulder pain is neuritic. Just how arthritis causes atrophy of muscles moving the affected joint is still matter for speculation; but the fact of its general presence is of daily observation. The atrophy may be very marked and the arthritis relatively mild. I once saw a boy with mild multiple arthritis in whom the muscular wasting was so obtrusive that the case was thought to be one of advanced progressive muscular atrophy. Indeed, I have known men high in the profession to be misled by this arthritic atrophy.

There are two more signs of value: signs, I regret to say, that seem to be infrequently looked for. These are fever and leukocytosis. In very mild and chronic cases they may not be found, but such are exceptional.

Repeated examinations, especially when the pain is at its worst, will nearly always show at one time or another (sometimes uniformly) a slight rise of temperature (from 99 to 100 F.) and a slight increase in

Fig. 5 (Case 1).—Same as Figure 3.

the leukocyte count (from 8,000 to 9,000 or 10,000, occasionally 12,000). These two signs naturally lead us toward the conclusion that the arthritis is of infectious origin, and this conclusion is supported by the fact that less obtrusive arthritis elsewhere can often be found. Even when the patient avers that he has no other trouble, painstaking inquiry and examination frequently will show that his statement is inaccurate. Other joints oftenest involved are the opposite shoulder and the articulations of the cervical spine. Perhaps next in order come the lumbar and the dorsal spine, the hip and the finger joints. Furthermore, it is not so very unusual to find a coexistent indurative headache or soreness and tenderness of the occipital or suboccipital region, troubles I believe to be almost invariably due to infection, the focus generally being somewhere in the head or the throat. Of like significance is tenderness of the trapezius or other neck muscles—a fairly frequent concomitant.

Fig. 6 (Case 1).—Same as Figure 4.

SCIATICA AND ARTHRITIS OF THE HIP

Thirty-odd years ago Dr. L. Putzel of New York called to my attention the frequent mistaking of chronic arthritis of the hip for sciatica. Since that time I have been interested in observing how much

more common is the arthritis than sciatica, and how frequently an erroneous diagnosis is made.¹ Outside of large charity hospitals I believe sciatica to be a relatively rare disease, while arthritis of the hip is one of the most common. This is no new assertion. Many years ago that wonderful observer Jonathan Hutchinson expressed the opinion that a certain number of cases of sciatic pain were due to "trouble in the joint." Since then similar statements have now and then been

Fig. 7 (Case 2).—Arthritic side; good flexion.

made, and recently the elder Bruce² has published a remarkable little book sustaining the thesis that the pathology of sciatica is "trouble in the hip joint." Inferentially he concludes that all sciatica is hip joint trouble. I cannot go so far as he because I am quite sure that occasionally one does see a true neuritis of the sciatic nerve or sacral plexus, and semioccasionally an example of neuralgic sciatica—or at least of sciatica in which the signs of neuritis, except pain, are absent.

Without touching on the reasons why arthritis leads to a diagnosis of sciatica, I shall simply mention a few signs and symptoms indicating the former and excluding the latter.

Sciatica, like brachial neuritis, causes no limitation of passive motion, with one notable exception—the Lasègue³ sign—whereas, arthritis of the hip is pretty sure to interfere with some movement or other. Especially is there one sign, so far as I know never present in sciatica, which is of great value in detecting arthritis of lower grades, but which I have not seen described.

Fig. 8 (Case 2).—Arthritic side; no Lasègue sign.

For convenience of record I call it the *fabere* sign, the letters of the word standing for flexion, abduction,

external rotation and extension. With the patient supine on a level surface, the thigh is flexed and then the external malleolus or ankle is placed above the patella of the opposite extended leg. In bad cases even this maneuver cannot be executed without pain, but many patients do it easily. If, however, the knee is now depressed, the ankle maintaining its position above the opposite knee, the patient will complain of pain before the knee reaches the level easily attained in normal persons.⁴ This procedure involves flexion, abduction, external rotation and then extension of the thigh, and will catch more cases of arthritis of the hip than any other test with which I am acquainted. Many of the patients have quite free motion of the joint, especially flexion, and in many of them it is not easy to detect any tenderness, though firm pressure with the point of the thumb over some part of the capsule may be much more painful than on the good side. Bruce thinks that the most frequent tender point is at the upper edge of the acetabulum. Heavy percussion of the sole, or the quick shoving of the head of the femur into the acetabulum, which is almost certain to cause pain in tuberculous hip joint disease,

Fig. 9 (Case 2).—Arthritic side; fabere sign; knee cannot be normally depressed.

generally fails to reveal anything in these nontuberculous cases. But rare indeed is the case which will not be detected by the test described above. In making the test, one must not be deceived by the patient's yielding to depression of the knee by tipping the pelvis. This he instinctively does to avoid pain. The pelvis must remain level.

Figures 1 to 6 illustrate a case in which a woman, aged 50, complained of pain about the left hip and from the hip to the knee (Case 1). She was sent to me by one of the best surgeons in the country with the statement, among others, that a careful examination, including roentgen-ray plates, showed nothing in the hip joint. But the case was one of arthritis of the hip and, incidentally, of the left knee. Figures 7, 8 and 9 are from another case of hip arthritis (Case 2). This patient had not only free and painless flexion of the thigh, but also good abduction and rotation; and yet in this test the knee would go no lower than the point indicated in Figure 9.

It goes without saying that other tests should be used as indicated in the shoulder cases: extension,

1. A number of times I have known the sciatic nerve to be injected or stretched when the trouble was in the joint.

2. Bruce, William: *Sciatica*, New York, 1913.

3. Very similar to Kernig's sign. With the knee well flexed, the thigh can be fully flexed without pain; but when, with the thigh thus flexed, an attempt is made to extend the leg, it cannot be got much beyond a right angle without causing pain.

4. If the soreness has all gone from the joint and only the relics of a previous arthritis are present, there will be no pain: only inability to bring the knee down to the normal point.

flexion, abduction, adduction, rotation, strong abduction with rotation, etc., *against resistance*.

As shoulder arthritis is often accompanied by a like condition of the cervical spine, so hip arthritis frequently is complicated by involvement of the lumbar spine, and probably of the fibrous structures overlying the sacrum.⁵ This may cause pain radiating down the extremity, thus assisting in the simulation of sciatica. Bruce believes that the pain down the leg is referred pain, like the pain in the knee of tuberculous hip disease.

If arthritis of the hip is present, naturally we would look for more or less muscular wasting. And this wasting, instead of being an embarrassment in the diagnosis, is an aid. Atrophy due to inflammation of the sciatic nerve of course involves all the muscles supplied by this nerve, no others, and probably the peripheral ones (leg muscles) more. The atrophy of hip arthritis, on the other hand, is found in the muscles that move the joint, especially, as Bruce has insisted, the gluteals. With the patient prone, a distinct flattening of the buttock often is apparent, and with the patient standing, a difference in the gluteal folds.

The other differential criteria distinguishing hip arthritis from sciatica may be taken over bodily from the shoulder cases. In neuritis of the sciatic nerve the Achilles jerk is diminished or lost, whereas in arthritis of the hip it is normal or, more frequently, increased. The knee jerk on the affected side is also nearly always increased. But in arthritis of the spine with considerable involvement of the nerves, these deep reflexes may be diminished.

Bruce⁶ says of sciatica, "I have rarely found the tender points of the books." I entirely agree with him. The first reason is that most of the cases are not sciatica. But even in this disease the classical points of Valleix are a delusion and a snare. In arthritis, more or less tenderness on the affected side is almost the rule.

How much of this is of organic origin and how much due to the hypersensitiveness of a long-suffering patient is not always easy to say. I know that in arthritis of the hip, if one firmly punches the gluteal point, the femoral point, the popliteal point and the submalleolar point, frequently one or more of them will be or seem to be tender. But if the examiner is not satisfied with this, and proceeds to punch other places, from the iliac crest to the heel, he will find muscular tenderness, joint tenderness, tendon or fascia tenderness, sometimes even periosteal tenderness, all of which has nothing to do with the sciatic nerve.

Concerning one striking sign or consequence of sciatica, I am in doubt. This is the high degree of scoliosis or lateral curvature sometimes observed. I do not know whether this can be caused by simple arthritis of the hip, but I think that it can.

25 East Washington Street.

TORULA IN MAN

REPORT OF A CASE WITH NECROPSY FINDINGS

PHILIP H. PIERSON, M.D.

SAN FRANCISCO

Torula is a form of yeast, producing lesions preferably in the central nervous system. These lesions macroscopically simulate tuberculosis, from which disease they must be differentiated as well as from lesions formed by coccidioides and blastomycetes. It is not my object in this paper to go deeply into the differential features of these three diseases but briefly to point

out the main differences and then present a case of torula infection in a man admitted to the San Francisco Hospital, September, 1916.

It is definitely agreed that coccidioidal granuloma and blastomycosis are two different diseases. Coccidioidal granuloma is a disease produced by an organism which sporulates but never buds in tissue. It varies greatly in size, having large forms up to 30 or 40 or even 80 microns in diameter producing many small ascospores which, in turn, escape from the capsule. It simulates tuberculosis in the type of its lesions and in the manner of spreading, principally by the lymphatics

and thence to the blood stream. It is largely confined to men in the San Joaquin Valley, California. It is nearly always fatal, as no treatment has been found that is of any benefit.

The organism of blastomycosis varies in size from 3 to 20 microns. It buds but never sporulates, the buds being from one-half to one-third the size of the parent cell. It grows to the size of the parent cell before reproducing itself. It produces nodules with or without caseation, and deep and superficial abscesses with polymorphonuclear leukocytosis. It may affect all organs, including, as a rule, the skin and the bones, and it is helped by iodids. On culture mediums both the blastomycetes and coccidioides grow mycelium sooner or later. The organism of coccidioidal granuloma also produces chlamydospores at the ends of the aerial hyphae in cultures.

Years ago torulae were found to have a marked pathogenic action in animals; but prior to this case there have been only six reported in man—two by Rusk, two by Stoddard and Cutler, and one each by

Fig. 1.—Low power view of lesion from cortex, several organisms seen with clear zone of gelatinous material surrounding them.

5. Cases thought to be lumbago with sciatica or alternation of these two affections are generally cases of arthritis of spine and hip or of the former alone.

6. Bruce, William: Sciatica, p. 11.