Lahey, F. H. LOSS OF BOTH EYES IN GRAVES'S DISEASE. [Boston Med. and Surg. Jl., April 22, 1920.]

A woman with such severe exophthalmos as to necessitate enucleation of the right eye in consequence of edema and ulceration of the cornea is here reported upon. A month later the superior cervical sympathetic ganglion was removed on the left side; at the same time the external canthus was incised. The exophthalmos, ulceration and edema continued to increase, and two days later the lids were sutured together. Two months later this eye had to be removed.

Rachford, B. K. SUBSTERNAL GOITER WITH PRESSURE SYMPTOMS. [Ann. Jl. Med. Sci., September, 1920.]

Rachford's patient was treated successfully, so far as relief of pressure symptoms is concerned, by rest, röntgen ray, fifteen drops of tincture of digitalis, three times a day, and a capsule containing $2\frac{1}{2}$ grams of the neutral bromid of quinin, 1 grain extract of ergot and $\frac{1}{80}$ grain extract of belladonna, three times a day, for a long period of time. At rare intervals this treatment has been interrupted for a period of a few weeks.

Ishihara, M. EXPERIMENTAL EXTIRPATION OF THYROID. [Bull. Naval Med. Ass. of Japan, June, 1920.]

Enlargement of the thymus gland followed excision of the thyroid in rabbits, due to a compensatory action existing between these organs. Degenerated corpuscles of Hassall were seen more frequently in the thymus gland of normal rabbits than in operated ones. Enlargement of the pituitary body was not observed.

Squier, T. L. IMPROVEMENT IN EXOPHTHALMIC GOITER SUBSEQUENT TO SEVERE FOCAL INFECTION. [Am. Jl. Med. Sci., September, 1920.]

Two instances of hyperthyroidism complicated by infections caused an increase in the thyroid symptoms. After the acute infection had subsided the situation was in part or entirely relieved. The improvement is attributed to actual loss of secretory tissue following the infection.

Trott, R. M. BLOOD PICTURE BEFORE AND AFTER GOETSCH EPINEPHRIN TEST. [Arch. Int. Med., September 15, 1920.]

If 4-8 minims of adrenalin be given subcutaneously to individuals with apparent or even externally unrecognizable hyperthyroidism, there will result, according to this observer, a rise in leucocyte count with definite lymphocytosis, which latter may be absent in ataxic goitrous state.