POLIOMYELITIS.

S. Flexner, of New York (*Journal A. M. A.*, September 24), gives a resume of the experimental contributions to our knowledge of human poliomyelitis. These he shows have yielded a number of important facts relating to the spontaneous disease in man. These facts include the discovery of the nature of the virus, of many of its properties, of certain important clinical and pathological peculiarities of this disease, of the phenomena of immunity, of a mode of 'spontaneous infection through the nasopharyngeal mucosa, and have also served as a basis on which to develop measures of prevention and on which it is hoped to work out a successful treatment. The advances which have been made will make it possible to accurately determine whether the disease is a single clinical and pathologic entity or whether it is a generic name covering groups of symptoms and due to the effects of certain lesions of the brain and cord, of which it is not merely one, but the most important, variety. It is improbable, he says, that poliomyelitis, like meningitis, may be found to rise from several independent causes and that epidemic poliomyelitis, like epidemic meningitis, may be distinguished among the varieties by being always the result of a particular micro-organism. We have already evidence that certain animals like the dog, poultry, and possibly the horse, are subject to poliomyelitis due to a special germ in each case.

TUBERCULOSIS IN MILK.

F. O. Tonney, Chicago (*Journal A. M. A.*, October 8), gives an account of the work done by the Chicago Department of Health in the examination of market milk for tuberculosis germs. The samples were collected by the health department inspectors, partly in the original containers and partly in sterilized bottles furnished by the department, and the effort was made to have the samples reach the laboratories within a few hours from collection. Guineapigs were used for the tests. After inoculation they were kept in clean

cages and were observed for three months. It is to be mentioned, however, that inasmuch as the negative animals were not post-mortemed, but allowed to live and be used for other purposes, the results of the series must be reported as minimum findings only. No cultural studies were attempted to determine the presence of human and bovine types of the organism. From a strictly scientific point of view it is to be regretted that cultures were not made from the organs to exclude the possible presence of Rabinovitch's butter bacillus. The autopsies were made when the laboratories were overworked, but they were in every case well marked and typical, and it is believed that the positive findings reported may be considered tuberculosis. The total number of milk samples examined was 163; of these, 51 caused the death of the animals injected from acute infection within three weeks before diagnosis of tuberculosis could be made; 112 remained available, of which 10, or 8.9 per cent, proved tuberculous. Comparing this with observations made in other cities and combining their figures with these, he finds the incidence of tubercle bacilli in the milk supplies to be about 8.3 per cent. In Chicago, the situation is being met by a regulation providing that all milk sold beginning January 1, 1914, shall be obtained from tuberculin-tested cows. In the interim milk not thus obtained may be sold, provided it is pasteurized according to the rules and regulations of the department of health. Under this ordinance about 54 per cent of the milk sold in Chicago at the time of writing is pasteurized; 24 per cent is tuberculin tested. It is calculated that before the close of the season the remaining 22 per cent will be pasteurized. During the summer of 1909, when only 30 per cent was pasteurized, a decrease of 521 was noted in the death rate of children under 1 year, chiefly among children suffering from diarrheal diseases. The milk-produced diseases of children are, he thinks, first in order, the diarrheal diseases responsible for about one-third of the deaths among children under 2 years in our large cities; second, is typhoid