produce similar symptoms. At a later date, local manifestations of the disease still proving rebellious, a third dose of 5,000 units was injected. Ten days later the eruption had become wide-spread, appearing all over the body. A different treatment was instituted, but the woman's people became dissatisfied, and the doctor in charge was dismissed.

Because of the fact that the patient's mother does not "believe in doctors," three weeks elapsed before another physician was called. At this time symptoms of an unusually malignant case of syphilis were in evidence. Complete anorexia, profound melancholia and weakness marked a serious condition. The characteristic eruption consisted of lesions as wide-spread and numerous as those which a severe case of discrete small-pox might exhibit. Iritis, ulcerative blepharitis and a mouth and throat as sore as that of the worst pellagra sufferer completed the patient's misery. (These symptoms are recited to emphasize what every case does not show so plainly, that the patient is injured more by a mistaken diagnosis than the doctor himself.) The Wassermann test was strongly positive. The woman's husband, confronted with the facts, admitted having become infected during his wife's pregnancy. He had been treated with salvarsan in a neighboring city.

CONCLUSIONS

Thorough physical examination when the physician was first called would have disclosed the fact that general glandular enlargement was present. This, combined with the throat described and with the elimination of the possibility of diphtheria by cultures, would have furnished food for thought.

The giving of antitoxin in the first place, it will be admitted, was justifiable. "When in doubt, give antitoxin" is a good motto.

The mistake in diagnosis, discovered later, should have been confessed to some member of the family, and the real nature of the malady divulged. This disease cannot be well treated without some measure of cooperation; in addition, the jeopardy in which one's reputation has been placed in instances of this sort can be minimized by thus making the best of a bad situation.

Death During Ether Anesthesia—Status Lymphaticus.—

M. B., female, aged 3½, was admitted to the Montreal General Hospital on Nov. 12, 1913, for bilateral talipes valgus. She had been a premature infant (eighth month) and had been artificially fed. At the age of 18 months she had acute anterior poliomyelitis. In September, 1913, she underwent the operation of tonsillectomy under general anesthesia. At the time of her admission she appeared a well-nourished child and presented no signs of rickets. Examination of the heart and lungs was negative. The anterior and posterior cervical glands were palpable. There was no apparent enlargement of the spleen. The urine was normal. Nov. 15, 1913, she was given ether by the open method for a tendon-grafting operation. The course of the anesthesia was uneventful until the operation was nearly completed, that is, for about an hour. Breathing then became somewhat gasping in character, and within a minute ceased. Artificial respiration was immediately begun. Oxygen, camphor and strychnin were administered, but without success. Respiration was never reestablished. At the necropsy the following conditions were found: The thyroid gland was normal. The thymus gland was greatly enlarged especially on the left side. There was slight enlargement of the heart. The bronchial lymph-glands were hypertrophied and showed areas of caseation. A caseous gland was present in the posterior mediastinum in close contact with the branch of the left pulmonary vein from the lower lobe. The spleen was enlarged. Peyer's patches and the solitary follicles in the large intestine were enlarged and very prominent. There was hypertrophy of the mesenteric glands. The pancreas, liver, kidneys, adrenals and brain were normal.—W. B. HOWELL, M.D., Anesthetist to the Montreal General Hospital, Montreal, Canada.

ANKYLOSIS OF THE MANDIBLE

R. C. CREASY, M.D., WILKES-BARRE, PA.

W. C., aged 13 years and 7 months, undersized and underweight, came to me for the first time Aug. 9, 1913, with a limited motion of the mandible permitting of but ¼ inch separation of the incisors and no lateral motion. At the age of 4 years and 2 months, while swinging, he fell 8 feet and struck his chin on the hard wood floor. As the child suffered intense pain and was unable to move the mandible, a physician was summoned. Palliative treatment was instituted, and the child fed through a tube for a period of two months. No attempt was made to use the mandible, and it ankylosed in the position I found it. Roentgenoscopy disclosed the condition to be a previous fracture of the ramus, the line of fracture running into the joint, and a mass of callous substance filling the joint.

Ether was administered and the limit of motion was still the same, thus eliminating the spasmodic condition of the muscles as being one of the causal factors for the limited motion. Under force with the use of two oral mouth-gags, the degree of force applied being the same, thus avoiding any lateral dislocations, I succeeded in an increase of the limited motion to 1½ inches; this form of treatment was carried out three successive times at intervals of one week. During the time intervening I permitted the boy to make good use of the oral screw. The results obtained by this procedure have been excellent.

New and Nonofficial Remedies

THE FOLLOWING ADDITIONAL ARTICLES HAVE BEEN ACCEPTED BY THE COUNCIL ON PHARMACY AND CHEMISTRY OF THE AMERICAN MEDICAL ASSOCIATION. THEIR ACCEPTANCE HAS BEEN BASED LARGELY ON EVIDENCE SUPPLIED BY THE MANUFACTURER OR HIS AGENT AND IN PART ON INVESTIGATION MADE BY OR UNDER THE DIRECTION OF THE COUNCIL. CRITICISMS AND CORRECTIONS ARE ASKED FOR TO AID IN THE REVISION OF THE MATTER BEFORE PUBLICATION IN THE BOOK "NEW AND NONOFFICIAL REMEDIES."

THE COUNCIL DESIRES PHYSICIANS TO UNDERSTAND THAT THE ACCEPTANCE OF AN ARTICLE DOES NOT NECESSARILY MEAN A RECOMMENDATION, BUT THAT, SO FAR AS KNOWN, IT COMPLIES WITH THE RULES ADOPTED BY THE COUNCIL.

W. A. PUCKNER, SECRETARY.

TYPHOID VACCINE (See N. N. R., 1914, p. 259).

H. M. Alexander and Co., Marietta, Pa.

Typhoid Vaccine (Immunizing).—Prepared according to the method used in the U. S. Army Laboratory. The preparation is marketed in two styles of package: syringes and ampules, containing, respectively, 500 million, 1,000 million and 1,000 million killed typhoid bacilli. The injections are to be made at intervals of ten days.

B. B. CULTURE.—B. B. Culture is a pure culture of *Bacillus bulgaricus* said to be made from a desirable strain; marketed in bottles containing about 90 Cc. (3 fluidounces).

Actions and Uses.—B. B. Culture is intended for use in the treatment of intestinal indigestion and the various toxic and nervous effects arising therefrom; also for the enterocolitis of infants.

Dosage.—From 4 to 8 Cc. (from 1 to 2 fluidrams) three times a day in sweetened water before meals. For babies of any age 1 teaspoonful every three hours. The date of issue is stated on each bottle.

Manufactured by the B. B. Culture Laboratories, Yonkers, N. Y. No U. S. patent. U. S. trademark No. 90,535.

Brevity.—As a matter of fact we find that every really great writer tries to express his thoughts as purely, clearly, definitely and shortly as possible. Simplicity has always been held to be a mark of truth: it is also a mark of genius.—Schopenhauer.