Eruption from Confetti.—Foveau de Courmelles recently reported to the Paris Acad. de Méd. a case of a cutaneous eruption produced by confetti. The probable mode of poisoning was by the penetration of the anilin dyes from the confetti alighting on cracks in the lips.

Experimental Hypnotism; Effect on Circulation.—Berillon contributes an article on this subject to the "Journ. de Med. de Paris," describing the effect of hypnotism on several hysteric subjects, including those otherwise sound and those with functional circulatory disturbances. The sphygmograms show a remarkable influence of the circulation under the suggestion by the hypnotizer. The pulse and also the heart action could be retarded, accelerated or rendered regular at will, and the arterial tension modified. He also mentions that one of his friends, in a series of sphygmographic researches on himself, was alarmed to discover, after a time, marked arrhythmia, which, however, subsided so soon as he abandoned his researches. The close concentration of the attention directed by suggestion on certain organs, certain functions or certain parts of the body, leads to an actual polarization of the nerve force at this point. In one case of traumatic neurosis, with palpitations, arrhythmia and incessant tremor, all these symptoms subsided in the hypnotic sleep, as also the pronounced tachycardia in another case.

Ankylostomiasis in Cuba and Mexico.—Physicians in these two countries are awaking to the realization of the fact that the numerous cases of progressive anemia observed there and hitherto assumed to be malarial cachexia are, in reality, in many instances true ankylostomiasis. Agramonte describes his experience at Havana in the "Revista Med. Cubana," 1, No. 4. He found the *Uncinaria americana* in 16 cases of progressive anemia, 12 of which had been previously diagnosed as malarial anemia or cachexia and 3 as splenic anemia. He reports the clinical course of a few cases and illustrates the differences between this form of the uncinaria and the better known *U. duodenalis*. Manuell has had a similar experience in Mexico, having found 5 cases of ankylostomiasis in a brief time so soon as he commenced to examine for it. He remarks that these cases in the past have probably been cited as examples of the failure of quinin in certain manifestations of malaria. Two of his patients had been employed in Yucatan, 2 in the tobacco fields of the national valley and 1 came from a mining district. The trichocephalus accompanied the parasite in every instance, and in one case ascariides were also present. He mentions in his communication to the "Revista Médica" of Mexico, xiv, 11, that no one before has ever mentioned finding the trichocephalus in Mexico.

Aqueous Preparations of Menthol.—Blondel found that if menthol were given to a dog at the same time or shortly after an emetic dose of ipecac, there was no vomiting. Chauffard also noted that ipecac given with menthol failed to produce an emetic effect. This power to inhibit vomiting renders menthol very valuable in vomiting of reflex origin. Crésantignes gives several formulas in the "Journ. de Méd. de Paris" for preparing an aqueous solution of menthol. Ordinarily it does not dissolve in water. He uses the tincture of quillaja saponaria, 5 gm. to 3 or 5 cg. of menthol, adding, after they are mixed and the menthol is dissolved, 10 gm. of neutral glycerin in distilled water q. s. to make 125 c.c. This is taken by the tablespoonful. For local application in case of frontal headache, he applies a mixture of 15 to 30 cg. of menthol dissolved in 10 gm. of tincture of quillaja and distilled water to make 155 c.c. This is applied on a folded compress. After three to five minutes the compress is replaced by one dipped in pure water, after which the menthol compress is applied again, and so on. For a mouth wash and gargle he prefers the formula: menthol 10 to 20 cg. in 20 gm. tincture of quillaja and distilled water, or a saturated solution of boric acid q. s. to make a quart. The "Medical Standard" says that the first formula given "illustrates the absurdity of some of the prescriptions recommended in the journals for medicinal use. This mixture is to be used as a gastric sedative. The addition of the quillaja defeats this very end, since it is a decided, often a violent irritant; it is an active poison—a cardiac depressant, and a bill forbidding its

use as a foam-producing ingredient of soda fountain drinks has been introduced in at least one state. The art of the pharmacist has in this case over-shadowed the practical end of the physician, who, as a matter of fact, would find the simpler spirit of peppermint or peppermint water just as effective."

Queries and Minor Notes.

ANONYMOUS COMMUNICATIONS will not be noticed. Queries for this column must be accompanied by the writer's name and address, but the request of the writer not to publish his name will be faithfully observed.

CAUSE OF DEATH WANTED.

WATERTOWN, S. D., April 16, 1903.

To the Editor:—The following case has proved somewhat of a conundrum to me especially as no postmortem examination was allowed.

Cecil G., about 5 years of age, with a previous good history except a recent recovery from a mild attack of la grippe, was placed under chloroform anesthesia for the purpose of removing enlarged tonsils. The anesthesia was not complete, first, because the operation was expected to be of such short duration that it was not deemed necessary to take the time; second, in consequence of some danger to blood getting into the bronchi if reflex was entirely abolished.

One tonsil was easily removed, but some difficulty was experienced in getting the other engaged in the tonsillotome, principally in consequence of the contraction of the throat, and finally it was seized by a vulsella and removed with serrated scissors.

The child was immediately inverted to facilitate the flow of blood from the mouth, the child meanwhile struggling and crying with returning consciousness, when it suddenly stiffened and breathing and heart action ceased at almost the same instant.

Artificial respiration for over an hour, with subcutaneous injections of normal saline solution, failed to obtain any result. What would be supposed to be the exciting cause of death?

H. H. CLARK.

Correspondence.

Priority in Technic.

BROOKLYN, April 20, 1903.

To the Editor:—In the April 4 number of THE JOURNAL one of your contributors, in an article entitled "A Few Practical Points in the Technic of Nephrorrhaphy and Herniotomy, and a New Modification of Alexander's Operation," publishes, as his own, three operative procedures so long identified with the names of other surgeons that I can not forbear calling your attention to the fact.

In the second paragraph of his article your contributor states that he has employed the methods described since 1890, claiming originality, by implication, for the latter and dating the claim from the time mentioned. It may be true that Dr. Canac-Marquis was the first to apply the "crossed" or figure-of-eight suture in nephrorrhaphy, although, if my recollection serves me right, I used it in the winter of 1889 and 1890 in perineorrhaphy, nephrorrhaphy, herniotomy and for the closure of abdominal and other incisions in general. This part of the subject is unimportant, however, since this was the period of development of the possibilities of the "crossed" or removable layer suture in my work, and prior to any publication of my own on the subject, or, so far as I then knew, of the publications of others as well. Since, also, your contributor does not employ the figure-of-eight suture to anchor the kidney, but only to close the wound and since, further, he does describe its use in closing the wound of the herniotomy operation, I feel that failure to give credit for these applications of the method justifies the present contribution.

In the transactions of the American Surgical Association for 1891 there appeared my first communication relating to a removable layer suture, illustrated by woodcuts. This suture was again described, with a modification relating to the method of finally securing the ends of the suture. Still later the method was again described in illustration of its application to the closure of the wound of the abdominal wall following the operation for appendicitis in a "Treatise on Appendicitis," edition of 1894. In the 1900 edition of the latter work this suture, together with a still further modification of the

method of securing its ends, is again described and illustrated. Quoting from the first of these, I said:

It occasionally happens . . . that the different layers of tissue passed through in the course of the operation must be sutured separately in order to insure a good result. Here catgut will not answer, for the reason that it is dissolved or, at all events, weakened, before it has answered its purpose, and the sutured layers separate widely in spite of every care. To meet the requirements of these cases I have devised a suture which, while it serves all the purposes of a deeply buried layer suture, at the same time can be removed. I have called this the "crossed suture." Silkworm gut or silk may be used for this purpose, but the former is preferable. Both ends of the thread are threaded with needles of the proper curve, and both are passed beneath the lowermost or the deepest layer to be sutured, at sufficient distance from the edge to obtain a good hold thereon. As the thread is drawn through either edge of the layer, the loop is drawn taut beneath. The thread is now crossed, the needle which passed the latter through the outer edge crossing over the inner edge, and that which passed the thread through the inner edge going to the outer. The needles are now passed through the next layer as before, from below upward, securing a firm hold as in the first layer.

The threads are again drawn taut and crossed as before, each time of crossing securing a layer until the surface is reached, the last layer, of course, being formed by the skin itself.

While Dr. Canac-Marquis concludes his operation by the use of "a lead nickel-plated shield threaded and shotted," I have for some years attained the same result in a slightly different manner. Quoting from my book, "A Treatise on Appendicitis," 1900:

The crossed sutures are tied in pairs on either side of the wound. In order to guard against cutting of the skin from the silkworm gut, the sutures are tied over "bolsters" of sterile gauze or, what I find exceedingly convenient for this purpose, sections of thick-walled tubing of soft rubber (aspirator tubing) are used, the silkworm gut being passed through the lumen of the tubing. This method gives all the advantages of buried layer sutures, and yet places completely within control the length of time the sutures are to remain in place. Removal is accomplished by cutting the suture beneath the bolster on one side, and making traction on the other end of the thread.

While my first article was in the hands of the printer, and after I had employed the method of thus placing a removable layer suture in practically all the situations in which it is applicable, Well's translation of the first edition of Pozzi's work on gynecology appeared. I then became aware, for the first time, through a very scant reference to the matter by Pozzi (Vol. II, pp. 980, 981), that a similar suture had been described by Heppner for the perineum.

Taking up, next, Dr. Canac-Marquis' "Technic of Herniotomy for Radical Cure of Hernia": Bryant, as far back as 1889, advocated the use of the hernial sac itself to form a pad of resistance to intra-abdominal pressure. In the "Transactions of the New York State Medical Association," 1889, under the heading "Treatment of Hernia," Dr. Bryant says:

I can not close this portion of the subject without calling to your attention a somewhat newer method of operating that I have practised on three distinct occasions. It certainly has some features that commend it. . . . It is performed by weaving the sac, after its separation and its neck has been tied with catgut, into the pillars of the ring.

He then illustrates slits through which the sac is woven back and forth, "and finally the whole mass is quilted with deep stitches of catgut."

Though the details of Bryant's old operation, since abandoned, differs somewhat from your correspondent's, still the fundamental principle involved remains the same, and, "honor where honor is due."

In regard to Dr. Canac-Marquis' "New Modification of Alexander's Operation for Shortening the Round Ligaments," I would recall to your attention the fact that in the "Annals of Surgery," 1896, occurs a paper by Robert Abbe, entitled "Fixation of Round Ligaments in Alexander's Operation." After referring to a previous method of his own, Abbe says:

The idea occurred to me of utilizing the round ligament itself for the double purpose of sewing up the inguinal canal and securing the ligament immovably.

This, I found, was easily accomplished when the ligament was drawn out four inches, as was always readily done. The ligament was drawn on until the uterus was firmly held well forward. The needle and loop now drew the ligament through new punctures on either side of the canal, crossing it to become a continuous suture for the canal. Three such stitches completely closed the canal up to the spine of the pubis, and left more than an inch and a half of ligament over. This was turned back over the last portion of the wound and stitched to it by one fine catgut stitch.

The use of living tissue as a suture thus serves the double purpose of fixing the uterus immovably and binding the canal permanently.

This is identical with your correspondent's method, except in the minor details of closing the superficial structures.

Unless your contributor can substantiate his implied claim as the originator of these three points in technic by publications bearing date earlier than 1889 in the case of the hernia operation, 1891 in the case of the removable layer or figure-of-eight suture, and 1896 in the case of the modified Alexander's operation, the credit of the method of weaving the pillars of the ring together by means of the sac itself should be given to Bryant, that of a removable suture to Heppner and myself, and that of employing the round ligament to close the inguinal canal and at the same time secure the uterus in position in Alexander's operation to Abbe.

GEORGE RYERSON FOWLER.

Marriages.

CHARLES P. PINCKARD, M.D., to Miss Mary Losey Graff, both of Chicago, April 25.

J. JULIUS RICHARDSON, M.D., to Miss Dorothy Flynn, both of Washington, D. C., April 20.

HUGH G. NICHOLSON, M.D., to Miss Roberta Coleman, both of Charleston, W. Va., April 16.

TASSO OLIVER FELIX, M.D., Downs, Kan., to Miss Neva Lee Trusdle of Kirwin, Kan., April 16.

WILLIAM A. CAMPBELL, M.D., Chicago, to Miss Jeannette E. Halsted, at Muskegon, Mich., April 15.

Deaths.

Matthew Gardner, M.D. McGill University, Montreal, 1871, a member of the American Medical Association; chief surgeon of the Southern Pacific Railway system; a member of the California State and San Francisco County medical societies; eminent in his profession and universally beloved, died at the Southern Pacific Hospital, San Francisco, April 18, a few hours after an operation for appendicitis, aged 56. He leaves an estate of a value estimated at from \$75,000 to \$80,000.

Joseph S. Eastman, M.D. Missouri Medical College, St. Louis, 1878, one of the founders and first president of the Oakland College of Medicine and Surgery, and a member of the state and county medical societies, died at his home in Berkeley, Cal., April 19, after an illness of six months, aged 48.

Henry Clinton Bunce, M.D. Medical Institution of Yale College, New Haven, 1850, assistant surgeon of the First Connecticut Heavy Artillery in the Civil War, died at his home in Glastonbury, Conn., April 15, after a protracted illness, aged 78.

William Flood Smith, M.D. College of Physicians and Surgeons, Baltimore, 1889, and thereafter professor of anatomy in the same institution, died from pneumonia at the Protestant Infirmary, Baltimore, April 14, after a short illness, aged 38.

John M. Allen, M.D. University of Pennsylvania, Philadelphia, 1840, surgeon of the Fifty-fourth Pennsylvania Volunteers in the Civil War, died at his home in Chester, Pa., from paralysis, after a short illness, April 20, aged 84.

Charles L. Howard, M.D. Medical College of Indiana, Indianapolis, 1896, of St. Paul, Ind., was instantly killed in a collision between an automobile which he was driving and a street car at Indianapolis, April 18.

John F. Gaylord, M.D. Medical Institution of Yale College, New Haven, 1878, a member of the Samoset Medical Association, died at his home in Plymouth, Mass., April 14, after a lingering illness, aged 45.

Sherman Hartwell Chapman, M.D. College of Physicians and Surgeons, New York, 1869, a specialist in diseases of the throat and lungs, died at his home in New Haven, April 15, from heart disease, aged 59.

James M. Brown, M.D. Jefferson Medical College, Philadelphia, 1875, a member of the American Medical Association, died at his home in Philadelphia, April 11, after an illness of several months.

L. Hart Kitchel, M.D. University of Buffalo, 1870, died from concussion of the brain and spine, resulting from a fall two months before, at his home in Alden, N. Y., April 20, aged 57.

Andrew G. Selman, M.D. Transylvania University, Lexington, Ky., 1843, a surgeon in the Civil War, who retired from practice in 1893, died at his home in Indianapolis, April 17, aged 81.