During the winter months following the epidemic we observed in patients admitted to hospital few late complications and sequelae that could be attributed to preceding influenza and pneumonia. This observation was rather surprising, and contrary to expectation. In a few instances we have seen chronic bronchitis with cough and râles persist for fully six months after pneumonia, without much impairment of general health. Such signs naturally suggest a suspicion of tuberculosis, but this was apparently excluded.

The first fifty fatal cases were subjected to necropsy without selection and thereafter only occasional cases were brought to necropsy for special reasons. The findings were the same as reported from other camps, in general extensive, often confluent, bronchopneumonia with marked compensatory acute emphysema.

The bacteriologic and pathologic studies of the disease disclosed no facts of peculiar importance and determined nothing as to the real cause of the disease.

SUMMARY

1. The epidemic of influenza at Camp Upton in its clinical aspect and severity was the counterpart of the disease as reported from the other great camps in the northeastern section of the United States.

2. Secondary bronchopneumonia was the one predominant complication. Bronchopneumonia with its complications, chiefly empyema, was the sole cause of death. Empyema was encountered in only 1.76 per cent. of pneumonia cases.

3. The statistical tables and graphic charts are arranged to show, by reading the tables crosswise and the charts up and down, the ultimate outcome (complications and deaths) that resulted from the original daily admissions of influenza, calculated up to Jan. 1, 1919, irrespective of the date on which the complications or deaths happened to fall. This arrangement portrays the relative severity of the disease at all stages of its progress.

4. Chart 2, based on Table 2, shows the curves of pneumonia incidence rate compared with pneumonia mortality rate, expressed in percentage, resulting from admissions of primary influenza week by week through the epidemic. This chart brings out visually the interesting and unexplained fact that the pneumonia incidence rate increased while the pneumonia mortality rate decreased progressively and strikingly from the start throughout the course of the epidemic. This fact, if confirmed in other military camps, tends to invalidate claims of special therapeutic efficacy based on results obtained after the early part of the epidemic, during a period of natural decline of mortality.

5. The use of digitalis in large doses at the beginning of pneumonia, employed as routine practice, resulted in no observed harmful effects in any case, and proved of distinct benefit. The pulse remained slow and regular, even in fatal cases, to the end.

6. The early diagnosis of the type of bronchopneumonia secondary to influenza seldom rested on clear physical signs in the chest. The physical signs were evasive and slight and needed to be carefully sought for and combined with the other clinical data or the roentgen-ray findings in determining an early diagnosis. The criteria for diagnosis are described in some detail.

7. Toward the end of the epidemic of influenza there was added a new affection, occurring in epidemic form, that symptomatically closely resembled influenza, needed

careful differentiation, and led to much confusion and error in diagnosis—a peculiar form of infectious tonsillitis, characterized particularly by swelling of the tonsils and a membranous deposit on them. This affection has been the chief cause of admission to hospital from late November to the present time (April 1, 1919). A similar affection prevails widely in other places and is masquerading in vital statistics under the guise of influenza.

HYPOPHYSIAL DYSTROPHY IN HYDROCEPHALUS

AUGUST STRAUCH, M.D.

Attending Physician at the Cook County Hospital, Children's
Department

CHICAGO

Besides primary disturbances of the hypophysis itself, such as tumor, hypoplasia, etc., there are a number of intracranial conditions, either in the vicinity of the gland or more remote, that, directly or indirectly, may lead to interference with its function, resulting in the development of pituitary adiposity. Not only neighboring new growths, but also severe hydrocephalus with dilatation of the third ventricle, of whatever origin, may exert deleterious pressure on or obstruct the hypophysis and thus interfere with its normal function.

Harvey Cushing¹ remarks that secondary changes probably occur in the pituitary body in every case of increased intracranial tension, often with gross deformity and resultant functional disturbances which frequently elicit recognizable clinical manifestations. The scanty literature on the relation of hydrocephalus to the pituitary gland has been reviewed by the above named writer and in *THE JOURNAL* by Pollok.²

As an instance of conspicuous manifestations of hypophysial dystrophy in connection with a hydrops ventriculorum, the following case of a patient admitted to the Cook County Hospital, Oct. 15, 1918, may be reported:

REPORT OF CASE

History.—The girl, F. M., aged 8 years, 10 months, had always been a strong, bright child and had had a somewhat large head. When 16 months old, she is said to have weighed 35 pounds. After December, 1917, the girl often came home from school with wet clothes, being unable to hold the urine long. She drank much water and had to urinate frequently. She had, during the winter of 1917-1918, daily attacks of supra-orbital headache, sometimes accompanied by vomiting, with occasional dizziness. Her appetite grew ravenous, especially for bread. Polydipsia and polyuria lasted up to the time of admission to the hospital.

In February, 1918, her body weight, which was then 72 pounds, began rapidly to increase, and she became obese. At first irritable and "nervous," she soon became apathetic and sleepy, quite in contrast with her previous alertness, and she ceased making progress in the school. She tired very easily, and had difficulty in keeping her balance; walking became difficult, and she finally had always to be led by her mother. After the end of February, she was unable to attend school. The upper extremities became weak, trembling, awkward and ataxic. During that year she grew rapidly—"one could see her grow;" there also occurred a marked aggravation of preexisting genua valga and a striking enlargement of the head, so that it became difficult to purchase a hat large enough for her.

1. Cushing, Harvey: *The Pituitary Body and Its Disorders*, 1911.

2. Pollok, L. J.: Hypopituitarism in Chronic Hydrocephalus, *J. A. M. A.* 64: 395 (Jan. 30) 1915.

Physical Examination.—Physical examination revealed a well developed, obese girl (Fig. 1), weighing 95 pounds, body length 135 cm., distance of lower edge of symphysis from floor 66.5 cm. There was a marked development of adipose tissue, especially on the neck, in the region of the angulus mandibulae, of the pectoralis muscles, on the lower part of the large protruding abdomen, and on the mons veneris, which was separated from the abdomen by a deep furrow. There was also around the hips and on the upper part of the thighs an excessive amount of adipose tissue. The circumference of the chest in the nipple line was 74 cm., of the abdomen in the umbilical line 83 cm., and around the trochanters 82 cm.

The general integument and mucous membranes were of a healthy color; the skin was very soft, delicate, smooth and without perspiration. The body was covered with a very fine lanugo, that on the outer aspect of the forearm being more developed. The labia majora were much developed, large and covered with dense lanugo and a few darker, longer and thicker hairs. The hair of the scalp was abundant and long. The skull was large, its horizontal circumference being 59.5 cm.; the measure from the meatus auditorius of the one side to that of the other was 39 cm. The skull in the region of the anterior fontanel was thinned, distinctly yielding on pressure, and tender.

The face was very fat. The veins of the right upper lid and temple were bulging, those of the left upper lid less so. There was slight exophthalmos. The pupils were large, reacted at times sluggishly, at other times promptly. The eye fundus was normal. The hands were graceful, somewhat pudgy, very soft and smooth, the fingers being somewhat tapering. The nails of both thumbs had a normal lunula, the index fingers a hardly visible lunula, while the other fingers had no crescent at all. The hands and feet were cold. Genua valga and pedes valgi were present.

There was motor ataxia of the arms, which on some days was very pronounced, on other days demonstrable only with finer tests. The gross muscular power of the arms and legs was much reduced. There was no spasticity. The patient was able to sit freely without ataxia, but was unable to stand or walk without support, and swayed and fell if not held up. The general asthenia forced her to spend all her time in bed or on a chair. The patellar reflexes were increased, that on the right side being more pronounced. Babinski's reflex was positive. There was ankle clonus, especially on the right side. There was involuntary micturition. The Wassermann test was negative. The systolic blood pressure was 95, diastolic 65. The pulse was between 100 and 110. The intellectual faculties were much impaired, and the school-learning was very deficient. On three occasions, 100, 150 and 210 gm. of glucose were given by mouth without causing glycosuria.

The roentgenographic examination (Fig. 2) revealed a greatly enlarged, shallow sella turcica and reduced clinoid processes. The stereoplates of the entire skull showed a hydrocephalus with open coronal suture and marked dilatations of the frontal bone.

The girl gained 3 pounds in the first two weeks of observation (98 pounds). The reflexes were changeable, the ankle clonus often being absent. Later incontinencia alvi supervened. By the end of November, bilateral optic neuritis was found, the vision of both eyes being 20/30. The hairs on the labia majora increased somewhat in size and number. The sensorium was free, the girl friendly and kind; however, she was quiet and often languid and drowsy. During the last three months of her life she had periods of marked apathy, even torpor, when suffering from measles, otitis media suppurativa acuta and bronchopneumonia, to which she succumbed, March 2, 1919. She grew 3 cm. in height after

admission to the hospital and lost in body weight during the last few weeks.

Postmortem Findings.—From the pathologic report, for which I am indebted to Dr. F. H. Stangl, resident pathologist at Cook County Hospital, the following features may be mentioned here:

The calvarium measures 3 mm. in thickness; its sutures are only fibrous. The brain is large, owing to an increased amount of ventricular fluid. The convolutions are markedly flattened. The brain is 16.1 cm. wide, 20.3 cm. long and 10.1 cm. high, and, with its contained fluid, weighs 2,190 gm. The infundibulum is 12 mm. long and kinked on itself. The hypophysis is but slightly flattened and measures 7 mm. by 11 mm. by 16 mm. Serial coronal sections of the brain reveal greatly dilated ventricles with reduction in the brain substance surrounding them. The right lateral ventricle measures 4.5 cm. transversely and 6.5 cm. vertically, the left one 3.8 and 6.5, and the third ventricle 2.0 and 2.5 cm., respectively. There is a shifting of the parts of the brain, especially of the occipital half, with resulting malposition. The aqueductus Sylvii is entirely closed by compression, and the right hemisphere of the cerebellum has undergone decrease in size from pressure.

The internal genitalia reveal no gross changes. The body of the uterus is 3 cm. long, 1.5 cm. wide at the fundus and 0.5 cm. thick. Sections of the hypophysis stained with eosin-haematoxylin have been examined microscopically by Dr. G. B. Hassin with the following result: The three portions of the organ were distinctly shown; they are surrounded by a connective tissue membrane consisting of collagen fibers with well stained oblong nuclei. The anterior or glandular part exhibited numerous acini, some of which appeared more densely stained than others. The epithelial cells of the acini or alveoles showed a large, well developed granular cytoplasm, containing an eccentric nucleus. Some epithelial cells were unusually large, exhibiting two or more nuclei. In other acini which were quite numerous the epithelial cells were very pale, homogeneous, irregular in shape, but with a distinct, well stained nucleus. Such cells often merged together forming a large colloidal mass containing several nuclei. The colloid often totally obliterated the acini, or filled up their lumina, sometimes appearing as very large diffuse masses or small globules freely scattered in the interalveolar spaces. The blood vessels were dilated and greatly congested; some were partially or totally filled with a colloid substance. There were to be seen foci

of hemorrhages in the tissue of the capsule and between the latter and the glandular tissue. The pars anterior of the hypophysis was in the center of the section—isthmus-like—continuous with the posterior portion, which appeared reticular, the stroma in the area of junction showing much wider meshes than in the rest of the posterior lobe. There were to be seen a few congested capillaries and still fewer colloidal bodies.

The lateral portions between the anterior and the posterior lobes of the hypophysis (of the pars intermedia) were occupied by strikingly large cuneiform clefts or rather cystic formations lined with epithelial cells and entirely filled with colloid. The latter contained cellular masses with distinct nuclei, situated here and there, but as a rule near the margins of the cystic clefts. These two colloidal formations can be seen with the naked eye, the larger one measuring 1.5 mm. in width and 3 mm. in length. The pathologic changes can thus be summed up as subcapsular hemorrhages, excessive hyperemia and vascularization of the anterior lobe, hyaline or colloidal thrombosis and excessive amounts of colloidal acini. The findings so far give no exact idea of the presence or absence of any pathologic process that caused the hydrocephalus. Further studies

Fig. 1.—Hydrocephalus, accompanied by hypopituitarism.

are being carried on, and a more complete report, especially of the histopathology of the brain is forthcoming, being in preparation by Dr. G. B. Hassin.

COMMENT

Certain features of the anamnesis of the case are suggestive and a part of the clinical syndrome is quite characteristic of hypophysial dystrophy. The manifestations of the latter, however, must be distinguished from those of internal hydrocephalus. The rapid development of adipositas nimia, with eunuchoid localizations, and the high tolerance for glucose, indicate constitutional, metabolic changes—such as are observed in functional deficiency of the posterior lobe of the hypophysis. The delicate skin, the lack of perspiration and the arterial hypotension likewise point to this organ. On the other hand, the abnormal increase of the size of the skull, with thinning of the region of the anterior fontanel, and the marked remissions and exacerbations of the nervous disturbances, leave no doubt about the existence of hydrocephalus internus. Its causal agency, however, cannot be ascertained clinically beyond dispute, the differential diagnosis considering an "idiopathic" hydrocephalus, or rather the exacerbation of a previous hydrops ventriculorum, and a possible distant obstructive tumor without focal manifestations.

The presence of the hydrocephalus gives satisfactory explanation of dystrophy, through the assumption of a pressure lesion of the hypophysial apparatus, and, with the absence of optic nerve disturbances, even as late as at the time of admission, militates against the assumption of a tumor of the hypophysis itself. The radiologic findings are in accord therewith, the shallow dishlike dilatation of the entrance of the sella turcica being merely a partial manifestation of the cranial changes that are due to the general increased intracranial tension.

Cushing at first explained the pituitary deficiency in hydrocephalus as due merely to the dilatation of the third ventricle, with consequent flattening of the gland. Since the demonstration of a possible discharge of the posterior lobe secretion into the cerebrospinal fluid, he sees in the stasis of the cerebrospinal fluid an equivalent in its effect to an experimental obstruction of the stalk, or to a compression by an immediately superimposed infundibular tumor. In connection with his ideas it is noteworthy that in our case a slight flattening of the gland and kinking of the infundibulum were found postmortem, according to the report given me.

Polyuria and polydipsia, which for a time were so marked in our case, have not infrequently been noted by other authors as a temporary symptom in hypophysial dystrophy, and in a few cases a serious diabetes insipidus was the predominant feature. It also has been encountered for a period as an after-effect of

surgical operations in the hypophysial region, even of simple sellar decompression (Cushing). The occurrence of polyuria observed by Schaeffer³ in bloodless, mechanical or chemical irritations of the hypophysis, and the well established diuretic effect of the extract of the nervous lobe, combine to indicate the cause of this clinical symptom to be an increased secretion resulting from irritative conditions, such as pressure, inflammation, etc.

Thus a state of initial stimulation of the posterior lobe in our case may be assumed preceding its inhibition or obstruction. The persistence of polydipsia, however, during the period of well developed adiposity, the latter indicative of hypofunction, cannot, of course, be attributed to a simultaneous hyperfunction; it must be explained on a different basis. We may consider it merely a habit formed during the illness. Several circumstances indicate this; the excessive drinking of water, for instance, had entirely lost its initial compelling character and its degree; it became easily controlled by persuasion, was subject to occasion and absent during the stay in the hospital.

On the other hand, marked polyuria has been observed also in tumors (gumma) in the vicinity of the hypophysis and in various affections of the brain stem (chiasma, subthalamic region, etc.); it likewise has been produced by irritation of certain parts in the medulla oblongata.

The polyphagia, repeatedly mentioned in the literature, has been attributed to posterior lobe insufficiency. If we interpret the rapid growth of the patient, the advanced development of the labia majora, and the growth of hair on the labia majora in the light of the present preponderant

Fig. 2.—Roentgenogram of sella turcica and clinoid processes in a case of hypopituitarism.

evidences of the function of the parts of the hypophysis, we may possibly have here an example of activation and hyperfunction of the anterior lobe, combined with posterior lobe insufficiency (dyspituitarism, Cushing). This unusual type differs thus from the common type (Froehlich's⁴) of hypophysial dystrophia adiposogenitalis, which is characterized by undergrowth and sexual hypoplasia due to simultaneous anterior lobe inhibition. Somewhat similar instances of dyspituitarism are those of Neurath⁵ and Case 34 of Cushing.

The disturbance of micturition and defecation finds its ready explanation in the large hydrocephalic effusion, with pressure lesion of the corresponding cerebral districts. Bladder disturbances, which, according to Frankl-Hochwart and Froehlich,⁶ are not infrequently observed in hypophysial dystrophy, have been explained mostly by compression of the pes pedunculi by a grow-

3. Schaeffer, E. A.: Quoted from Falta.

4. Froehlich: Wien. klin. Rundschau, No. 45, 1901.

5. Neurath: Ueber Fettkinder, Wien. klin. Wchnschr., No. 2, 1911.

6. Frankl-Hochwart and Froehlich: Arch. f. exper. Path. u. Pharmacol. 63: 347, 1910.

ing tumor of the pituitary body. However, both authors consider also the possibility of the reduced function of the posterior lobe and the lack of the pituitrinum infundibulare being a responsible factor.

The presence of a marked degree of hydrocephalus in our case renders it futile to distinguish the psychic alterations of hypophysial origin from those due to the ventricular effusion.

The supervening optic neuritis would have made surgical interference urgent. Balkenstich operation after Anton and Bramann was contemplated as the simplest décompression operation for the hydrocephalus, with the possibility in view of thus relieving and reactivating the pituitary body. The intercurrent diseases, however, rendered it impossible.⁷

4557 Broadway.

THE SURGICAL TREATMENT OF RHINOPHIMA

JOSEPH E. FULD, M.D.

Attending Surgeon to Park Hospital; Instructor in Operative
Surgery, College of Physicians and Surgeons, Columbia
University; Assistant Visiting Surgeon,
City and Gouverneur Hospitals

NEW YORK

Rhinophima is the most advanced stage in the development of acne rosacea, characterized by marked hypertrophic changes in the cutaneous and subcutaneous tissues of the nose, producing a firm, reddish or purplish lobulated, and, in extreme cases, pendulous tumor masses. This condition is frequently so disfiguring and repulsive as to prevent the person afflicted with it from earning a livelihood or mingling in society, and it is in just such cases that surgery offers the only means of relief.

Fig. 1.—Incision made in removing the left half of the growth: A B, midline incision made down to the cartilage; A C, lateral incision encircling the growth.

The pathologic changes which lead up to this condition may well be divided into three stages. The first

7. In addition to the references already given, the following will be found of interest.

Falta, W.: The Ductless Glands (translation from the German).
Strauch, A.: Infantilis, Am. J. M. Sc., 148: 247, (Aug.) 1914.

stage is marked by a simple venous engorgement. The second stage shows the veins and the capillaries permanently enlarged. In the third stage, there is an enormous hyperplasia of the connective tissue elements

Fig. 2.—The procedure for removing the growth.

of the skin, and the sebaceous glands are so enlarged that the nose presents a honeycombed and unsightly appearance.

TREATMENT

It is at this stage that the electric needle, ointments and the roentgen ray are no longer of therapeutic value, but where surgery is the only cure. When the tumors are pedunculated, a condition which frequently occurs, their removal requires no special description. There is, however, a marked difference in the operative procedure involving extensive hypertrophy without pedunculation, as in the case of the patient on whom I operated, as shown in the accompanying photographs.

Instead of removing the tumor masses in one piece, in my recent cases, I excised first one half then the other half of the growth, making the operation much easier to perform.

OPERATIVE PROCEDURE

Under general anesthesia, an incision is made down to the cartilage in the median line of the nose, as shown in Figure 1, from A to B, and a second incision all around the growth from the midline outward from A to C. The forefinger of the left hand is now inserted into the left nostril, with the palmar surface forward, as a guide. The median edge of the tumor mass is grasped with mouse-tooth forceps (assistant to hold these), and all the diseased tissue, including the entire left lower half of the nose, with the whole thickness of the skin, is dissected off with a knife, the bony and cartilaginous framework of the nose being exposed. The hemorrhage is controlled by pressure with hot pads or forceps and ligations, if necessary. The foregoing steps are repeated on the right side. It is important to leave a little skin near the opening of the