Ryerson for spina bifida, double club-foot and double retroverted knees; complete recovery followed except impaired function of one leg, which is improving. One case of complete exstrophy of the urinary bladder in a female child with absence of the pubic arch; this child died soon after birth. Three cases of facial paralysis, one congenital; this child died when about 2 years of age probably from a cerebral growth; the remaining two children, in whom the facial paralysis was caused by forceps, recovered.

There were seven cases of hemorrhage of the newborn. Sterile horse-serum was used in one of these, after a large bowel hemorrhage. The child recovered. Of the six that died, one lived about eight months.

There was one case of anencephalus, and one case of imperforate esophagus, in which the child died of starvation on the tenth day after birth. A catheter would pass $4\frac{1}{2}$ inches from gums to the end of the proximal portion. From the stomach the esophagus led upward into the trachea.

SUMMARY OF MATERNAL RESULTS IN 1,110 DELIVERIES

	No.	Per Cent.
Normal	953	85.4
Artificial	163	14.6
Lacerations:		
No perineal lacerations	676	60.5
Lacerations of the first degree	56	5
Lacerations of the second degree	388	34.8
Lacerations of the third degree	10	.9
Maternal classification:		
Primiparas	404	36.2
Multiparas	712	63.8
Presentations:		
Head	1064	95.3
Breech	36	3.2
Transverse	5	.44
Face	5	.45
Positions: Head and Transverse:		
Left occipito-anterior	802	71.8
Right occipito-anterior	211	18.8
Right occipito-posterior	53	4.7
Left occipito-posterior	3	.26
Breech:		
Left sacro-anterior	22	1.9
Right sacro-anterior	14	1.2
Face:		
Left mento-anterior	3	.27
Right mento-posterior	2	.18
Convalescence:		
Good	1	90.82
Death from puerperal sepsis	1	.08
Post-partum hemorrhages	16	1.43
Adherent placenta	7	.6
Placenta prævia centralis	2	.18
Eclampsia in labor	3	.27

(These three cases were: one ante-partum, one intra-partum, and one post-partum).

Phlegmasia alba dolens, severe type	3	::
Phlegmasia alba dolens, mild type	4	::

In 1905 just prior to this series I had a rare specimen in a primipara. Following the birth of a normal infant, came a "fœtus amorphus anideus" a little over 4 inches long and $2\frac{1}{2}$ inches thick, with no head or extremities. It had a mass of black hair like that of a scalp, which covered about one-quarter of its surface. Dr. J. Clarence Webster reported this specimen and stated that it was the second of the kind he had studied.

The total number of infant deaths was thirty-eight, or 3.4 per cent. Of the deaths eleven occurred ante partum, eighteen post partum, and nine intra partum.

The care of this series of cases has required more sacrifice of time, labor, patience and comfort than a similar series would have required in a maternity hospital, and probably more than is generally given. But since most cases must be attended in the home, it is imperative that the profession should provide as high-grade care for the parturient woman and her child as any department of medicine can give, in order to save lives and unnecessary illness.

6517 Stewart Avenue.

BLOOD-CHANGES CAUSED BY THE HYPO- DERMIC ADMINISTRATION OF THE CANCER PROTEID*

J. WALTER VAUGHAN, M.D.

DETROIT

During the past six years, over two hundred cases of recurrent or inoperable malignant disease, which have come under my observation, have been treated by some method of vaccine therapy. Two forms of vaccine have been used in this work—namely, cancer residue and a vaccine made of the cancer-cell in its entirety. The preparation of cancer residue has been described in previous articles, so that it is sufficient to state that this is the water-soluble portion of the cancer-cell which is left after splitting off the toxic radical by means of heating the cancer-cells in a 2 per cent. solution of sodium hydroxid in absolute alcohol.

Cancer-cell vaccine is prepared by grinding cancer-tissue finely in a meat-grinder, after which it is rubbed up as a suspension in alcohol in a sterile mortar. Next, it is rubbed through a very fine-meshed wire sieve, the alcohol filtered off, and the collected cell substance air-dried. This is weighed and then placed in normal saline solution, making a 2 per cent. cell suspension. To this 0.5 per cent. phenol is added for the purpose of rendering it less likely to become contaminated. In the preparation of both residues and vaccines it has been found that satisfactory results can be obtained only when the tumor is of firm consistency and without necrotic or infected areas. The average injection of a 1 per cent. residue is from 5 to 20 minims; that of a 2 per cent. cancer-cell emulsion is from 5 to 10 minims.

When this work was first begun, it was based on the belief that it is possible to compel the human body to form specific ferments for the splitting up of any given proteid by the introduction of that specific proteid, as such, in a dead form, within the human body.

The most natural place to look for the formation of such a ferment is in the white blood-cells. For this reason daily differential leukocyte counts were made from the beginning of the work. Over 20,000 such counts have been made by my assistants and myself so that conclusions drawn from this portion of the work cannot be much in error. In the early stages of this work it was ascertained that those cases in which there was a marked decrease in the percentage of polymorphonuclear leukocytes, with a corresponding increase in the percentage of large or small mononuclear leukocytes, received benefit from vaccine injections, while cases in which the percentage of polymorphonuclear cells remained high, or were influenced but little, were but slightly benefited, or not at all. All cases treated were inoperable or recurrent after operation, so that by far the majority were absolutely hopeless at the start. In approximately 20 per cent. of these cases all visible signs of malignant tissue disappeared. This disappearance, however, has not always been permanent and in my cases treated by the vaccine method, which have been without treatment for from two to four years, the percentage of non-recurrences is reduced to ten. A review of these cases shows that the disappearance of evidences of tumor-cells depended, first, on the relative proportion of mononuclear leukocytes present, and sec-

* Read in the Section on Pathology and Physiology of the American Medical Association, at the Sixty-Third Annual Session, held at Atlantic City, June, 1912.

* From the Chase Cancer Research Fund, Harper Hospital.

Chart 1.—Rabbit, Cage 340; 5 minims of 1 per cent. sarcoma residue injected subcutaneously at 8 a. m., 9 a. m. and 10 a. m. The line designated as P indicates, in this and the following charts, count of polymorphonuclears; S indicates small and L large mononuclears.

ond, on the amount of cancer-tissue present. No estimate of the value of the use of a vaccine could be given by the microscopic picture of the tumor-cells. In other words, the malignancy of any given tumor is dependent on the bodily resistance of the patient to the growth of cancer-cells, and not on the nature of the growth itself. Time is too short to permit my going into this interesting branch of the subject in detail, but other experiments, which I shall mention later, tend to corroborate this view; suffice it to state that, in animal experiments, the chemical nature of the tumor-cells must have some influence, since a tumor can be trained to "take" almost invariably. My contention, however, is that the pathologist is not always able to state the degree of malignancy of a given tumor by an examination of it, but that a differential leukocyte count will give more accurate information on the subject.

Since a vaccine treatment would benefit only a very small percentage of cancer sufferers the next step was to see if it was possible to produce an active ferment for the splitting up of malignant cells in the bodies of animals, which, in turn, might be available in the treatment of the cancer patient.

For this reason sheep and rabbits were injected intravenously, intra-abdominally and subcutaneously with both cancer residue and cancer vaccine and frequent blood-counts made. All told, over 500 animals have been used, and the blood changes have been the same in all except about ten animals. Of these two were proved to be pregnant at the time of injection, which may have had some influence on the usual reaction. A glance at the accompanying charts will show that while the percentages of polymorphonuclear and small mononuclear leukocytes are not affected with any degree of regularity,

Chart 2.—A, rabbit given subcutaneous injection of 0.5 c.c. rectal adenocarcinoma (Morris); B, rabbit injected with 0.5 c.c. small round-celled sarcoma; C, rabbit given intraperitoneally 0.5 c.c. rectal adenocarcinoma (Morris); D, rabbit given intraperitoneally 0.5 c.c. rectal adenocarcinoma (Morris).

the proportion of large mononuclear cells is invariably increased from 100 to 400 per cent. within from twenty-four to forty-eight hours. This increase, however, is of short duration and recedes with rapidity after reaching its height.

Chart 1 shows the average blood change obtained through the injection of an active residue. The preparation used was made from a small round-celled sarcoma of the mediastinum. Three subcutaneous injections of 5 minims each of a 1 per cent. residue were given at hourly intervals, the first blood-count being made before the injections were commenced. The second count made seven hours later, showed an increase in polymorphonuclear cells from 21 per cent. to 37 per cent. and a corresponding decrease in small mononuclear leukocytes. The third count made twenty-five hours after the first showed the characteristic increase of large mononuclear cells which are here registered at 27 per

Chart 3.—Sheep injected with 0.5 c.c. sarcoma residue.

cent. with a normal at the first injection of 10 per cent. Two and one-half hours later the percentage of this form of cell had returned to normal, which was again followed by a slight increase. It is probable that the highest registration of large mononuclear cells occurred between the second and third counts.

Chart 2 represents four rabbits which were given 0.5 c.c. each of cancer-cell vaccine by different methods. The first was given a single subcutaneous injection of rectal adenocarcinoma. The fourth represents the same amount of the same tumor given intraperitoneally, as does also the third. The second shows the injection of vaccine made from a rapidly growing round-celled sarcoma of the neck. From these charts it can be seen that the intraperitoneal method of injection gives a more rapid reaction than the subcutaneous.

Chart 3 represents the blood changes occurring in a sheep following the injection of 0.5 c.c. of sarcoma residue.

Chart 4 shows more frequent differential counts following repeated subcutaneous injections.

Chart 5 illustrates daily counts following a single intraperitoneal injection of cells from a breast adenocarcinoma. This is rather an extreme reaction inasmuch as the proportion of large mononuclear cells is increased from 6 per cent. to 35, about 500 per cent.

Chart 6 shows a very slight reaction following the injection of a breast-cancer residue which has almost lost its activity.

Chart 7 and A in Chart 8 illustrate two rabbits giving the average reactions. B in Chart 8 represents

Chart 4.—A, rabbit injected subcutaneously with 5 minims of 1 per cent. s. r. sarcoma at 12 m., 2 p. m. and 3 p. m.; B, rabbit injected subcutaneously with 0.1 c.c. of 2 per cent. mixed residue at 12:15 p. m., 2 p. m. and 4 p. m.

Chart 5.—Rabbit given a single intraperitoneal injection of cells from a breast adenocarcinoma.

Chart 6.—Rabbit injected with a breast-cancer residue which has almost lost its activity.

the effect of giving an intraperitoneal injection of 10 c.c. of breast-carcinoma cell-emulsion into a rabbit sensitized to sarcoma vaccine, the injection being given when the percentage of large mononuclear cells was at its highest point.

In order to ascertain just what bearing this change in percentage of large mononuclear leukocytes had to the formation of a specific ferment, several rabbits were sensitized to the cancer-cell and then at varying percentages, 10 c.c. of cancer-cell emulsion were injected intravenously into the animals. In unsensitized rabbits there was no noticeable effect. Rabbits with a percentage of above 30 of large mononuclear cells usually died within one to three hours, and rabbits with a lower percentage but with a marked increase were made very sick, but recovered.

Sickness is immediate after injection; the animal at once falls on its side and begins violent scratching. The respiration is labored and, when death ensues, always stops before the heart-beat. Rabbits which die from a dose after being sensitized, however, usually have a stage of apparent rest from one-half to three hours after the first stage of excitability, which lasts

for from three to five minutes. A possible explanation of this, in view of work that I shall mention shortly, is that the first stage of excitement is due to the destruction of cancer-cells and consequent liberation of their toxic radical by the specific ferment present in the blood-serum, while the fatal result which follows later is due to the reaction between the cancer-cells and the large mononuclear leukocytes.

The increased percentage of large mononuclear leukocytes is but a transitory affair, however, which usually lasts from four to ten hours, and it is impossible to produce fatal results in rabbits by intravenous injections of cancer-cell emulsion after this stage is passed. Consequently, I have applied the term "transitory sensitization" to this phenomenon.

If we bleed a rabbit at the height of this transitory sensitization and obtain the serum, this, when mixed with cancer-cell emulsion and incubated for one hour, will produce marked symptoms of poisoning when injected intravenously into a normal rabbit. The severity of symptoms depends on both the amount of cancer-cell emulsion and the serum. Given 5 c.c. of cancer-cell emulsion and 10 c.c. of the serum, a rabbit of from 500 to 1,000 gm. will be very sick but will invariably recover.

With this experiment in mind injections of both sheep and rabbit serum were given in twelve cases of inoperable carcinoma, the serum being obtained when the percentage of large mononuclear cells was at its highest point following injections of small amounts of cancer proteid into the animals. Of these twelve cases, eight were given rabbit serum and four sheep serum. Injections of from 1 to 5 c.c. were made every second or third day. Of the patients treated with sheep serum, one became sensitized after three weeks. Two others developed an acute nephritis.

Chart 7.—A, Rabbit 54 Br., given 5 minims of 1 per cent. sarcoma at 1 p. m., 2 p. m. and 3 p. m.; B, Rabbit 54 Bl., given 5 minims of 1 per cent. sarcoma at 1 p. m., 2 p. m. and 3 p. m.

Of the eight given rabbit serum four developed acute nephritis with blood and albumin in the urine at the end of four weeks. While in all cases there seemed to be an improvement in the condition of the malignant growth, yet the use of the serum was discontinued, since in over 50 per cent. of the cases in which it was tried the serum complications rendered it impracticable.

The next step was to ascertain whether the specific ferment could be removed from the large mononuclear leukocytes in such a way as to be free from the dangers which were attributed to the serum. For this purpose rabbits were sensitized to the cancer-cell and, when the percentage of large mononuclear leukocytes was at its highest point, the blood was collected under sterile conditions in 0.33 $\frac{1}{3}$ per cent. acetic acid. This was next centrifugated until the leukocytes were thrown down and the supernatant fluid decanted. The remaining

mononuclear leukocytes was 25 or above, would, when injected intravenously into a rabbit of from 500 to 1,000 gm., kill within from one to five minutes. This is well illustrated in the accompanying table.

TABLE SHOWING RESULTS OF INJECTING INTRAVENOUSLY INTO RABBITS CANCER-CELL EMULSION PLUS LEUCOCYTE EXTRACT INCUBATED ONE HOUR

Animal	Weight, Gm.	Rabbit Leukocyte Extract	+ Vaccine	Result
343 G.	046.5	10 c.c. carc. 100....	5.0 c.c. 100	Died, 1 min.
107 G.	750	10 c.c. sarc. 51 res..	5.0 c.c. 100	Died, 1 min.
157 W.	1,500	10 c.c. sarc. 100....	5.0 c.c. 100	Very sick one-half hr.; rec.
53 Bl.	1,050	7 c.c. sarc. 51 res..	5.0 c.c. 100	Died, 5 min.
96 G.	1,317	7 c.c. sarc. 51 res..	2.5 c.c. 103	Died, 4 $\frac{1}{2}$ hr.
251 W.	645	10 c.c. sarc. 51 res..		Not sick.
251 Br.	842	15 c.c. carc. 100....		Not sick.
167 W.	700	10 c.c. normal saline.	5.0 c.c. 100	Not sick.

Chart 8.—A, Rabbit 167 W., given 5 minims of 1 per cent. sarcoma at 8:30 a. m., 9:30 a. m. and 10:30 a. m.; B, Rabbit 54 T., given 5 minims of 1 per cent. sarcoma at 1 p. m., 2 p. m. and 3 p. m.; at 12:15 p. m. 10 c.c. of breast carcinoma-cell emulsion (Mel.) was injected intraperitoneally; rabbit died, 8-10 hours.

leukocytes were next placed in a sterile mortar and mixed with a sufficient quantity of sterile quartz sand to cover. This was then ground up with vigor for fifteen minutes with the frequent addition of sterile normal salt solution until five times the volume of the leukocytes obtained had been added. Next, this normal saline extract was separated by passing through a Berkefeld filter, and tests were made for the presence of the specific ferment by adding varying amounts of the leukocyte extract to cancer-cell emulsion, incubating for one hour and then injecting intravenously into rabbits. Through repeated experiments it was ascertained that 5 c.c. of 2 per cent. cancer-cell emulsion, plus 10 c.c. of leukocyte extract, which was obtained when the percentage of large

The entire mass of leukocytes, when removed from the filter and mixed with 5 c.c. of cancer-cell emulsion and incubated one hour, failed to have any effect on a rabbit when injected intravenously, thus showing that most of the specific ferment was soluble.

Again, 10 c.c. of the leukocyte extract, prepared in the same manner, from a normal rabbit which had not been sensitized, added to 5 c.c. of cancer-cell emulsion and incubated one hour, produced no effect when injected intravenously into a normal rabbit. This would tend to show that the sensitized animal possesses some specific chemical substance which reacts with cancer-tissue which the normal rabbit does not possess. That this substance is specific is further shown by the fact that 10 c.c. of leukocyte extract from a sensitized animal, plus 5 c.c. of a 2 per cent. normal-skin vaccine, incubated one hour, produced no effect when injected intravenously into a rabbit of 450 gm.

The same chart shows that sarcoma residue or vaccine sensitizes to carcinoma as well as sarcoma, and vice versa, so that the probable conclusion is that the chemical change within the cell is the same for both sarcoma and carcinoma.

While it is not my intention at the present time to make mention of the use of leukocyte extract in the treatment of cancer, yet, because of the added danger to the patient and also because of the added proof of specificity, attention should be called to one fact that tends to show the presence of a specific ferment.

Even as small an amount as 1 c.c. or more of leukocyte extract when injected directly into the tumor may cause sudden and severe symptoms. In four cases, when this procedure was adopted, I have had the patient complain within from one to five minutes of difficulty of respiration. Next, he would lose consciousness, which would be accompanied by rather violent muscular twitchings and lowered pulse-rate. This stage would last from five to

ten minutes and would be followed by a period of rest, from fifteen minutes to one hour in duration, which in turn would be followed by a violent chill, and temperature ranging from 103 to 106 F. This would last from one to six hours and would be followed by from twenty-four to forty-eight hours of extreme exhaustion. At no time have I obtained such a reaction, even with doses of 5 c.c., when given subcutaneously away from the tumor, although a chill from one to three hours after injection has been observed.

It is my desire now, however, to call attention only to the animal experiments as described above. These have occupied two years' time and appear to throw some light on the cancer problem. I do not wish to create an impression that I advocate the use of sensitized leukocyte extract in the treatment of malignant disease, or that its use will be of any benefit to the sufferers from cancer. I have used it only in absolutely hopeless cases and only over a period of six months; consequently, any report of the subject from this point of view must be reserved for another year at least.

The report of the animal experiments is given, however, with the hope that those who are interested in this branch of cancer study will either corroborate or disprove the work, since it is only through the help of many that the truth will eventually be found.

In addition, I would state that a residue or vaccine remains active only as long as it is capable of producing blood changes in the animal injected. I have had a residue remain active for three years. The vaccine made from the cell in its entirety is not so stable, lasting from one week to six months. As a consequence I prefer to keep my cancer-tissue in alcohol and prepare only small amounts of vaccine at a time.

Washington Arcade.

ABSTRACT OF DISCUSSION

Dr. J. W. McCONNELL, Philadelphia: What does Dr. Vaughan mean when he speaks of the impossibility of pathologists telling much about the activity of the tumor—does he not refer to the mitotic figures? The general opinion is that these are present in large numbers, and do not necessarily indicate the activity. What effect on the individual is produced by injecting this leukocytic extract?

Dr. L. J. HIRSCHMANN, Detroit: I have observed some of Dr. Vaughan's work since he started; and, in answer to the inquiry of Dr. McConnell, I wish to state that in a number of cases of carcinoma of the bowel there has been a uniform regularity in the decrease of the polymorphonuclears and in the increase of the large mononuclears, when the leukocytic extract and residue have been injected. One could read the blood-picture without seeing the patient, and get a pretty good idea of the clinical condition of the patient on that day. It is merely suggestive, but it is uniform. After reading the blood-picture, it was not necessary to see the patient to tell whether he was improving or not.

Dr. G. W. McCASKEY, Ft. Wayne, Ind.: Two points of especial interest occurred to me during the reading of the paper. The first has reference to the biologic significance of the large mononuclear leukocytes. It appears perfectly clear from the leukocyte curves, that there is a very definite reaction on the part of these cells to the injection of the cancer residue. This fact would seem to me to point to the conclusion that these cells play a very definite rôle, and probably should not be regarded as transitional in any sense. It would seem to indicate that they form a class by themselves just as definite as the small mononuclears or polymorphonuclears.

The other point has reference to the bearing which this work might have on the embryonic theory of neoplasms. This theory seems to be predominant at the present time, and yet these investigations point, it seems to me, very

clearly, to some sort of specificity in malignant neoplasms. It would, of course, be quite possible that a specific toxin could act in conjunction with embryonic tendencies. I believe that this actually occurs in various pathologic conditions, as I have endeavored to point out in the pathology of pernicious anemia. We clinicians are greatly indebted to the pathologists for work such as this, so suggestive in its bearings, even if it does not as yet lead to definite and final conclusions.

Dr. ISAAC LEVIN, New York: How much necrosis and sloughing was there in these tumors that Dr. Vaughan was handling? Were the differences in the leukocytes going parallel with clearing up of this sloughing and necrosis? Does Dr. Vaughan mean by that the clinical improvement of the cases, or was he handling solid tumors without any central softening, necrosis or sloughing? This matter seems of great importance from our knowledge and understanding; whether there is a real specificity in that difference in the leukocytes of the blood, whether there is a specific relationship with cancer, or whether it is due to an amount of dead tissue present in the body.

Dr. J. W. VAUGHAN, Detroit: In reply to Dr. Levin, lately I have used an active ferment in a few cases with small tumors, when operation was absolutely refused, on condition that the patients would agree that, if benefit was not observed within two weeks, they would submit to operation. Among these cases were some breast tumors, which were not ulcerated at all. These tumors, when the ferment is injected, will swell up and soften, and afterward gradually contract down and harden again. At this time, it is proper to inject again; and they go through the same process again. An open tumor, one that is ulcerated, will do the same way after injection, and there is apt to be considerable bleeding for twenty-four hours afterward. This shows an increased vascularity following the injection of the specific ferment.

In reply to Dr. McCaskey's question with regard to the significance of this work in connection with the function of the cells, it is my belief that just as polymorphonuclear cells protect the body against outside infection, so the large mononuclear leukocytes protect the body against broken-down body tissue cells, and against autogenous growths or cancer cells. We are constantly having local injuries and broken-down tissue cells, and I believe that it is the function of the large mononuclear leukocytes to take care of these. I have invariably noticed that when a vaccine is used, the betterment in the patient ran in accordance with the increase in large mononuclear cells. For example, in one case which maintained 27 per cent. of large cells for weeks, which is a higher count than you usually are able to get, the tumor disappeared with rapidity. Transitional leukocytes I usually class as mononuclear leukocytes.

The question concerning the effect of the ferment on the human individual, was covered in my answer to the questions of Dr. Levin. If cancer vaccine is injected into a man who has cancer an increase in the large mononuclear leukocytes results. I have had cancer vaccine injected into myself; and I have found that I get a higher increase, when the vaccine is active, in such a case than I would in the case of a person who had cancer. In other words, there is a much higher increase in a normal individual. Concerning the prognosis in cancer through a study of the microscopic sections of any tumor, I believe that it is customary to estimate the malignancy by the presence or absence of mitotic figures. In our work, Dr. Sill, who is our pathologist, has run closely along with me in this work. We found that there apparently is no relationship between the percentage of large mononuclear leukocytes and the presence or absence of mitotic figures so far as we could estimate.

The Importance of Open-Air Exercise to the Gouty.—In the case of middle-aged men in affluent circumstances it is difficult to get them to moderate their appetites and regulate their diet, but they will often accede to the suggestion of taking more exercise. Motoring is good, but a long walk in the country every Sunday will do more to keep down their gouty proclivities. Exercise in a stuffy gymnasium is irksome and monotonous, and is soon abandoned.—William Murrell in *Clin. Jour.*