REVIEWS.

A PRIMER ON EXPLOSIVES FOR COAL-MINERS. C. E. MUNROE and C. HALL.

Bulletin 423 of the United States Geological Survey. Washington, 1909.

An octavo pamphlet of sixty-one pages, including contents, with twenty-one half-tone plates; it is intended to check the improper use of explosives in coal-mines. Combustion and explosion, and the nature of "low" and high explosives in general and as applied to coal-mining, are simply explained. Squibs, fuse, and detonators, the various modes of coal-mining with explosives, including the proper charges and placing of shots, loading and firing, the advantages of electric firing, and the application of explosives to driving shafts and entries, are dealt with. Throughout, the sources of danger, and precautions for avoiding them, are pointed out. A chapter on the transport and storage of explosives (by B. W. Dunn), a description of the tests for permissible explosives, carried out in experimental galleries containing gas or coal-dust mixed with air, and a list of the publications of the department on the subject, are appended.

O. E. MOTT.

ALLEN'S COMMERCIAL ORGANIC ANALYSIS. Fourth edition. Vol. IV. Edited by W. A. DAVIS, B.Sc., A.C.G.I., and SAMUEL S. SADTLER, S.B. 8vo. Pp. viii+466, with 7 illustrations. London: J. and A. Churchill, 1911. Price 21s. net.

In the volume under consideration a large proportion of the space is given to the essential oils and certain constituents of them, such as the terpenes, camphor and

other ketones, but this would appear to be fully justified in view of their commercial and technical importance, as well as the wide scientific interest associated with their more exact chemical investigation. It is not to be expected, however, that even within the space assigned to them, such a large group of natural products, so diverse in character and composition, could be described with the same degree of completeness as in a more specialised treatise on the subject, especially as methods of analysis and tests for the detection of adulterants and impurities must likewise be accorded the consideration which their importance demands.

In presenting a survey of the contents of the present volume, or more particularly of the sections devoted to the essential oils and allied subjects, it would seem desirable and of interest to record some general observations, and also to note some details of statement where inaccuracies appear to exist, without, however, attempting to enter upon a more extended discussion, for which the limitations of space would not permit.

The "Hydrocarbons of Essential Oils" are considered in a chapter of twenty-five pages (pp. 163-187), and these are followed by a chapter entitled "Ketones of Essential Oils," which comprises twenty-eight pages (pp. 189-216). A larger section (pp. 217-301), bearing the simple title "Volatile or Essential Oils" is devoted to a broader treatment of the subject, and includes their extraction, composition, general characters, and analysis, the general information under the latter heading being followed by a more detailed description of the methods employed for the detection or estimation of certain compounds. This comprises, for example, the detection of sulphur compounds, estimation of free acids, phenols, alcohols, esters, aldehydes, and ketones, together with the determination of such factors as the iodine value, methoxyl number, optical activity, and refractive index, and tests for some of the more common adulterants. A knowledge of these methods is naturally of very special importance to the analyst, inasmuch as the commercial valuation of a number of essential oils depends, not only on general tests for their identity and purity, but frequently on the quantitative determination of some particular constituent. A chapter entitled "Special Characters of Essential Oils" comprises within the limits of 128 pages (pp. 303-430) a description of the more important individual oils, and this is accompanied by a tabular representation of their principal characters, constituents, and solubilities, which occupies the last thirty pages of the work.

Notwithstanding the general excellence and the many valuable features of the present volume, there are also some apparent defects as well as obvious errors, and it is thought desirable that these should be noted for consideration in a subsequent revision, in order that a work which has achieved so high a reputation may continue to merit the confidence of those who consult it. In the chapter on the "Hydrocarbons of Essential Oils" there appears to be, for example, no reference to the so-called "olefinic terpenes," such as *myrcene*, which occurs in bay oil, sassafras-leaf oil, and the oil of hops, or the isomeric *ocimene*, from *Ocimum basilicum*. On the other hand, under the subject of "Pinene" (p. 180), this terpene is stated to form "a notable proportion of the oil from bay-leaves." It was shown, however, some years ago, that the true bay oil (not laurel oil) contains no pinene, as had previously been asserted, and its presence would therefore be direct evidence of adulteration

(*cf.* p. 315). The name Thoms is wrongly printed on p. 189 as *Thorns*, and on p. 210 as *Thomas*.

In the chapter on "Special Characters of Essential Oils" no consideration appears to have been given to such important articles of commerce as the oils of cinnamon and cassia, the oils of wintergreen (*Gaultheria*) and sweet birch, or the oil of bitter almonds, although the comparatively unimportant oils of hyssop and parsley are described. Under "California Bay Oil" (pp. 315, 316) the only information given regarding its constituents is the statement that "the oil contains a hydrocarbon and appears to contain cineol." It has thus here been overlooked (*cf.* p. 446) that this oil was very thoroughly investigated several years ago, and that its principal constituent, a ketone, $C_{10}H_{14}O$, designated "umbellulone," has been specially studied with respect to its constitution (*J. Chem. Soc.*, 1904, **85**, 629; 1906, **89**, 1104; 1907, **91**, 271; 1908, **93**, 252). Under "Nutmeg Oil" (p. 358) there occurs the statement that "the presence of *myristicol* has been noticed by Gladstone and by Wright," although in the investigation referred to in the paragraph immediately following it was shown that the fraction of nutmeg oil to which the designation "myristicol" had previously been applied consisted of several alcohols, all of which have now been definitely identified. A statement concerning the composition of mace oil (p. 359) is not quite correctly expressed, inasmuch as the crystalline compound obtained by Semmler does not pre-exist in the oil, but was formed by the action of alkalis on the liquid constituent now known as *myristicin*, and, being isomeric with the latter, has been designated *isomyristicin* (*J. Chem. Soc.*, 1907, **91**, 2039).

In the "Table of Essential Oils" (p. 434) there occurs, in connection with the oil of *Asarum Europæum*, the word "asarol," a name which was originally given to the chief constituent of the oil of *Asarum Canadense*; but as this compound has since been found to be identical with *d*-linalool, the latter designation should be adopted (*J. Chem. Soc.*, 1902, **81**, 63). On p. 445 the known constituents of oil of hedeoma (American pennyroyal) are given as hedeomol and pulegone, without consideration of the results of a more recent investigation, which are described on p. 377.

In a section immediately following the above, the name *Hedeoma* is incorrectly applied to the leaves of *Mentha pulegium*, which is known as "European pennyroyal," but the two genera are botanically quite distinct. The statement regarding the constituents of oil of hops (p. 445) should be revised in accordance with the more recent investigation of Chapman (*J. Chem. Soc.*, 1903, **83**, 505), and on p. 450 the statement that the known constituents of oil of nutmeg are "chiefly pinene and myristicin" would suggest some amplification (*cf.* p. 358).

The observations above noted, together with a considerable number of typographical errors, would indicate that in these relatively minor details a little more care might have been exercised. Nevertheless, the intrinsic merits of the work are such that it may be hoped it will receive the recognition accorded to its predecessors, and, in view of the large amount of valuable information it contains, that it will continue to maintain its place as a trustworthy guide in the manifold operations of chemical analysis and investigation.

F. B. POWER.

RESINS, INDIA-RUBBER, ESSENTIAL OILS.

Dr. M. B. Blackler is the author responsible for the section on resins, with which this volume opens. Those familiar with this subject know that, with a few exceptions, a particular resin cannot be identified by ordinary analytical processes alone, and consequently the analyst who is faced with a problem of this kind has to base his judgment on a great variety of factors, and must as a rule create his criteria from his knowledge of the properties and characters of resins as a whole. Dr. Blackler has kept this fundamental difficulty in mind, and has written and arranged his matter with a view to enabling his readers to obtain a clear idea of the differences which distinguish the chief groups of resins from each other. In the present nebulous condition of resin analysis this is undoubtedly the best method of procedure.

The errors of the article are for the most part inherited from previous editions of this work. On p. 6 it is implied, without any qualification, that abietic acid contains two carboxyl groups; whilst on p. 23 it is correctly pointed out that this view is erroneous, and that the acid is monobasic. The table on p. 6 includes "alban," one of the resins of guttapercha, as a "resene," although some years ago Van Romburgh and Cohen proved that guttapercha alban is an ester. The misleading information about "anime," including the statement that "it appears in general to be closely related to elemi and copal," is reprinted unchanged from the third edition.

Competent Indian authorities appear to be agreed that all shellac contains orpiment, and not only "some of the dirty yellow shellac of commerce," as the author asserts on p. 67. The third edition stated that French turpentine is called "Gallipot, or Barras," and Dr. Blackler not only repeats this erroneous statement on p. 76, but adds at p. 21 the equally inaccurate assertion that French colophony comes on the market under the name "galipot." Under "African balsam," p. 83, it is re-asserted that this is probably derived from a *Copaifera*, whereas it is now perfectly well known that it is obtained from *Daniella thurifera*. The old mistake is made of including the oleo-resin elemi under the "gum resins," and its botanical origin is wrongly given. The statement on p. 17 that a variety of acaroid resin is produced in the Bahamas is incorrect, though the author can no doubt quote good authority for his assertion.

The occasional application of the term "gum" to these products is unfortunate, since it is important, in view of the use of this term in commerce, for such widely different materials as true gums, resins, camphor, opium, and rubber that technical writers should restrict it to its proper application.

Practically all the important resins are dealt with, but descriptions of some of the Euphorbia resins and of gamboge ought to have been included, and as Chinese lacquer is now an article of European commerce, some reference should have been made to this interesting product.

T. A. HENRY.

INDIA-RUBBER.

The article on "India-rubber," which comprises india-rubber, rubber substitutes and guttapercha, is by Mr. E. W. Lewis, A.C.G.I. It covers fifty-five pages, of

which roughly forty-two pages are devoted to rubber and five pages to guttapercha. The balance of the article, under the heading Rubber Substitutes, deals with fatty (oil) substitutes, with "recovered" or "reclaimed" rubber, and with certain proprietary articles. In the article on india-rubber, Mr. Lewis deals first with latex and its analysis, and then gives a brief sketch of the work on and our knowledge of the composition and of the constitution of the rubber molecule. This is followed by some notes on the physical and chemical properties of crude rubber. The next section—namely, that on the analysis of crude rubber—is of a practical character, and should be decidedly useful to the general analyst who has no special experience of crude rubber. It might perhaps have been desirable to omit the reference to the Schneider process, or to have made the reference to it of a somewhat briefer character. Mr. Lewis would probably be the first to agree that the method has very little, if any, practical value. The remarks dealing with the vulcanisation of rubber, if somewhat brief, are sound, as one would expect from a chemist of Mr. Lewis's practical experience. Considerable space is devoted to the analysis of vulcanised rubber, and for those who desire further information there are ample references to the literature on the subject. On the whole, this section of the article has been dealt with admirably, but it is necessary for the reviewer to point out that recent work has conclusively shown that pyridine is not a suitable solvent for the separation of tar, pitch, and bituminous matter from many—possibly the majority—of rubber mixings (*cf.* Britland and Potts, *ANALYST*, 1910, **35**, 539). There are some useful notes on the analysis of ebonite and on the methods for determining the physical properties of rubber. Mr. Lewis deals very briefly with the latter, and also with the mechanical properties of rubber, and, in taking this course, he appears to have been well advised, inasmuch as the determination of the physical and mechanical properties of rubber require special apparatus and plant which are not likely to be present in the general analyst's laboratory. Without such special apparatus, attempts to determine the mechanical and physical properties of rubber are likely to do more harm than good, but Mr. Lewis has given references to the literature on this subject for those who desire to pursue it further. On the whole, Mr. Lewis is to be congratulated on handling a difficult and intricate subject in a sound and scholarly fashion.

P. SCHIDROWITZ.

THE MICROSCOPICAL EXAMINATION OF FOODS AND DRUGS. By HENRY GEORGE GREENISH, F.I.C., F.L.S., Professor of Pharmaceutics to the Pharmaceutical Society. London: J. and A. Churchill, 1910. Price 12s. 6d. net.

This is a second edition of Professor Greenish's book, which originally appeared in 1903. A section has been added on fibres, and some additions have been made to the previously existing sections of the work, which is one primarily designed as a working textbook for students. The well-known skill and care of the author as a microscopical expert have resulted in a set of drawings which, as far as they go, afford clear indications of the histological structure of many of the materials with which analysts as well as pharmacists are called upon to deal; but, as only typical articles are selected, the list—and consequently the general utility of the book—is limited. The first section is devoted to starches and the second to hairs and textile

fibres, the remaining sections dealing with spores and glands (lycopodium, lupulin, etc.), woods, stems, leaves, flowers, barks, seeds, fruits, rhizomes, and roots, about fifty different typical drugs being dealt with, as well as some of the more common food substances, the analysis of which calls for microscopical examination, some of the more common adulterants of powdered foods and drugs being included. The working instructions are good throughout, being the outcome of long personal experience in this important branch of the analyst's art.

BERNARD DYER.

IDENTIFICATION OF THE COMMERCIAL DYESTUFFS. Being Volume III. of a Method for the Identification of Pure Organic Compounds. By S. P. MULLIKEN. New York: J. Wiley and Sons. London: Chapman and Hall. 1910. Price 21s. net.

The introductory chapter refers to the difficulty and complexity of the problem involved in the identification of the several hundred definite chemical species of colouring matters liable to be met with in current industrial dyes. In Chapter II. the author gives directions for the general analytical procedure in the examination of an unknown dyestuff, dividing his tests into four groups: (*a*) Preliminary; (*b*) generic, divisional, and sectional; (*c*) co-ordination; and (*d*) special tests.

The reactions indicated under these headings are likely to prove of service to the analyst who is occasionally called upon to examine commercial brands of dyes, many of which are mixtures of several specific colouring matters.

The tests are arranged in a definite order and are referred to by numbers. The preliminary tests include the identification of indigoid dyes and sulphide colours by vat-dyeing experiments with sodium hydrosulphite and sulphide respectively. The generic tests contain descriptions of the colour-discharge effects produced by Rongalite (sodium formaldehydesulphoxylate) and the regeneration of the colour in certain cases by oxidation with potassium persulphate.

Among the co-ordination tests will be found the effect on dyed fabrics of such general reagents as concentrated sulphuric acid, aqueous sodium hydroxide, nitrous acid, and chromic acid (after-chroming).

The application of the foregoing reactions will generally enable the analyst to refer the colouring matter to its proper class, and special tests are now applied for the purpose of identification. Among these special tests may be mentioned—(i.) The precipitating action of sulphuric acid, sodium hydroxide, tannic acid, and the sulphates of calcium, chromium, and copper; (ii.) the dyeing of wool with different mordants; (iii.) the reduction of azo-dyes with sodium hydrosulphite, and the identification of the fission products; (iv.) the absorption spectra of dye-solutions.

Whenever possible, verification tests should be carried out by making comparative dyeings with the identified dye and standard specimens of the same colouring matter.

The greater portion of the volume is devoted to a tabulation of the commercially important dyestuffs arranged in accordance with their reactions towards the reagents employed in the generic or sectional tests, and indicating the tinctorial effects of the dyes on wool or cotton. The last chapter gives useful suggestions for the identification of dyestuffs in mixtures.

G. T. MORGAN.