of the eye, and in some sections a number of very thin membranous excrescences accompany it. This condition is apt to remind one of a persistent hyaloid artery. Yet, it is impossible to find any connection between this appendix and the central retinal blood-vessels in the papilla. It probably owes its origin to former hemorrhages and is similar in nature to what Manz termed proliferating retinitis.

No microbes could be found in the tissues of the eye.

From this description it is seen that the pathologic conditions in this eye, which undoubtedly caused a sym-

all the tissues—the severest inflammatory signs, either still actively present or shown by their results, are found in the retina, and especially in the posterior parts, where the chorioid is but little affected, quite contrary to the usual experience.

The character of the sympathizing inflammation in this case seems to have been preeminently a hemorrhagic and proliferating one. From this it seems that Fuchs laid out the limits of the range of what may be called a sympathizing inflammation somewhat too narrowly.

316 Metropolitan Bldg.

SCLEROSIS OF THE LIGAMENTUM PECTINATUM AND ITS RELATION TO GLAUCOMA *

F. H. VERHOEFF, A.M., M.D.

BOSTON

In the pathologic examination of glaucomatous eyes I have several times noted a condition that seemed of considerable significance in regard to the pathogenesis of glaucoma, namely, sclerosis of the ligamentum pectinatum with an open filtration angle. The condition, I find, has been previously described by a few other observers, all of whom have evidently regarded it as the cause of the advanced glaucoma with which it was associated. Brief mention is made of it by Schmidt-Rimpler,¹ and single cases have been recorded by Tartuferi,² Sarti³ and De Vries.⁴ The latter explained the sclerosis as due to toxins carried in the aqueous humor, an explanation which suggests that we have here a form of glaucoma dependent on the same cause as arteriosclerosis.

Thomas Henderson,⁵ in his theory of glaucoma, which, as I understand it, is a variation of the old retention theory, postulates a sclerosis of the ligamentum pectinatum. He maintains that with age the trabeculae of the ligament become thicker so that finally in some cases glaucoma ensues, owing to the cutting off of the access of aqueous humor to the canal of Schlemm and the consequent rise of intra-ocular venous pressure. At the same time, as a result of congestive edema, the iris swells, causing its root to become adherent to the ligamentum pectinatum. The actual existence of this advanced sclerosis, however, he has not demonstrated, no doubt because at the stage when glaucomatous eyes are usually obtained for histologic examination, adhesion of the iris root had been of such long standing that it was impossible to determine whether the sclerosis of the ligament was the cause or the result of the adhesion. It is true that in the few recorded cases in which glaucomatous eyes were examined at very early stages sclerosis of the ligament was not noted, but it is possible that they were not examined with special reference to this point.

It is obvious, therefore, that if it could be shown that cases of glaucoma occur in which there is primary sclerosis of the ligamentum pectinatum with the filtration angle partly or wholly open, Henderson's theory would receive strong support, since it would not be

Fig. 14.—Optic nerve and papilla. Infiltration and edema. Newly formed connective tissue covers retina and papilla and is detached, together with the lamina interna on the right, by hemorrhage, forming the large vesicle reaching beyond the macular region. From this newly formed tissue a tentlike process has grown into the vitreous chamber.

Fig. 15.—Optic nerve-head. Edema by lymphocyte infiltration, especially around the central blood-vessels. Retina thrown into folds by edema. Newly formed connective tissue covering the papilla.

pathetic inflammation of the fellow eye, while showing much which Fuchs considers as essentially characteristic of the sympathizing inflammation, differ in other points very materially from his description. While he states that the retina and optic nerve are usually but slightly affected, in this eye—aside from the considerable inflammation in the anterior parts near the wound which affects

* From the Massachusetts Charitable Eye and Ear Infirmary.

1. Schmidt-Rimpler: Graefe-Saemisch Handbuch der Gesamten Augenheilkunde, 1908, II, 104.

2. Tartuferi: Cited by Sarti, note 3.

3. Sarti: Bull. d. Soc. med.-chir. di Bologna, 1882, lxiv, 147; Centralbl. f. Augenheilk., 1895, p. 488.

4. De Vries: Nederl. Tijdschr. v. Geneesk., 1907, II, 263; Nagel's Jahrbuch, 1908, p. 680.

5. Henderson: Glaucoma, London, 1910.

expected that the assumed edema of the iris would always be sufficient to cause peripheral synechia. Even if Henderson's theory was not accepted in its entirety—and his argument does not seem altogether convincing—sclerosis of the ligament would no doubt be accepted as an essential cause of glaucoma. The evidence, however, that I have to bring forward, seems to point in an opposite direction, so that sclerosis of the ligamentum pectinatum as a cause of glaucoma yet remains to be demonstrated.

Following are brief summaries of the findings in ten cases in which there was sclerosis of the ligamentum pectinatum with an open filtration angle:

CASE 1 (2-1780).—Clinical History.—The patient, a man aged 38, was struck in the left eye while boxing twenty years ago. Cataract resulted, which was removed by discissions thirteen years ago. Nine years ago he was again struck in the left eye with a fist. After this the patient could see but little, and the eye was painful at times. Eighteen months ago paracentesis for acute glaucomatous attack was performed. The right eye was normal. The left eye was enucleated Dec. 18, 1909.

Pathologic Condition.—There were advanced secondary glaucoma following discission of the lens, with prolapse of the vitreous into the corneal incision, equatorial staphyloma, deep cupping of the optic disk and advanced atrophy of the retina,

Fig. 1.—Case 1. Primary glaucoma. Hyaline fibrous tissue extending over surface of ligamentum pectinatum and membrane of Descemet. The underlying trabeculae are still well seen.

iris and ciliary body. The pupil was free; the filtration angle was everywhere open. The ligamentum pectinatum was covered with fibrous membrane continuous with the anterior boundary layer of the iris. The spaces of Fontana and the canal of Schlemm were obliterated.

CASE 2 (2-891).—Clinical History.—The patient, a man aged 32, had an old injury of the left eye with traumatic cataract. Ten years ago the cataract was extracted, after which there was much pain. The tension was +3. The right eye was normal. The left eye was enucleated May 25, 1904.

Pathologic Condition.—There were advanced secondary glaucoma, old perforation of the cornea, traumatic cataract, ciliary staphyloma, deep cupping of the optic disk, advanced atrophy of the retina and atrophy of the ciliary body, marked at the site of the staphyloma, but slight elsewhere. The pupil was free; the iris was almost normal, except at the pupillary margin, where there was beginning ectropion. The filtration angle was everywhere open. The ligamentum pectinatum was covered with a smooth layer of hyaline tissue, which was lined by flattened pigmented cells. The trabeculae were visible, but the spaces of Fontana and the canal of Schlemm were obliterated (Fig. 3).

CASE 3 (2-2097).—Clinical History.—The patient, a man aged 73, twenty-seven years ago was struck in the right eye with a chip from a hammer. The eye soon afterward became blind and has since been very irritable at times. The tension was +1. The left eye was normal. The right eye was enucleated Nov. 10, 1911.

Pathologic Condition.—There were advanced secondary glaucoma, an old perforating wound of the cornea and lens, a small foreign body encysted on the posterior part of the ciliary

body, slight siderosis bulbi, localized chorioretinitis in the lower outer quadrant (due to movement of foreign body before it became encysted?), deep cupping of the optic disk and moderate atrophy of the retina. The pupil was free, partly excluded. The iris and ciliary body were almost free from atrophy, but showed slight recent iridocyclitis. The filtration angle was everywhere open. The ligamentum pectinatum was in places almost free from sclerosis. In these places the stroma cells of the ligament were greatly pigmented (pigment bleaches, hence not iron pigment) and the spaces of Fontana contained chronic inflammatory cells. Elsewhere, and to some extent also here, the ligament was covered with a thin hyaline layer lined by pigmented cells continuous around the angle with the anterior boundary layer of the iris. The underlying trabeculae and spaces of Fontana were almost normal.

CASE 4 (2-557).—Clinical History.—The patient, a man aged 48, was struck in the right eye with a piece of steel twenty-two years ago; the left eye was normal. The right eye was enucleated May 1, 1901.

Fig. 2.—Case 4. Primary glaucoma. Dense hyaline fibrous tissue on surface of ligamentum pectinatum. The underlying trabeculae are greatly compressed so that the new tissue might readily be mistaken for the ligament itself.

Fig. 3.—Case 2. Secondary glaucoma. Ligamentum pectinatum covered with a smooth layer of hyaline tissue lined by flattened pigmented cells. The almost complete absence of iris atrophy indicates that the separation of the peripheral synechia occurred early. The atrophy of the ciliary body here is due to a ciliary staphyloma.

Pathologic Condition.—There were advanced secondary glaucoma, traumatic cataract, equatorial staphyloma, foreign body in the vitreous chamber, slight siderosis bulbi, deep cupping of the optic disk and advanced atrophy of the retina, iris and ciliary body. The pupil was free. The filtration angle was mostly open, being closed only along a small part of the circumference. The ligamentum pectinatum was covered with a thin dense hyaline layer presenting a smooth surface lined by endothelium. The spaces of Fontana and the canal of Schlemm were obliterated.

CASE 5 (2-2057).—Clinical History.—The patient, a man aged 53, received a perforating wound of the corneal limbus of the left eye thirty-four years ago. The foreign body was retained in the globe. Four years ago the patient was told that he had a cataract in the left eye. The eye was painful and moderately injected; the cornea was clear, the pupil free, the lens cataractous, and vision was nil. The right eye was normal. The left eye was enucleated Oct. 4, 1907.

Pathologic Condition.—There were secondary glaucoma, an old perforating wound of the cornea limbus, remains of the foreign body encysted in the sclera near the optic disk, slight siderosis bulbi, moderate cupping of the optic disk and moderate atrophy of the retina. The filtration angle was open on one side and closed on the other. On the closed side the ciliary body and the root of the iris were greatly atrophied. Elsewhere the ciliary body was almost free from atrophy. The ligamentum pectinatum in places was covered with a delicate hyaline membrane. The spaces of Fontana were not obliterated. The iris was practically normal except at the point where the root adhered and at the pupillary margin (probably due to the fact that the peripheral synechia became separated on one side before there was atrophy of the iris root, so that the circulation of the iris was not seriously interfered with even on the adherent side).

CASE 6 (2-998).—Clinical History.—The patient was a man aged 41. Eleven years ago the left eye was inflamed and painful (iritis?). The present attack of pain and redness began five days ago. Vision was *nil*. There were posterior synechia

Fig. 4.—Case 9. Primary glaucoma. Ligamentum pectinatum covered with a new-formed elastic membrane continuous with the membrane of Descemet. Elastic-tissue stain.

Fig. 5.—Case 7. Primary glaucoma. Iris in act of pulling away from ligamentum pectinatum.

and hyphemia. The tension was +1. Right eye, vision 20/50. The left eye was enucleated Aug. 1, 1904.

Pathologic Condition.—Advanced secondary glaucoma was present and old iritis with exsclusio pupillae; no cyclitis, old or recent. There were deep cupping of the optic disk, advanced atrophy and complete separation of the retina (subretinal coagulum), recent hemorrhagic retinitis and obstruction of both central retinal vessels from endovasculitis. The iris was highly atrophied; the ciliary body only slightly so. The filtration angle was closed on one side only. The ligamentum pectinatum was not greatly altered, even where the iris was adherent, the trabeculae and spaces of Fontana everywhere being well seen. On the open side, along the surface, the spaces of Fontana were closed off by proliferation of the stroma cells. On one side, at the site of the adhesion, there was evidently a state of tension, broken strands of iris tissue bridging the angle. The strongest adhesion here was that between the anterior part of the ligament and the iris. It is evident that ultimately the picture seen in Case 8 (Fig. 7)

would have resulted. The internal spaces of Fontana were dilated, due, probably, to traction on the trabeculae. The dilated spaces of Fontana and the irregular spaces formed by the breaking adhesion were frequently found filled with blood, which was continuous with an interstitial hemorrhage in the root of the iris. The canal of Schlemm was almost obliterated.

CASE 7 (2-1646).—Clinical History.—Patient was a woman aged 61. About six years ago a cataract began in the left eye. For several years past the eye had been painful and for the past eighteen months it had been absolutely blind. The tension was +1. The right eye was normal (?). The left eye was enucleated Jan. 23, 1909.

Pathologic Condition.—There were advanced primary glaucoma, deep cupping and cavernous atrophy of the optic nerve, advanced atrophy of the retina, moderate atrophy of the iris and almost no atrophy of the ciliary body and active recent iritis with posterior synechia and pupillary membrane. The filtration angle was open almost everywhere. The ligamentum pectinatum showed three grades of sclerosis: a, vascularized connective tissue over its surface without complete obliteration of underlying spaces of Fontana; b, thick layer of dense fibrous tissue on its surface, with almost complete obliteration

Fig. 6.—Case 10. Hemorrhagic glaucoma. Iris in act of pulling away from ligamentum pectinatum.

Fig. 7.—Case 8. Primary glaucoma. Iris root pulling away from ligamentum pectinatum and held only by a strand of iris tissue containing a blood-vessel.

of the spaces of Fontana (Fig. 2), and c, condition intermediate between a and b (Fig. 1). Where the angle was closed the adhesion was evidently old, but was in the act of pulling away from the ligament. Here the spaces of Fontana have disappeared, and irregular strands of iris tissue, clearly under tension, some of them containing capillaries, stretched between the surface of the ligament and the iris (Fig. 5).

CASE 8 (2-1900).—Clinical History.—The patient, a man aged 67, has had no vision in the left eye for fifteen years. Three years ago he began to have attacks of redness and pain. Six weeks ago he began to have pain in the right eye and to see colored rings around lights. Vision of the right eye was 20/30; field considerably contracted. The left eye was enucleated April 16, 1910.

Pathologic Condition.—There were advanced primary glaucoma, deep cupping of the optic disk and advanced atrophy of the retina and iris. The ciliary body was moderately atrophic. The pupil was free. The filtration angle was everywhere free, except as noted below. The ligamentum pectinatum in places

was covered with a layer of vascularized hyaline tissue, sometimes containing elastic tissue and presenting a smooth surface lined by endothelium. In other places the spaces of Fontana were partly opened and the surface of the ligament was ragged, showing portions of iris tissue adherent to it. Here, the opposite surface of the iris was also ragged. In one place the angle was bridged by a strand of iris tissue containing a blood-vessel (Fig. 7). The spaces of Fontana were for the most part obliterated, and the trabeculae were not easily recognized.

CASE 9 (2-1792)—Clinical History.—The patient, a man aged 66, twenty-eight years ago was struck in the left eye with a stone, but the eye gave no trouble until two years ago, when it became inflamed and painful; it had been subject to recurring attacks since then. The cornea showed an old central scar with calcareous deposits—atheromatous ulcer. After the calcareous deposits were scraped out the patient neglected treatment and ulcus serpens ensued. The right eye was not glaucomatous; vision was 20/30. The left eye was enucleated Nov. 28, 1909.

Pathologic Condition.—There were advanced primary glaucoma, deep cupping of the optic disk, advanced atrophy of the retina, moderate atrophy of the iris and ciliary body, active ulcus serpens without perforation and purulent iritis. The filtration angle was for the most part open; in places partly closed. The ligamentum pectinatum was covered with a new-formed elastic membrane continuous with the membrane of Descemet (Fig. 4). The spaces of Fontana were not obliterated and the trabeculae were well seen. Where the angle was partly open, the adhesion covered about two-thirds of the ligament. Here it was sometimes seen that the adhesion was in the act of separating at the apex of the angle, leaving only the anterior part adherent.

CASE 10 (2-1935).—Clinical History.—The patient, a man aged 57, had hemorrhages into the vitreous of the right eye five years ago. Three years ago the vitreous was clearer, vision equaled hand movements. One year ago he had an acute attack of glaucoma, which lasted about one month. There was recurrence of tension and pain. The left eye showed sclerosis of the retinal vessels, no glaucoma and normal vision. The right eye was enucleated Dec. 10, 1910.

Pathologic Condition.—There was advanced hemorrhagic glaucoma. The retina was atrophic and covered with hemorrhages. There was deep cupping of the optic disk. Complete obstruction of the central retinal vein. The filtration angle was open around one-half of its circumference. On the adherent side, the iris and ciliary body were highly atrophic and there was ectropion uveae; on the open side there was distinctly less atrophy and no ectropion. The ligamentum pectinatum on the open side was covered wholly or in part with a layer of vascularized connective tissue, having a smooth surface and continuous around the angle with the iris. At the place of transition between the closed and open angle, the adhesion was in the act of pulling away from the ligament, the tissue intervening between the iris and ligament evidently being under tension (Fig. 6). In places the apex of the angle was freed, leaving the anterior portion of the ligament adherent, but in other places the anterior portion was being freed first.

In each of the above ten cases the glaucoma was in an advanced stage, with deep cupping of the optic disk and marked atrophy of the retina. Three cases were of primary, one of hemorrhagic and six of secondary glaucoma. Of the latter, one was due to old iritis of unknown etiology, with exclusion of the pupil, one followed dissections of a traumatic cataract, and four followed perforating wounds of the globe in three of which there were retained foreign bodies and siderosis bulbi. In only the first three cases was the filtration angle open all around. In the others there were peripheral synechiae of greater or less extent which might well have been overlooked if the eyes had been sectioned without this point in view.

These cases seem to me to offer convincing evidence that the sclerosis of the ligamentum pectinatum that has

thus far actually been observed in glaucoma, is not the cause of the latter, but the result of preceding peripheral anterior synechiae. The most important evidence, perhaps, is furnished by the last five cases in which the iris is found in the act of pulling away from the ligament (Figs. 5, 6 and 7), thus showing plainly that peripheral synechiae had once been complete. This is strongly supported by the fact that the changes in the ligament in all the cases involved chiefly its surface. Moreover, the fact that the condition in question occurred in secondary, as well as in primary, glaucoma, is almost alone sufficient to prove that the sclerosis was not the cause of the increased intra-ocular pressure. In three of the cases, it is true, there was slight siderosis, so that it is conceivable that here the sclerosis was due to iron pigment deposited in the ligament which later disappeared, but this explanation is improbable from the fact that it does not apply to the other seven cases in which the changes in the ligament were similar.

The question as to the cause of the preceding peripheral synechiae has no direct bearing; necessarily, on the question as to the cause of sclerosis of the ligament, and need not be discussed here. In the case of primary glaucoma, it no doubt involves the fundamental question relating to the cause of the increased intra-ocular pressure, which thus far has never been satisfactorily answered. In the case of secondary glaucoma, Fuchs⁶ has shown that peripheral synechia frequently is due to an inflammatory condition of the iris and ligament which leads to their contact.

In all of the above cases the changes in the ligament were essentially similar, differing chiefly in degree only, and consisted in the occurrence of new tissue on its surface with more or less obliteration of the spaces of Fontana. The new tissue consisted of vascularized connective tissue, hyaline or elastic tissue, or in one case partly of tissue closely similar to the iris stroma. These changes were perhaps produced in this way. During the persistence of the adhesion of the iris root to the ligament, the openings of the spaces of Fontana into the anterior chamber became permanently obliterated by cell proliferation. If the iris then pulled away early, a comparatively smooth surface remained; if late, more or less tissue from the iris was left. Over this surface the corneal endothelium grew, and beneath it was gradually formed a hyaline layer, or even an elastic membrane (Fig. 4), identical with and continuous with the membrane of Descemet. The obliteration of the spaces of Fontana deeper down was probably due chiefly to compression.⁷

Granting that sclerosis of the ligamentum pectinatum in the above cases was due to peripheral anterior synechiae which became separated, how was this separation brought about? In three of the traumatic cases the iris was almost free from atrophy, so that no doubt the separation occurred early, as was indicated also by the fact that the changes in the ligament were comparatively slight. Here, evidently the traction exerted by the sphincter pupillae was alone sufficient to accomplish the separation. Fuchs⁶ suggests that peripheral synechiae

6. Fuchs: Arch. f. Ophth. (Graefe's), 1909, lxi, 254.

7. It may be well to emphasize here the fact that this paper is concerned only with the type of sclerosis of the ligament described in the literature and illustrated by the above cases. I have sections of an eye (2-1042) in which there is an entirely different type of sclerosis. Here the internal portion of the ligament is relatively normal; while in places there is marked cell proliferation immediately about Schlemm's canal. In this case there is no cupping of the disk and the filtration angle is everywhere open. On the other hand, there is albuminuric retinitis and endovasculitis of all the intra-ocular vessels, so that the sclerosis involving the wall of Schlemm's canal is no doubt simply a part of a general angio-sclerosis.

associated with traumatic iritis may be freed in this way, without, however, discussing the possible effect on the ligament thereby resulting. In the cases in which there was marked atrophy of the iris, it is evident that the freeing of the adhesion must have taken place relatively late, and that it could not have been due to traction of the sphincter alone. The explanation of these cases, no doubt, lies in the cicatricial contraction of the atrophied iris. This contraction, being resisted all around by the ciliary body, must have produced a state of tension in the iris which caused traction on the peripheral synechiae in a direction practically at right angle to the plane of the iris. The pull was therefore not toward the pupil, as at first might be supposed, but backward, or even slightly toward the periphery. The reason that such separations of peripheral synechiae do not occur more often is no doubt due to the fact that the adhesion is apt to be especially firm in just those cases in which the cicatrization of the iris is greatest.

233 Charles Street.

ABSTRACT OF DISCUSSION

DR. M. FEINGOLD, New Orleans: What constitute the essential features of glaucoma? Is it the increased tension alone that constitutes glaucoma? Or are other symptoms of glaucoma correlated and equally essentials of the disease, such as the affection of the optic nerve? Most authors agree that glaucoma is essentially and solely increased tension and that all symptoms found in glaucoma are either symptoms leading to glaucoma or symptoms that are the consequence of increased tension. In the former group would belong such symptoms as shallow anterior chamber, small cornea, cyclitis, occlusion and seclusion of the pupil, adherent leukoma, etc. To the latter group would belong the defective epithelium and edema as well as the loss of sensitiveness of the cornea, atrophy of the iris, oval pupil, and above all the glaucomatous excavation. If it be true that glaucoma be purely and simply nothing else but increased tension we must ask, how is increased tension brought about?

Problems seen in daily practice, convince one that no one of the theories of glaucoma can satisfactorily account for all cases of glaucoma, and that for the present we are compelled to call in several of the theories to explain a single case of glaucoma. Sclerosis of the pectinate ligament, as proclaimed by Henderson to be the sole cause for glaucoma, may possibly in the future be demonstrated as the primary factor in causing glaucoma. Dr. Verhoeff's pictures prove that, at least in his case, the sclerosis of the ligament existed with peripheral anterior synechia, these anterior synechiae having been separated during the course of the disease. Such pictures are rare because in most instances the iris is so much in contact with the cornea that the coexistent sclerosis of the ligament cannot alone be blamed for the glaucoma. Ranvier remarked years ago that the pectinate ligament becomes sclerotic with age, but no undoubted case of pure sclerosis of this ligament has been found as yet as the sole cause for glaucoma. Still such possibilities may be admitted. Could not syphilis, that plays so important a part in the morbidity of the colored race, could not syphilis with its consequent vascular changes, be held responsible for the simple glaucoma that we see in the colored race almost to the exclusion of any other form of primary glaucoma?

DR. WALTER BAER WEIDLER, New York: Will Dr. Verhoeff explain just what he means by a sclerosis of the ligamentum pectinatum; whether there was any infiltration present or not; whether there are any cases of hypopyon or whether there was any pus present in any of the specimens shown? I thought I saw deposits in the anterior chamber.

DR. F. H. VERHOEFF, Boston: By sclerosis of the ligament I mean, in a general way, thickening of the framework of the ligament. In these cases there was also new formation of

connective tissue on the surface. The leukocytes in the filtration angle shown in some of the illustrations were due in one case to an active ulcerus serpens, and in another to recent iritis. These, of course, were of too recent occurrence to have had anything to do with the sclerosis of the ligament.

THE KRÖNLEIN OPERATION AS AN EXPLORATORY PROCEDURE IN AFFECTIONS OF THE ORBIT

ARNOLD KNAPP, M.D.
NEW YORK

The purpose of this paper is to draw attention to the facility of the osteoplastic resection of the temporal orbital wall, commonly called the Krönlein operation, as an operative procedure, and its adaptability as a diagnostic step to obtain exact information of conditions present in the depth of the orbit such as morbid processes in the orbital walls, in the optic nerve and in the soft parts.

I take it for granted that this operation is accepted to be the best method of removing circumscribed tumors in the depth of the orbit with the preservation of the eyeball and, in many instances, with the preservation of sight and unimpeded ocular motility.

The operation, furthermore, permits the best determination of the extent of many orbital tumors, often their operability, and the securing of a proper specimen for microscopic examination as a preliminary to an orbital exenteration if found necessary, or, in the case of rapid malignancy, the abandonment of further operations. The question of the advisability of the Krönlein operation in inflammatory conditions of the orbit seems to me to be still open, for it is now known that most inflammatory conditions in the orbit are secondary to disease of the nasal accessory sinuses, in which case operation at the internal orbital wall is the best.

The method of performing the Krönlein operation is sufficiently well known not to require a detailed description, yet the following points are of practical importance in my experience. The cutaneous incision should be slightly to the outer side of the periosteal incision and of the bony wound, so that the line of cutaneous cicatrix shall not be directly over and adherent to the bony margin of the orbit. The incision of the periosteum along the bony margin should be carefully made with the point of the knife directed against the bone, so that a perforation of the orbital periosteum with premature prolapse of the orbital contents can be avoided. The separation of the orbital periosteum can best be done with a sharp flat periosteal elevator. The bone-flap is formed by a medium-sized osteotome and hammer from the orbital surface, and forcibly displaced outward.

This displacement is aided by two incisions of the deep muscular tissues and periosteum, each radiating from the anterior extremities of the bony wedge. If the bone in the deeper part of the wedge has been fractured, more bone can be removed with the rongeur forceps in order to give the greatest possible access to the orbit. The bone externally at the orbital margin and malar process must not be removed, as the wedging of the bony flap in this position is necessary to keep it in place and prevent deformity when the bone flap is replaced at the end of the operation.

The horizontal incision of the orbital periosteum is held apart with small retractors and avoiding the