This grouping or clustering together of this class of nebulae is very remarkable and suggestive.

There are quite a number of such clusters the most remarkable of these being one composed of very small

Yerkes Observatory, Williams Bay, Wisc., 1897 June 1.

Note 1906 Aug. 30. These were put in an envelope and addressed to the A. N. nine years ago, but for some unknown reason were never put in the mails.

E. E. Barnard.

Group of small nebulae.

On 1890 Aug. 20 I swept up with the 12 inch of the Lick Observatory a group of small nebulae in approximately for the date

$$\alpha = 15^{\text{h}} 34^{\text{m}} \quad \delta = +44^{\circ} 0'$$

from a pointing.

The enclosed sketch (field 43' di.) was made. Following are descriptions

1	Pretty bright. Small round. Much brighter in middle
2	Faintish. R g b M
3	" "
4	Faint "
5	Not faint "

The star in the following edge of field was estimated 7.5 mag. and is supposed to be BD. +44°2493 6^m.8 whose place for 1855.0 is 15^h33^m29^s.4 +44°5'0". A note says "another star 7^m.5 p. this 40' and is 25' south (BD. +43°2560).

The smaller nebula close following 1 is marked with a question as to whether it was a faint star.

Yerkes Observatory, Williams Bay, Wisc., 1906 Sept. 1.

E. E. Barnard.

On a group of small nebulae

observed near Brooks comet (1889 V) Oct. 28, 1889.

By *E. E. Barnard.*

On 1889 Oct. 28 while observing Brooks comet (1889 V) with the 36 inch of the Lick Observatory I found a group of small nebulae some 15' following the comet. Several of these were measured with the great telescope.

They were referred to a $9\frac{1}{2}$ mag. star.

	Pos Angle	Distance
<i>a</i>	226°7 (4)	145.1 (3)
<i>c</i>	281.2 (2)	181.2 (2)
<i>b</i>	306.1 (5)	144.2 (4)

α is $5'' \pm$ n. p. a 10^m star.

Following are descriptions of these nebulae — the letters referring to the enclosed sketch (size of field $6' \pm$):

<i>a</i>	R.	not F.	m b M
<i>b</i>	p B.	R.	m b M
<i>c</i>	v S.	c F.	b M
<i>d</i>	p S.	p F.	g b M
<i>K</i>	p S.	F.	g b M

There are probably a number more nebulae here.

Comparison star is BD. $-3^{\circ}57'04$ (9^m5) whose position is

$$1855.0 \quad 23^h40^m5^s2 \quad -3^{\circ} 7'6.$$

South following it is BD. $-3^{\circ}57'06$ (9^m2) on the above date, with the 12 inch refractor, I determined the $\Delta\alpha$ and $\Delta\delta$ of these two stars

$$\Delta\alpha = 0^m21^s06 \text{ (12 transits)} \quad \Delta\delta = 1'27''.6 \text{ (2 measures)}$$

Yerkes Observatory, 1906 Sept. 1.

E. E. Barnard.

The nebula NGC. 6302.

$$1860.0 \quad 17^h4^m17^s \quad -36^{\circ}56'.$$

By *E. E. Barnard.*

This nebula was discovered by me in 1880 at Nashville, Tennessee, while hunting comets with my 5 inch refractor. Subsequently observing it with his 16 inch at Rochester N. Y., Swift found it to be triple.

In 1892 I examined and measured the nebula with the 36 inch of the Lick Observatory and found it to be a very remarkable object. With the great telescope it was seen to consist of three rather bright small nebulae. Two of the nebulae had faint streamers running north preceding, while the third, the following component, had two nebulous

arches springing from it and extending some distance following. The entire nebula, especially the following part, looked like a ghostly bug of some kind; the third mass being the head and antennae. From its singular appearance, I have called it the "Bug Nebula". The enclosed drawing was made from three or four nights study of it with the 36 inch, and very fairly represents its remarkable appearance.

Careful measures were made of the nebula, but the drawing is not from these measures. I also send a rough diagram which indicates the portions measured.

• K

North

In the following measures the letters refer to those on the diagram.

<i>CA.</i>			
	Pos. Angle	Distance	
1892 July 29	$261^{\circ}5$ (2)	$38''.2$	(2)
<i>CK.</i>			
1892 July 29	$294^{\circ}1$ (3)	$152''.66$	(3)
Aug. 5	294.0 (4)	151.85	(5)
" 12	294.0 (3)	152.78	(3)
	$294^{\circ}0$	$152''.43$	

K is a $13\frac{1}{2}$ magnitude star n. p.

<i>CB.</i>	
1892 July 29	Dist. $11''.55$ (2)

<i>CP.</i>	
1892 July 29	Dist. $28''.31$ (2)

Yerkes Observatory, 1906 Aug. 30.

The figures in parenthesis represent the number of settings.

The E and W diameter of the brightest part of *C* = $8''.7$, *C* is somewhat elliptical.

Estimations were made of the relative brightness of the different parts of the nebula.

C is 5 times as bright as *B*

C is 4 times as bright as *A*.

Another estimate made *B* very slightly brighter than *A*.

The brightest part, *C*, was estimated to be of 12 mag. The other two portions were about 14 mag.

Too much reliance must not be placed on the exactness of the position of this nebula as given in A. N. 2588 Bd. 108 p. 369.

E. E. Barnard.