329

Author(s): John H. Lawlor

Source: *The Mathematical Gazette*, Vol. 5, No. 88 (Oct., 1910), pp. 338-339

Published by: The Mathematical Association

Stable URL: <http://www.jstor.org/stable/3603426>

Accessed: 19-06-2016 20:12 UTC

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at
<http://about.jstor.org/terms>

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.

The Mathematical Association is collaborating with JSTOR to digitize, preserve and extend access to *The Mathematical Gazette*

parallel to AB . Then $AM = a\theta = BL$. By projection the coordinates of B are seen to be :

$$(i) \ x = OP - PL - BL \cos \frac{\theta}{2} = 2PL - PL - BL \cos \frac{\theta}{2} = 2a \sin \frac{\theta}{2} - a\theta \cos \frac{\theta}{2}.$$

$$(ii) \ y = a\theta \sin \frac{\theta}{2}.$$

(iii) The normal to the locus of B is evidently

$$BP = \sqrt{BL^2 + PL^2 + 2BL \cdot PL \cos \frac{\theta}{2}} = a \sqrt{\theta^2 + 4 \sin^2 \frac{\theta}{2} + 2\theta \sin \theta}.$$

(iv) The radius of curvature at B of the locus is by the formula,

$$\frac{\alpha \left(4 \sin^2 \frac{\theta}{2} + \theta^2 + 2\theta \sin \theta \right)^{\frac{3}{2}}}{4 \sin^2 \frac{\theta}{2} + \theta^2} = \frac{BP^3}{BL^2 + PL^2} = \frac{2BP^3}{OB^2 + BP^2} = \frac{2n^3}{r^2 + n^2},$$

if r denote the rad. vector OB and n the normal BP .

$$(2) \text{ Equation to } BL \text{ is } x \sin \frac{\theta}{2} + y \cos \frac{\theta}{2} = 2a \sin^2 \frac{\theta}{2}.$$

Hence the locus of the foot of the perpendicular from O on BL is $r = 2a \cos^2 \theta = a(1 + \cos 2\theta)$, whose pedal equation is $r^3 + 3rp^2 - 8ap^2 = 0$. The pedal equation to the envelope of BL is thus $r^2 + 3p^2 - 8ap = 0$.

M. T. NARAYANA IYENGAR.

328. [A. 1. a.] *Note on the factorization of the trinomial $x^2 + px \pm q$.*

Beginners learning algebraical factors are easily misled by such trinomials as $x^2 + 37x - 210$; the more careless will give factors involving 30 and 7, while others, noticing that these numbers "do not work," will be confident that there is a misprint in the signs. It affords useful practice to give groups of four such expressions for factorization, e.g.

$$x^2 + 37x - 210, \quad x^2 + 37x + 210, \quad x^2 - 37x + 210, \quad x^2 - 37x - 210,$$

$$\text{or} \quad x^2 - 29x + 210, \quad x^2 + 29x - 210, \quad x^2 + 29x + 210, \quad x^2 - 29x - 210,$$

and so on.

To construct such examples it is necessary to find a formula giving values of p and q for which $x^2 + px + q$ and $x^2 + px - q$ will both factorize. The required formula is

$$p = k(a^2 + b^2) \quad \text{and} \quad q = k^2 \cdot a \cdot b(a^2 - b^2),$$

where a and b are integers, $a > b$, and k is any odd integer not containing a squared factor.

$$\text{Then } x^2 + k(a^2 + b^2)x + k^2 \cdot a \cdot b(a^2 - b^2) = \{x + k \cdot a(a - b)\}\{x + k \cdot b(a + b)\},$$

$$\text{and } x^2 + k(a^2 + b^2)x - k^2 \cdot a \cdot b(a^2 - b^2) = \{x - k \cdot b(a - b)\}\{x + k \cdot a(a + b)\}.$$

C. C. WILES.

329. [A. 2. a.] The scheme used by many elementary text-books in assisting the student to memorize the ratios $x : y : z$ when determined from

$$a_1x + b_1y + c_1z = 0,$$

$$a_2x + b_2y + c_2z = 0,$$

usually takes some form like this :

$$\begin{array}{ccc} a_1 & \nearrow & b_1 \\ & \searrow & \nearrow c_1 \\ a_2 & \nearrow & b_2 \\ & \searrow & \nearrow c_2 \end{array}$$

Might I suggest the following scheme which impresses the cyclic order on the youthful mind?

The explanation is obvious, and so is the rule of signs to be adopted in following the dotted line. The idea is capable of extension in other directions that will be obvious.

JOHN H. LAWLOR.

330. [K. d. 5.] *To inscribe a triangle in a given triangle so that the sum of the squares of its sides shall be a minimum.*

Let ABC be the given triangle.

Consider a triangle PQR inscribed in ABC , and let the circum-circles of the triangles AQR , BRP , and CPQ intersect in the common point O .

Then we know that the minimum triangle similar to PQR inscribed in ABC is XYZ , where OX , OY , and OZ are perpendicular respectively to BC , CA , and AB .

Consequently the sum of the squares of the sides of the triangle PQR cannot be a minimum unless the perpendiculars at P , Q , and R to BC , CA , and AB respectively are concurrent.

Let XYZ be the triangle for which $YZ^2 + ZX^2 + XY^2$ is a minimum.

This expression is equal to

$$OA^2 \sin^2 A + OB^2 \sin^2 B + OC^2 \sin^2 C.$$