quence of ossification of the vocal cords and esophagus or from tuberculosis secondary to ankylosis of the ribs. The disease may persist 10 to 15 years. The author has no inkling of a suggestion of vegetative nerve mechanisms.

del Valle, R. PROGRESSIVE MUSCULAR ATROPHY. [Riv. d. Med. y. Cir. Pract., Oct., 1919. J. A. M. A.]

Del Valle reports pronounced improvement in the case of progressive muscular atrophy in a man of 46 under electricity and hot sand baths twice a day. The best results were noted when the sand bath just preceded the faradization. The impotence of the hands from the muscular atrophy retrogressed under this mixed treatment in five months to such a degree that further treatment was not required. He was a working-man, and the first symptoms had been noted three years before. Repeated chilling of the hands was the only causal factor that could be discovered, and this suggested the advisability of the hot sand baths. The advantages of the sand bath are the unusually high temperature that can be used, the fact that perspiration can proceed unhindered, and that there is no maceration of the skin. No benefit was apparent from a course of strychnin, and it was abandoned.

Guillain, G., and Barré, J. A. PERIODIC PARALYSIS. [Ann. de Méd., Dec., 1919.]

A soldier, 36 years of age, had a paralysis of the arms which recurred about 2 p.m. with great regularity. It was not accompanied by pain but there was great weakness. No paresthesiæ. After 4-5 hours the weakness passes and the muscles were normal during the day, for occasional fatigue on writing. He had had this for sixteen years in mild forms, but the actual paralysis did not develop until he was in the trenches. The attacks vary in intensity. During the severer ones the normal electric reactions disappear. The cranial nerve innervations never showed paralysis. The tendon reflexes disappeared during the severer attacks, but the superficial reflexes showed no changes. No sensory disturbances. The blood Wassermann was positive on two occasions, but the spinal fluid shows no sign of neurosyphilis. The authors assume an intermittent toxic accumulation hypothesis.

Campbell, A. W. THOMSEN'S DISEASE. [Med. Jl. Australia, Dec. 13, 1919.]

A man, 27 years of age, for ten or eleven years had had a good muscular development to all appearances, but whose real power was slight. Every movement was hampered by the characteristic rigidity and tendency to tonic contraction, of Thomsen's disease and whose muscles were unduly firm in relaxation and show an alteration of electrical excitability, suggesting the myotonic reaction. The familial history was absent.