FIG. 1. Acquired atresia of the vagina showing pinhole opening where the septum had been incised for the relief of symptoms at the time of puberty.

FIG. 2. Operation for acquired atresia of the vagina. The transverse septum has been dissected out, and the upper portion of the mucous membrane of the vagina united to the lower by interrupted catgut sutures.

FIG. 3. Operation for congenital absence of the vagina. The four flaps taken from the labia minora and the thighs are approximated over a glass form. The wounds where the flaps were removed are represented as sewed up with interrupted sutures. The pouch made by the four flaps is turned into the opening previously made by separating the bladder from the rectum.

FIG. 4. Operation for congenital absence of vagina. The glass form has been removed and the seams of the pouch completely sewed. The untied sutures seen issuing from the pouch were previously passed through the vault of the artificial opening made between the bladder and rectum. The pouch is inverted into the artificial opening and the stitches tied so as to prevent any prolapse of the pouch before it heals to the raw surface of the opening.

Original Articles.

OPERATIVE TREATMENT OF ATRESIA OF THE VAGINA.*

BY W. P. GRAVES, M.D., BOSTON.

ATRESIA of the vagina may be either acquired or congenital. The acquired form is by far the more common, being the result of an extensive adhesive vaginitis. Congenital atresia is due to defective embryological development and is always accompanied by other defects involving the genital organs.

ACQUIRED ATRESIA OF THE VAGINA.

The majority of cases of this condition represent the end result of a long-continued plastic vaginitis usually gonorrheal in origin and acquired before the age of puberty. The vaginal mucous membrane of the adult is peculiarly resistant to the gonococcus, and is rarely infected excepting secondarily, and then not frequently. The delicate undeveloped vaginal mucous membrane in the child, on the other hand, is extremely susceptible to a gonococcal infection, which causes a peculiarly profuse purulent vaginitis and one that persists with much obstinacy. Such a vaginitis on healing has a great tendency to plastic adhesions of the vaginal walls. The most common situation for these adhesions to take place is at the junction of the anterior and middle third of the vagina where the two walls are normally in constant contact. In this way the vagina becomes completely occluded, and when menstruation is established the catamenial flow is obstructed and retained in the vaginal pouch behind the adhesion. As this accumulation increases each month, pressure is exerted on the cicatricial adhesion, and it is gradually stretched out into a well-marked membrane or diaphragm, which under sufficient pressure from behind may protrude from the vaginal orifice. Sometimes the anterior and posterior walls of the vagina may be completely glued together throughout the whole length of the vagina, a condition which is illustrated by Case 4 of this article. The plastic adhesion may take place at the orifice and cause atresia of the hymen, or the labia may be glued together and the vulva completely obliterated. Such conditions are due to the fact that, in all cases of purulent vaginitis in children, the vulva is involved, and is probably often the primary locus of infection, the epithelium of the vulva, like that of the vagina, being extremely susceptible to infection in children while very resistant in the adult. The present paper is confined to only those cases where the adhesion takes place in the vagina.

Acquired atresia of the vagina may also occur in adult women and may be the result of adhesions forming after a vaginitis following severe lacerations at childbirth. Adhesions from this cause usually occur much higher up in the vagina, and sometimes may include only the cervix, in which

case atresia of the cervical canal results. Senile vaginitis is not an uncommon cause of atresia of the vagina, but does not usually give rise to serious trouble because of the absence of menstruation. It sometimes happens, however, that secretions may collect behind an atresia even in women who have passed the menopause, and such a condition is illustrated by Case 3.

When the atresia has been acquired in early childhood, symptoms do not ordinarily manifest themselves until the age of puberty, though Kelly reports the case of a child where a large amount of mucus accumulated behind the atresia, causing a considerable tumor and indicating the necessity for an operation. At puberty the menstrual blood accumulates in the vagina behind the adhesive membrane, distends it and makes a tumor. If the condition is not relieved, the accumulation of blood added to by each menstrual period gradually distends first the vagina and then the uterus, and a very large tumor occupying a considerable portion of the abdominal cavity may be formed. In severe cases, as in Case 4, a double hematosalpinx may form. The pain is at first periodic, but may become continuous as the tumor enlarges.

The operation for atresia is one that is attended with considerable danger of fatal sepsis, and every precaution should be taken to prevent infection. After evacuating the old menstrual blood, which is always of a thick tarry consistency, the cavities of the vagina and uterus should be most carefully irrigated. Where there is a definite membrane occluding the vagina, the method of operation is of extreme importance. Cases 1 and 2 illustrate well the results which may ensue after an operation which consists only of an incision in the occluding membrane and the evacuation of the contents. The opening made by such an incision sometimes closes up entirely, but more often results in one or more pinhole apertures through which the menstrual blood may partially flow. A certain amount of mucus and blood is dammed back into the vagina, becomes foully infected and produces the dire results seen in Cases 1 and 2. In order to leave the vagina in a permanently normal condition the occluding membrane should be completely dissected out and the wound edges approximated with great care. Such a wound will heal up by first intention even in the face of a badly septic discharge. The details of the operation for dissecting out an occluding membrane are as follows:

OPERATION FOR ACQUIRED ATRESIA OF THE VAGINA.

After the septum has been incised and the contents of the space beyond have been removed, there should be a thorough cleansing of the parts. The edges of the septum should then be trimmed flush with the walls of the vagina. All scar tissue should be removed so as to prevent a future contraction. When the septum has been completely cleared away it will be found that there remains a denuded area which extends like a ring around the inner circumference of the vagina. The

* The drawings illustrating this article have been previously published in the *American Practice of Surgery*, vol. vi.

mucous membrane of the upper portion of the vagina should then be drawn down and approximated to that of the lower portion by interrupted catgut sutures. Such an operation should restore perfectly the normal caliber of the vagina and should not be followed by a tendency to the development of a stricture. The use of antiseptic douches twice or three times daily should be begun the day after the operation and should be kept up until the discharge completely disappears.

The following are four cases of acquired atresia which have occurred in my experience.

CASE 1. Mrs. A., age forty-five. Referred by Dr. Charles N. Cutler. Patient gave a history of a long-continued severe vaginitis at the age of ten, following an injury to the external genitals from falling astride a fence. At the time of puberty she had the usual symptoms of atresia, namely, great pain at the time for a monthly period, but without appearance of blood. She was operated upon, the operation apparently consisting of a puncture of the occluding membrane with an evacuation of the secretions. The patient was married at about the age of twenty-four. She menstruated regularly from the time of the operation, the menstrual flow being sometimes foul. When I saw the patient she was suffering from an extremely severe pruritus vulvæ. Examination showed a white leathery appearance of the external genitals which represented an early stage of the condition called kraurosis. Vaginal examination showed a well-developed membrane across the vagina situated at the junction of the outer and middle third. Through the membrane was a pinhole opening which doubtless represented the puncture which had been performed early in life. The patient had menstruated incompletely through this opening, and as the secretions, which were continuously partially dammed back by the occluding membrane, had become infected, a chronic pruritus had been caused by the irritating discharges. An operation was performed on the patient according to the method described above for acquired atresia, the contour of the vagina being entirely restored to its normal character. The condition of kraurosis, though relieved for a time, continued, however, to develop and necessitated a complete removal of the vulva, by which the patient was entirely cured.

CASE 2. Miss L., age twenty-two. Referred by Dr. Charles N. Cutler. Patient had a history of early vaginitis, monthly pain without blood at the age of puberty, and an operation of puncture of the vaginal membrane with evacuation of contents of the vagina. The patient menstruated incompletely and had frequent discharges of such a foul nature that she was extremely offensive to people about her. Examination showed a condition exactly like that of Case 1, namely, a membrane at the junction of the middle and outer third of the vagina, with a pinhole opening through which exuded a purulent discharge of a nauseating odor. An operation was performed by the method described above, a large amount of bloody purulent material being evacuated from the upper portion of the vagina, which had been distended. The vaginal mucous membrane showed a condition of the most severe inflammation. The wound of the operation healed by first intention and the patient after a period of local treatment for the inflamed vaginal mucous membrane made a complete recovery.

CASE 3. Mrs. M., age fifty-one. Referred by Dr. Charles N. Cutler. This patient had had several attacks of severe pelvic pain which were relieved after the sudden discharge of a large amount of muco-

purulent matter. Examination of the patient showed that she had had a plastic vaginitis of the upper part of the vagina so that the cervix was entirely occluded. Pelvic examination showed that the patient also had evidences of an old pelvic inflammatory disease. The cervical canal was opened without ether, and after an unsuccessful attempt to keep the canal patent, I decided to perform a hysterectomy. The pelvis was found full of old adhesions. The pelvic organs were removed and the patient made a good recovery.

CASE 4. Miss E., age fourteen. Previous to my seeing her this patient had had two operations for supposed congenital atresia of the vagina. She had the typical symptoms of atresia accompanying functioning pelvic organs. Examination showed occlusion of the vagina less than an inch from the introitus. Rectal examination showed a very large tender uterus with tubal masses on both sides of the pelvis. There were signs of a mild pelvic peritonitis, and the patient was running some temperature. At the operation the anterior and posterior walls of the vagina were found completely glued together and the surfaces were separated with considerable difficulty. When the vault of the vagina was reached, no signs of a cervix or of a cervical canal could be seen. The cervical canal was found after much trouble and the uterus was evacuated of a large amount of dark, semi-fluid material. The uterine cavity was thoroughly irrigated. An incision was then made in the median line of the abdomen and both tubes were found to be enormously distended with blood. There was some pelvic peritonitis. The tubes were removed and the pelvis was drained. This operation did not prove satisfactory as it was found impossible to keep the cervical canal patent. A glass-stem pessary was kept in the uterus several times for long periods. The pessary, however, kept up an inflammatory condition and the cervical canal each time closed up within a few days after its removal. The patient's condition finally became a serious one, and it was necessary to perform a hysterectomy, which was followed by a complete recovery. Although this case had been pronounced one of congenital atresia by several surgeons who had seen her, I have classified it as of the acquired type, as all the pelvic organs were perfectly normal. Although no history could be gained, this girl unquestionably had had a vaginitis in childhood which had caused a complete atresia of the whole length of the vagina.

CONGENITAL ATRESIA.

Many cases of congenital atresia do not require operation because they are frequently associated with defective development, or partial absence of the internal pelvic organs, so that menstruation has not been established. It sometimes happens, however, that secondary sexual characters are normally developed, and operation for the purpose of coition is demanded by the patient. In other cases the internal pelvic organs may functionate normally, and it is important to establish an artificial vagina for the purpose of drainage. The establishment of a permanent artificial vagina is a matter of extreme difficulty, and only a few successful cases have been reported. The operation described in this article is based on successful operations by Burrage and Abbey, in which flaps were turned in from neighboring areas. One writer has been successful in grafting in the vaginal tissue removed from a case of cystocele. The following is the technic of the operation which I

have devised for congenital atresia based on the operations of Burrage and Abbey, and which in the only case where I have had an opportunity to try it was successful.

The patient is placed in the perineal position. A superficial transverse incision is made just below the urethra. Through this transverse slit the bladder and rectum are slowly dissected apart with the finger. The forefinger of the left hand should be kept in the rectum, and some blunt instrument like a urethral dilator should be held in the bladder by an assistant. The dissection should be carried by the right forefinger, the left forefinger behind and the blunt instrument in front acting as guides to prevent tearing into the wall of the rectum or bladder. In this way an artificial opening can be made of the exact proportions of the normal vagina. Care should be exercised not to enter the abdominal cavity. The next step is to line this cavity with a pouch of skin made up of flaps turned in from the surrounding parts. To accomplish this, the two labia minora are dissected off, from above downward, in such a manner as to leave a pedicle sufficiently large to furnish good circulation. The two surfaces are then split apart so that two paddle-shaped flaps remain. There should then be dissected from the inner side of the thigh two similar flaps which have their bases at the two lower corners of the artificial opening. All four flaps are then sewed together over a glass form, as shown in the cut (Fig. 4). Before the flaps are sewed together several catgut sutures, with the ends left long, should be placed in the vault of the artificial cavity. When the skin pouch has been nearly completed, the glass form is removed and the catgut sutures are brought out through the skin pouch. The pouch is then inverted and the sutures are tied in such a manner that the pouch fits snugly into the artificial cavity.

The history of the case in which the preceding operation was successfully performed follows:

Miss M., age twenty-seven, was a young woman of apparently normal development and attractive appearance. She gave a history of never having menstruated. She had had normal sexual feelings and was engaged to be married. Examination showed normal development of breasts, hips and external genitals. Further examination revealed complete absence of the vagina, there not even being a dimple to indicate the introitus. By rectal examination no sign of uterus or ovaries could be felt.

The operation described above was performed, and the pouch formed by the turned-in flaps from the labia minora and from the thigh healed by first intention with the exception of a small area in the vault, and a permanent functioning vagina was established.

CANCER OF THE SCROTUM.

BY ROBERT M. GREEN, M.D., BOSTON.

CANCER of the scrotum, or chimneysweep's disease, was apparently first described as a pathologic entity in 1775 by Percivall Pott. In a pamphlet, or *Chirurgical Observation*, entitled "Cancer Scroti," he writes of it as follows:

"There is a disease peculiar to a certain set of people, which has not, at least to my knowledge, been publicly noticed; I mean the chimney-sweeper's cancer.

"It is a disease which always makes its first attack on, and its first appearance in, the inferior part of the scrotum, where it produces a superficial, painful, ragged, ill-looking sore, with hard and rising edges; the trade call it the soot-wart. I never saw it under the age of puberty, which is, I suppose, one reason why it is generally taken, both by patient and surgeon, for venereal, and, being treated with mercurials, is thereby soon and much exasperated; in no great length of time, it pervades the skin, dartos and membranes of the scrotum, and seizes the testicle, which it enlarges, hardens and renders truly and thoroughly dis-tempered; from whence it makes its way up the spermatic process into the abdomen, most frequently indurating and spoiling the inguinal glands. When arrived within the abdomen, it affects some of the viscera, and then very soon becomes painfully destructive.

"The fate of these people seems singularly hard; in their early infancy they are most frequently treated with great brutality, and almost starved with cold and hunger; they are thrust up narrow and sometimes hot chimnies, where they are bruised, burned and almost suffocated; and when they get to puberty, become peculiarly liable to a most noisome, painful and fatal disease.

"Of this last circumstance there is not the least doubt, though perhaps it may not have been sufficiently attended to, to make it generally known. Other people have cancers of the same parts; and so have others beside lead-workers the Poitou colic and the consequent paralysis; but it is nevertheless a disease to which they are peculiarly liable; and so are chimneysweepers to the cancer of the scrotum and testicles.

"If there be any chance of putting a stop to or preventing this mischief, it must be by the immediate removal of the part affected, — I mean that part of the scrotum where the sore is, — for if it be suffered to remain until the virus has seized the testicle, it is generally too late even for castration. I have many times made the experiment, but though the sores, after such operation, have, in some instances, healed kindly, and the patients have gone from the hospital seemingly well, yet in the space of a few months it has generally happened that they have returned either with the same disease in the other testicle, or in the glands of the groin, or with such wan complexions, such pale, leaden countenances, such a total loss of strength, and such frequent and acute internal pains, as have sufficiently proved a diseased state of some of the viscera, and which have soon been followed by a painful death.

"If extirpation ever bids fair for the cure of a cancer, it seems to be in this case; but then the operation should be immediate, and before the habit is tainted. The disease, in these people, seems to derive its origin from a lodgement of soot in the rugæ of the scrotum, and at first not to be a disease of the habit. In other cases of a