tively few) there is a partial active immunity which accomplishes an early abortion of the disease, or that the infecting organism was avirulent. We consider bronchopneumonia the one characteristic clinical and pathologic finding in our epidemic (82 per cent.) In other words, *the so-called complication is the disease.*

2. Bacteriologic examinations thus far made reveal an organism which seems best classified as an atypical pneumococcus and which we consider to be the probable cause of the epidemic.

Further clinical, bacteriologic and immunologic studies are under way.

808 Brandeis Theater Building.

PHLEGMONOUS GASTRITIS

REPORT OF CASE

F. I. SHATARA, M.D.

BROOKLYN

Phlegmonous or submucous gastritis is a rare condition. It was first described by Benel in 1656. In 1904, Robson and Moynihan collected eighty-five instances, and up to the present about 100 cases in all have been reported.

The condition is an acute suppurative inflammation affecting oftenest the submucous coat of the stomach, and extending to the muscular and serous coats more frequently than to the mucous. It is sometimes primary, but usually secondary, occurring in the course of such diseases as typhoid, scarlet fever, smallpox or puerperal septicemia. Two forms have been described, a diffuse and a circumscribed.

The etiology is not definitely known. Of the several theories that have been advanced, the bacterial seems to be the most plausible. Streptococci are the commonest micro-organisms present; colon bacilli have been occasionally demonstrated. In the case here reported, streptococci were demonstrated in the stomach wall by the Gram-Weigert stain. In secondary cases it is probable that the infection is carried by the blood stream. It is held by some that the invading organisms enter through traumatic or other lesions in the epithelium. Symmers, at Bellevue Hospital, however, failed to produce the lesion in animals by feeding them ground glass and inoculating streptococci and pneumococci by way of the blood stream or stomach tube, and the association of phlegmonous gastritis with such lesions as ulcer and cancer is almost unknown. The pyloric region of the stomach seems to be the site of predilection, although the process may involve the entire stomach.

The onset is sudden with symptoms of intense gastric irritability and profound general disturbances. The issue is rapidly fatal. Sometimes the symptoms of general peritonitis, as in the case here reported, are superimposed, and it is probable that many cases are similarly obscured and thus wrongly diagnosed.

Pain, epigastric tenderness, nausea, vomiting, chills, fever and hiccup are the outstanding symptoms. Hiccup was marked in the present case.

The course is usually from one to seven days.

The diagnosis during life is at best a guess. Chvostek is said to have succeeded in making one such diagnosis.

REPORT OF CASE

History.—C. W., man, aged 29, married, colored, laborer, American, applied for treatment at the dispensary, Dec. 19,

1917, complaining of a vague abdominal pain, general in character, intermittent in type, increasing usually after meals, but never severe enough to confine him to bed or to interfere with his appetite. The patient was constipated. There was no history of vomiting, gaseous or sour eructations, or jaundice. Physical examination revealed a slight abdominal tenderness, more marked on the right side, but no masses and no rigidity. Over the aortic area there was a faint systolic murmur transmitted to the carotids, but no other signs of cardiac trouble. The patient was advised to enter the ward for observation, but refused.

December 26, he returned to the dispensary, complaining of the same vague abdominal pain. He was again advised to enter the ward, and the withdrawal of blood for a Wassermann test was suggested, but he refused both.

Present illness.—On the morning of Jan. 19, 1917, the patient awoke with a headache, and was unable to take his breakfast; this he interpreted as a sign of severe illness. He applied to the dispensary at 9 a. m., and was given half a grain of codein and ordered to bed. At 10 a. m. he vomited and complained of pain in the epigastrium. At 4 p. m. his condition grew worse, the abdominal pain increased, nausea became distressing, and he vomited again. The vomitus was brownish in character, mixed with mucus, but free from blood or pus. Physical examination revealed general abdominal tenderness and rigidity, more pronounced on the right side. There were no signs of free fluid and no obliteration of liver dullness. The facial expression was anxious. The patient had a leukocyte count of 15,000, temperature of 101.6, pulse 100, and respirations 22. A ruptured gastric ulcer was suspected and he was admitted to the male surgical ward of the Englewood Hospital under the writer's care.

The patient slept about seven hours during the night. Next morning he repeatedly vomited green fluid. At 8 a. m. he had a leukocyte count of 27,400; polymorphonuclear neutrophils, 92 per cent.; lymphocytes, 8 per cent.; temperature, 103.2; pulse, 108, and respirations, 24.

He was operated on at 11:30 a. m., by Drs. Bradner and Pitkin. About 2 ounces of serosanguineous fluid were found in the peritoneal cavity. No definite perforation was located. A posterior gastro-enterostomy was impossible on account of excessive thickening of the stomach wall. A gastrostomy was performed, and it was noted that the stomach wall had lost its elasticity and had become so indurated that it was difficult to suture it to the gastrostomy tube. Two cigaret drains were inserted into the peritoneal cavity, and the patient was returned to bed. His general condition was fair, pulse 116, respiration 24. The rectal drip was employed and pituitary solution given.

About 11 p. m. the patient became very restless and thirsty, and complained of pain in the operative area. He also had a hiccup and there was a bloody discharge on his dressing. An irrigation with warm saline through the gastrostomy tube was performed; the return was at first bloody but later became clear. The temperature was up to 104.4, pulse 128, respirations 26. He slept about four hours subsequently.

Next morning, nutritive enemas were started.

In the drainage bottle to which the gastrostomy tube was connected there were 16 ounces of wine colored fluid.

About 8 a. m. the patient's temperature went down to 102, and his pulse to 118, but his respirations were accelerated to 32. At 3 p. m. the abdomen was distended and tympanitic, but there was no evidence of free fluid. He was given an alum enema with good result.

His general condition grew worse, his pulse rose to 138, and became thready, he was given stimulation but did not respond, and he died at 7:50 p. m.

Necropsy.—This was limited to the abdomen. There was about 1 quart of bloody fluid in the general peritoneal cavity. The fluid had apparently oozed through numerous minute perforations in the stomach.

The stomach wall was greatly thickened, measuring from a third to half an inch, and on compression, gastric contents oozed through several pin-point perforations. These perforations were clearly demonstrated by opening the stomach and holding it up against the light. The gastric mucosa looked and

felt leathery and smooth, the rugae having been obliterated. The process was diffuse.

The intestine was distended, and the intestinal wall was slightly congested. The mucous membrane was normal. The mesenteric nodes were enlarged.

The liver was pale and opaque; its substance was soft and mushy, reminding one of the liver in cases of acute yellow atrophy. The splenic tissue appeared normal except for a slight congestion.

The remaining organs were negative.

Numerous microscopic sections of the stomach were prepared in the pathologic laboratories at Bellevue Hospital. They were examined by Dr. Symmers, who found that the mucosa of the stomach was normal except for a superficial necrotic layer. The entire stomach wall beneath the muscularis mucosae was richly infiltrated by polymorphonuclear leukocytes, and was markedly edematous. The blood vessels of this portion of the stomach were deeply injected. Gram-Weigert preparations revealed the presence of innumerable streptococci lying chiefly in the stomach wall immediately under the muscularis mucosae. Microscopic sections from the intestinal tract revealed no pathologic changes.

27 Schermerhorn Street.

THE INCIDENCE OF TYPHOID CARRIERS IN A STATE INSANE ASYLUM *

CHARLES KRUMWIEDE, JR., M.D.

BRONXVILLE, N. Y.

AND

ELBERT M. SOMERS, M.D.

BROOKLYN

In September and October, 1907, the Brooklyn State Hospital had some cases of typhoid fever. At this time, Drs. Smith and Agnew of this institution sent to Miss W. Carey Noble, of the Bureau of Laboratories, the stools of all the convalescent patients; continued examinations showed that two women (E. O. and A. M.) had become chronic bacillus carriers. In 1912, there were a few cases of typhoid fever, and from these developed another chronic carrier (M. R.). Again, in 1915, a few inmates contracted the disease.

As previous reports have shown the tendency to a high carrier rate in asylums, we thought it worth while because of this history of institutional fever, to cooperate in a survey to determine the incidence of typhoid carriers. Under the direction of one of us, the institution was listed by wards, and ward residence since 1907 indicated for each inmate. Our intention was to include only the resident population at the time these lists were compiled. Specimens of feces were collected and brought to the laboratory as soon as possible after evacuation, and were there examined for the *Bacillus typhosus*. All the inmates in the female wards have been examined, yielding the following results:

Wards 1 and 2, population (February, 1916), 91: No cases of typhoid developed in this ward. One resident had typhoid in Ward 9 in 1907. No carriers were found.

Wards 3 and 4, population (March, 1916), 65: There were two cases of typhoid in this ward in 1907, and both the patients became carriers (E. O. and A. M.). Two additional carriers (M. C. and H. G.) were found, both without a history of typhoid. M. R., a chronic carrier, who developed typhoid in Ward 9 in 1912, was transferred to this ward later in the same year, and died here in September, 1915.

Ward 5, population (April, 1916), 80: One case of typhoid developed in 1907. Another patient had typhoid in Ward 9 in 1912. No carriers were found.

Ward 6, population (April, 1916), 132: No cases of typhoid developed in this ward. No carriers were found.

Ward 8, population (April, 1916), 72: No cases of typhoid developed in this ward. Three residents gave histories of typhoid, two before admission to the hospital, one in 1907 in Ward 9. Examinations were negative. Two carriers (J. M. and M. W.) were found, both without a history of typhoid. M. W. died in February, 1916. The changes in ward residence of these carriers and of other carriers, then unrecognized, may have led to contact infection and the development of the carrier state, without recognizable symptoms of typhoid infection.

Ward 9, population (May, 1916), 76: Eight cases of typhoid have developed in this ward: two in 1907, four in 1912, and two in 1915. (Three of these persons are now residents in other wards, and noted there.) One 1912 patient (M. R.) became a chronic carrier and was transferred to Ward 3 (which see). Examinations of the four residents were negative. One typhoid carrier was found (M. B.) who denies a history of typhoid fever.

It will be seen from this summary that, in a total population of 516 female patients, the survey yielded five chronic carriers, in addition to three previously known. Of the last, all continued to excrete bacilli after having had typhoid fever, two in 1907 and one in 1912. Of the other five, none gives a history of typhoid fever. Too much reliance cannot be placed on negative histories of typhoid fever, especially from patients of this type, as the symptoms of a mild infection may have been overlooked.

In spite of this, however, we believe that it is not an unreasonable assumption that some of these patients coming in contact with cases or carriers ingested the bacilli, became normal or contact carriers, and later developed into chronic carriers. That is, their natural immunity prevented the development of a generalized disease; but in spite of this resistance, the bacilli multiplying in the intestine, small numbers may have invaded the body proper and through the lymphatic and vascular systems may have reached the liver and gallbladder, where they localized and gave rise to chronic inflammation. After their localization in the biliary passages, a chronic carrier state would be likely to result.

The striking thing in this investigation is the large number of carriers, eight among 516 persons, or 15.5 per thousand. This rate is higher than usually encountered in institutions and much higher than among the general population. The unusually high incidence is shown by applying the rate to the female population of New York State over 25 years of age. This would give the extraordinary result of 45,000 carriers.

In view of these results as well as because of the frequency of institutional typhoid fever, we do not believe that any one would question the advisability of vaccination of all inmates of insane asylums. Unfortunately, however, this procedure is left to the discretion of the institutional authorities and is not carried out as generally as it should be.

A considerable amount of institutional typhoid fever due to the failure to vaccinate has come to our notice during the last few years. Because of this, we feel that vaccination should be mandatory not only in asylums but also in all custodial institutions, and a regulation to this effect should be added to the sanitary codes of all health departments.

*From the Bureau of Laboratories, Department of Health, New York City, and the Brooklyn State Hospital.