lesion presents undermined borders and deep maceration, and is typically characteristic of the disease in this location.

The back of the scalp is markedly involved (Fig. 4). The lesion takes the form of a segment of a circle, and is about 6 inches long. The advancing border consists of a ridge of scaly papules. There are no vesicles to be seen. The color is dull red somewhat masked by the hair. There is no alopecia.

The Organism.—Scrapings were made from the edge of most of the lesions, including the scalp, and fungi corresponding in type to the *Epidermophyton inguinale* were demonstrated in potassium hydroxid preparations.

Cultures were made from the edge of the scalp lesion on Sabouraud's medium and on glucose agar. Only *Staphylococcus aureus* grew in the Sabouraud tubes. On the glucose agar, *Staphylococcus aureus* appeared in a few days. About ten days later, small yellowish colonies of fungi made their appearance and slowly grew until in about three weeks they were about one-half inch in diameter (Figs. 5 and 6). The color was a dirty white with a slight tinge of yellow. A week or so later the colonies were covered with fluffy, white tufts. Subcultures were made on glucose agar. The organism

described shows a lesion on the back of the scalp apparently autoinoculated from the extensive lesions elsewhere on the body. This location is not a flexure (except perhaps in very stout persons, and this patient is slender), so that it is apparent that at times inoculation may occur in places where there is no excess of heat and moisture.

Wall Building.

TREATED AND UNTREATED OSTEOCHONDRITIS JUVENILIS OF THE HIP*

WALLACE BLANCHARD, M.D.

Fellow of the American College of Surgeons

CHICAGO

Perthes' disease, or osteochondritis juvenilis of the hip, results in far more extensive bone atrophy than has been supposed. The bone atrophy and rarefaction

Fig. 4.—The lesion on the scalp.

Fig. 5.—Culture of *Epidermophyton inguinale* from the scalp.

Fig. 6.—Culture of *Epidermophyton inguinale* from the scalp.

reproduced true to type, but the cultures were also contaminated with staphylococcus.

Smears made from the cultures at about the third week showed the same organism observed in the scales from the lesions.

Hairs from the scalp lesion were examined in potassium hydroxid, but no organisms could be found.

Treatment.—The lesions were at first treated with Whitfield's ointment² with but little improvement. The patient was then given 3 per cent. pyrogallie acid for the scalp, pubic, crural, perineal and anal regions, and 3 per cent. chrysarobin for the remaining lesions. By May 22, 1917, the lesions had all disappeared.

COMMENT

Sabouraud³ states that "it [the *Epidermophyton inguinale*] does not attack the hair; it requires the bottom of a flexure for its inoculation." The case

are so extensive that they have changed our previous conception of the pathology of the disease.

Up to the present time, the only joint and bone changes recognized in Perthes' disease have been a partial or total destruction of the head of the femur with a stumping of the neck, together with a thinning of the femoral shaft. Most authors have agreed with the unsustained suggestion, made by Freiberg¹ at the last session of this Association, that the disease is caused by tonsillitis or some other local infection. Experiments have been made on animals by Allison² to determine what influence a slight injury or displace-

* Read before the Section on Orthopedic Surgery at the Sixty-Eighth Annual Session of the American Medical Association, New York, June, 1917.

1. Freiberg, A. H.: The Evolution of Osteochondritis Deformans Coxae Juvenilis, THE JOURNAL A. M. A., Aug. 25, 1916, p. 658.

2. Allison, Nathaniel: Am. Jour. Orthop. Surg., October, 1915.

2. Seven per cent. salicylic acid and 14 per cent. benzoic acid in white petrolatum (Whitfield: Brit. Jour. Dermat., 1911, 23, 375).

3. Sabouraud: Brit. Jour. Dermat., 1911, 23, 385.

ment to the epiphysis might have on the hip, and his results were negative.

Since I desired to make a comparative study of diseased and normal hips, I had roentgenograms taken of cases of Perthes' disease, to include all of the pelvis as well as both the femoral shafts on single large plates. In these comprehensive pictures, I soon discovered decreased bone density and some atrophy of the pelvis on the diseased side in the acute cases.

The bone atrophy and lessened density are of about the same degree in the ilium, ischium and pubic bone as in the shaft of the femur. The atrophy of the ischium and pubic bone usually reduces the obturator foramen to one-half the size of the opposite foramen, as is well shown in Figure 4. On examining roentgenograms of both legs in one picture, it soon became apparent that all the bones of the leg suffered moderate atrophic changes. Decreased density is also seen in all the surrounding muscles and other soft parts.

The shrinking of the head of the femur leaves a vacant space in the acetabulum which shows principally adaptive changes.

Atrophy of bones and muscles is one of the first symptoms, and is as equally constant in the functionally used diseased leg as in the unused diseased leg. The atrophy becomes less as the distance increases from the hip.

Early and continued mechanical treatment that protects the head of the femur from weight-bearing, jar and concussion will usually preserve the head in its rounded shape; but observation of a number of treated and untreated cases shows that the best available mechanical treatment has little or no effect in either preventing or causing atrophy of the bones and muscles of the leg of the diseased side.

I have not been able to locate a tonsillar or other source of infection in a single case.

The moderate but extensive atrophy and decreased density of bone and adjacent tissues precludes the possibility of the disease being due to a circumscribed joint infection or an epiphyseal injury. The disease is undoubtedly trophic in character and is probably caused by an extensively disturbed blood circulation that impairs nutrition from a median pelvic line to the foot.

It is unfortunate that this rather recently discovered disease of the hip joint has been differentiated from tuberculous hip disease, with which it has been so long confused, if the treatment is to be neglected because it runs a milder course and with much less resulting deformity.

Patients who have had Perthes' disease which has been diagnosed as tuberculous hip disease, and who have promptly received the best mechanical and general treatment usually accorded to tuberculous hips, have invariably made the best possible recoveries.

The neglected and untreated cases of Perthes' disease after one or two years show the heads of the femurs to be nearly destroyed, with the necks stunted, and in coxa vara or coxa valga positions. Abduction is partly lost, and movements of the hips in other directions are slightly limited. The legs are from 1 to 3 cm. short, and the patients walk with a limp.

The friction and concussion of weight-bearing on softened bone cause the erosion of the head and the stunting of the neck of the femur. The disease is self-limited. The extensive bone atrophy, together with the obliteration of the head of the femur, runs its

course in about one year, and this is followed by a longer period of bone hardening and rebuilding. One proof of this limited course is that if the head of the femur is protected from the outset of the disease by adequate mechanical treatment so that it retains its rounded shape for two years, it again becomes hard, firm and self-protecting.

Roentgenograms show a decreased density of bone in the first year of the disease, and during the second and third years the deepened shadows indicate that the diseased bones have become harder and more dense than normal bones. This hardening resembles the eburnation of soft bones after rickets, in which they frequently become as hard and firm in a child of 5 years as in a man of 25.

Fig. 1 (Case 1).—Atrophy and greatly reduced density of the pelvic bones, femur and muscles of the diseased side in acute Perthes' disease in a boy, aged 5 years.

There is perhaps no destructive joint lesion that is quite as amenable to treatment as Perthes' disease. The promptness with which the head of the femur rebuilds under mechanical protection is quite surprising.

If an adequate protective mechanical treatment is commenced even late in the course of the disease, when the head of the femur has been destroyed, and the treatment is then continued for a year or two, the head of the femur will regenerate to nearly its full rounded shape, in the same manner that the head of the femur rebuilds for several years after the reduction of a congenital hip dislocation, with the result that the ultimate shortening will be lessened fully a half. The limp is relieved in proportion to the regeneration of the head of the femur.

This rebuilding is a further proof that Perthes' disease is not an infective epiphysitis, since after the head of the femur has been destroyed by an epiphysitis, it does not regenerate.

Fig. 2 (Case 2).—Reduction of the head of the femur to a small pyramid of bone in E. M., a girl, aged 8, who had had Perthes' disease for one year; the leg is 2 cm. short; the patient walks with a limp.

Roentgenograms of untreated cases of Perthes' disease, taken from seven to eight years after the disease has run its course leaving only remnants of a head or a stunted neck, show that unaided Nature has been partly successful in rebuilding the head to its normal shape.

In Perthes' disease, lengthening of the leg rarely happens, but I have one case to report.

A. H., a boy, aged 6 years, has had pain in his hip, and walked for one year with the left leg abducted. The roentgenogram (Fig. 5) shows the usual atrophic changes. The abduction leverage on the softened neck and shaft of the

Fig. 3 (Case 2).—Regenerated head of the femur in E. M. at the age of 15 years; the patient had mechanical treatment during the second year of the disease, and walks without a limp.

femur during the one year of acute disease has straightened the normal outcurve of the shaft and elevated the neck about 35 degrees, causing a 4 cm. lengthening of the left leg. The abduction protected the head of the femur, which shows only atrophic changes. The hardening stage is somewhat

advanced, and nothing but an osteotomy at the neck can restore the normal relations. The boy walks with a limp to the right.

In marked contrast to the depression seen in tuberculous, hip disease, patients with Perthes' disease usually retain the irrepressible spirits of healthy childhood.

In going over thirty years of hospital and private histories of tuberculous hip disease, I have been able to cull out over forty cases of undoubted Perthes' disease in which the patients were treated as having tuberculous hips and discharged cured with results too suspiciously good to have been obtained in tuberculous cases. In a number of these cases, the roentgenograms confirm the late diagnosis of Perthes' disease. It is a consolation to know that even if the diagnosis was incorrect the patient received the best possible treatment.

The condition is not likely to be confused with any other than tuberculous hip disease. The patient is usually from 4 to 10 years of age, walks with a slight limp, and complains of occasional pain. There is

Fig. 4 (Case 3).—Head of the femur, in which only two small buttons are left, in an untreated case of Perthes' disease in a girl, aged 8 years; the leg is 2.5 cm. short, and the patient walks with a limp.

always some atrophy of the thigh and less of the leg. Abduction is limited, but movements in all other directions are free. These symptoms closely resemble the symptoms of an insidiously advancing tuberculosis of the hip joint, but the roentgenogram will usually decide. If it is too early for pathologic conditions to show decisively in the roentgenogram, one should be taken a month or several months later, but the mechanical treatment should be begun immediately, since in either condition the treatment will be the same, and the treatment cannot be commenced too early. With the hip joint well protected, constant exercise in the open air and a liberal raw food diet seem to be of greatest service in the treatment of Perthes' disease.

Every orthopedic surgeon is familiar with the plaster-of-Paris spica and with a variety of hip splints that afford good protection to the head of the femur, and I present the accompanying pictures of patients with Perthes' disease, the one in a plaster-of-Paris spica and the other in my strap iron hip splint with high shoe and crutches, only to emphasize the necessity of very complete protection to the hip joint in order to get the best possible ultimate results.

CONCLUSIONS

1. Osteochondritis deformans juvenilis seems to be a nutritional disease of the bones and muscles of the hip and leg.

Fig. 5 (Case 4).—Lengthening of the leg in Perthes' disease, a rare resulting deformity, in a boy, aged 6.

2. The femoral head that continues in functional use during the acute stage of the disease becomes obliterated wholly or in part by the attrition of weight-bearing on softened bone.

3. The head of the femur that is mechanically protected throughout the acute stage of the disease suffers only the same atrophic changes as the adjacent bones.

4. In convalescence, the destroyed head redevelops rapidly under mechanical protection, and redevelops more slowly and less perfectly without protection.

5. The disease progresses with one year of bone and muscle atrophy and lessened density followed by several years of increasing density and redevelopment.

6. The bone hardening resembles the eburnation after rickets.

7. From the foregoing facts it may be inferred that this disease of growing bone is nutritional character, and is due to a blood circulation that is disturbed from a median pelvic line to the foot, and that the head and neck of the femur are only incidentally affected.

8. The question of etiology remains of chief interest.

15 East Washington Street.

ABSTRACT OF DISCUSSION

DR. JOHN JOSEPH NUTT, New York: Although the author of the paper said that the etiologic factors are of the most interest, it seems to me that it is of more practical interest

to decide on the best method of treatment. A great many advocate treating these patients not at all; some advocate treating them slightly; and some advocate treating them much more thoroughly. If we treat them too slightly and neglect them, I think we will have worse results than by making mistakes in diagnosis and treating them as tuberculous hips.

One point that I have never thought of before is that as the head becomes flattened and lessened, the line formed by the lower edge of the shadow of the neck does not quite fall in line with the line showing the upper border of the obturator foramen.

DR. PERCY W. ROBERTS, New York: I merely wish to ask Dr. Blanchard what, in his experience, have been the acute symptoms; or, perhaps, to put it differently, in what proportion of cases has he seen such symptoms as pain and spasm?

DR. FREDERICK J. GAENSEN, Milwaukee, Wis.: I want to emphasize what I consider a very important point made by Dr. Blanchard. The tendency has been to let these patients go on about as they pleased, feeling that ultimate recovery, with but slight deformity, would result. I think he has sounded an important note of warning in advising against this, and I endorse his recommendation of judicious supportive treatment.

I wonder whether Dr. Blanchard did not make a mistake in referring to Dr. Freiberg as having written a paper on the infectious origin of osteochondritis. I think it was Dr. Kidner.

I became much interested in this condition when Calvé, in 1912, at Berc-sur-mer, showed me some cases of what we now know as Perthes' disease, which he described as pseudo-coxalgia. When I visited Perthes at Tübingen in the same summer he showed me examples of the same group of cases. I told him at that time that it corresponded exactly to the type of cases that Calvé had shown me. He was much interested, and I let him have the reprint that Calvé had given me. At first he thought the cases were not the same; but later changed his opinion, and regarded them as identical. If we want to add another name to the already hyphenated appellation of the disease, I think the name of Calvé should find a place in it.

DR. SAMUEL C. BALDWIN, Salt Lake City: I did not hear Dr. Blanchard say anything about the symptoms, or whether in his cases he found much pain. Last year, at the clinical meeting of the American Orthopedic Association at Philadelphia, Dr. James K. Young showed roentgenograms of several cases in which there had been a great deal of destruction of the bone. I have had one very interesting case recently, which illustrated what Dr. Blanchard has said about these cases having been treated for tuberculous hip disease, years back, when they were really cases of Perthes' disease.

About fifteen years ago, I had a boy whom I treated for hip disease. Last summer his sister was brought to me with a condition that looked exactly like Perthes' disease of the

Fig. 6 (Case 5).—A girl, aged 5 years, with Perthes' disease, in a plaster-of-Paris spica with high shoe and crutches.

hip, but it was in the second metatarsal bone. When I saw the girl first she was about 16 years old. There was no swelling or tenderness, and she had had practically no pain in the joint. If she walks a great deal and gets very tired the joint will sometimes pain her just a little. The head of the second metatarsal bone looks exactly like the head of the femur in a case of Perthes' disease.

Fig. 7 (Case 6).—A boy, aged 5 years, with Perthes' disease, in a strap iron hip splint with high shoe and crutches.

About fifteen years ago, as I have said, I treated her brother for hip-joint disease and he got well a great deal more rapidly than these patients usually do. He is well now, with very little limp, and I have wondered whether this was a case of hip disease or what we now know as Perthes' disease. His mother told me that he had been at work for a number of years and had a very slight limp. I do not know whether this condition has been described in any other bones than the head of the femur or not, but in the case of the sister I was impressed by the

shortening of the toe, such as we get in the head of the femur in Perthes' disease, and by the lack of other symptoms.

DR. JAMES W. SEVER, Boston: I should like to make a plea for the calling of this condition Legg's disease rather than Perthes' disease or Legg-Perthes' disease.

In the routine treatment of the cases at the Children's Hospital, Boston, use is made of the abduction splint; or what we used to employ is the old Taylor splint, and what we use now is the abduction splint, just as we do in tuberculosis of the hip. The patients are guarded to prevent walking on the affected limb, which is kept abducted and on a certain amount of traction. A plaster-of-Paris spica is not used, and the patients go on crutches. They do not get much deformity in the neck of the femur. They do, however, get some flattening and overgrowth of the edges of the epiphysis; and they recover with some limp, but very little deformity.

DR. EMIL S. GEIST, Minneapolis: Regarding the condition of toe deformity, we saw yesterday, at the clinic of Dr. Hibbs, a roentgenogram of that case. The condition has recently been well described by Dr. Freiberg. It bears a relation to osteochondritis of the hip, in that these toe cases are usually the result of injury, and the question arises whether there is a similarity between these two conditions in that regard.

DR. WALLACE BLANCHARD, Chicago: Many cases of Perthes' disease run a very mild course and remain unrecognized for several years.

I think that there was no mistake in my quotation from Dr. Freiberg. I got the information from the Transactions of the Section on Orthopedic Surgery of the American Medical Association, 1916.

The obturator foramen frequently becomes one-half the normal size at the end of the first year of the disease. It usually becomes gradually smaller until it ultimately diminishes to one-third the size of the opposite foramen.

One gentleman asked whether these patients come to us early. They seldom come early, and many times not until the second or third year of the disease.

Dr. A. T. Legg announced a paper before the American Orthopedic Association at Pittsburgh, in June, but failed to appear. In the announcement, he said that he was going to give the histories and descriptions of seventy-five cases of regenerated and flattened heads of femurs, and that twenty-five of these had been regenerations of the head after the reduction of congenital dislocation of the hip. The others he could not account for; and it occurs to me that most of these, and perhaps all, are rebuilt heads after unrecognized Perthes' disease.

CEREBROSPINAL FLUID PRESSURE*

A. L. SKOOG, M.D.

KANSAS CITY, MO.

While there is included herewith the report of a case with data and pathologic material adding strong support to the theory that cerebrospinal fluid is a secretion from the choroid plexus cells, I trust that it may serve a purpose in another direction, namely, that it may encourage more frequent observations of spinal fluid pressure, and especially the utilization of a manometer for accurate registration of subarachnoid tension. We are accustomed to see present day case reports and hospital charts with a lumbar puncture record giving the cell count, and Wassermann, colloidal gold and globulin tests; but seldom do we see a recorded figure for the spinal fluid pressure. There are a number of pathologic states involving the central nervous system in which a spinal fluid pressure reading is of as much importance as (in some of more importance than) a cell count or Wassermann test.

Before launching into the more practical facts, there are several historical, anatomic and physiologic considerations to be reviewed.

Relative to pressure and the circulation of the cerebrospinal fluid, it is essential to know through what organ the fluid makes its appearance, and by what route the exit is made. The choroid plexus was known to Herophilus, Galen, Vesalius and others.¹ While this gland attracted medical men of the early

Fig. 1.—Rachiocentesis apparatus.

Christian era, its function was appreciated only quite recently. Even now, with a considerable amount of valuable experimental data at our command, there remains some hesitancy in accepting the conclusion that the gland exists essentially for the purpose of secreting the cerebrospinal fluid. The work of

* Read before the Section on Nervous and Mental Diseases at the Sixty-Eighth Annual Session of the American Medical Association, New York, June, 1917.

1. Schäffer: Ueber den Bau und die Function der Epithelzellen des Plexus Choroideus, Beitr. z. path. Anat. u. z. allg. Path. (Ziegler's), Festschrift für Prof. Julius Arnold, 1903, p. 101.