Lueders, C. W. LABORATORY DIAGNOSIS OF HYPERTHYROIDISM. [Arch. Int. Med., Vol. 24, No. 4.]

In Lueders' opinion the laboratory affords the most reliable means toward the recognition of early hyperthyroidism. The sugar tolerance test seems an important aid in the detection of borderline or early cases of hyperthyroidism. The epinephrin test did not prove in the study of the cardiac neuroses, diagnostic of hyperthyroidism. It seemed rather an index of the sensitization of the sympathetic nervous system. Its value as a diagnostic test is increased when blood and urinary sugar estimations are recorded with pulse rate and blood pressure. Intramuscular injections gave best results. Tests for nitrogen loss and acidosis seemed suggestive as aids in the diagnosis of toxic hyperthyroidism. Further studies are being made to discover if borderline cases or early hyperthyroidism reveal such changes. Creatinuria did seem present in the thyroid disorder group, and when taken in conjunction with the other tests, was of value in the diagnosis of early hyperthyroidism.

Richter, H. M. THYROIDECTOMY. [J. A. M. A., Oct. 25, 1919.]

The basic fault in the operation for toxic goiter, according to this observer is not so much in faulty technic as in leaving too much goitrous tissue and an amount of thyroid sufficient to produce the reactions observed in many cases. An excessive degree of activity is excited by the operation in the thyroid left behind. The removal of "two thirds" to "three fourths" of the thyroid will not do; we should leave only a few grams. Richter has carried out the radical operation in a series of cases, but the records are faulty in determining how many of the series were complete. But in a consecutive series of more than 100 cases, dating back over five years, there was only one death. A few preliminary ligations were made before he felt safe in doing the radical operation, and he thinks such are preferable in a very limited number of more critical cases. In several, he found that a larger amount than was intended was left behind, and the greater part of what was left was strangulated by sewing through and through with heavy catgut. While it thus left an amount to undergo necrosis, no bad results followed, and it reduced the operative time below what would have been required for further excision.

Biggs, M. O. WASSERMANN REACTION IN HYPERTHYROIDISM. [Missouri State Med. J., Vol. 16, No. 10. J. A. M. A.]

Biggs calls attention to the large percentage of hyperthyroid cases in State Hospital No. 1, Fulton, Mo., giving a positive Wassermann in which it was impossible, either from clinical history, physical examination or inquiry into the family records, to establish any syphilitic infection or taint. The patients were insane, with two exceptions, and their

cases had been diagnosed under the head of "thyroigenous psychosis." These individuals all became unmanageable to a greater or less degree at their respective homes, necessitating their confinement in an institution for the insane. The psychosis of each, in the main, was characterized by wild delusions, irritability which sometimes developed into acute excitement, slow speech and deliberate mentation, absence of suicidal or homicidal tendencies, lassitude and indifference to surroundings and apparent feeble-mindedness in some. There was in these cases a total lack of the symptoms which are manifested in a psychosis which results from syphilis infection, either acquired or inherited. The physical signs were such as one would find in the average case of this type. In most of the cases an exophthalmic state was found to exist with marked tachycardia and other symptoms which accompany conditions of this kind. In some of the cases the enlargement of the thyroid was more lateral and varied in size to a great extent. The changes in the skin, teeth, blood, and temperature of the body, are typical of this class of cases. The prevailing mental tone associated with the disease was fear and apprehension, frequently associated with hallucinations of hearing and vision; voices were heard saying disagreeable things and with these hallucinations occurred anxious and agitated states. A brief case history of twelve patients is given.

Kamamura, K. TRANSLATION OF THE THYROID GLAND WITH INTACT BLOOD SUPPLY. [Journal of Experimental Medicine, July, 1919.]

The author, using the Carrel technic of blood vessel suturing, made successful autoplasmic thyroid transplants in dogs, the thyroid gland remaining in good condition and functioning for several months after the transplantation, even when the circulation had been interrupted for one and one half hours. The author was not as successful in obtaining permanent homoplasmic transplantation of the glands. The circulation through the transplanted vessels is reported as being as good as normal.

Levi, L. THYROENDOCRINE FEBRICULA. [Presse médicale, May 12, 1919.]

Levi reports several cases of thyroid instability and testicular insufficiency in children and lays stress on the instability of body temperature associated with this combination. These children are sensitive to cold and subject to chills, but also exhibit in succession different types of fever, viz., temporary slight fever; later, more prolonged fever, and finally, actual febriculæ, rather persistent and leading to a suspicion of some hidden infection possibly tuberculosis. These febriculæ are due either to some slight inflammation causing fever by autoinfection—from adenoid and lymphatic tissues—the latter provoking a reaction of the heat centers, rendered hypersensitive by the coexisting hypothyroidia; or, to a paroxysmal growth tendency, the patient being subject to the