reflex symptoms are more objective than the sensory symptoms, and that the extramedullary tumors are almost all located in the dorsal region of the cord. The extent of the area of rigidity is instructive. He speaks of his late researches on the innervation of the abdominal reflexes according to the segment of the spinal cord and involved roots.

Vaglio, R. SPINA BIFIDA STATISTICS. [La Pediatria, Jan., 1920.]

An examination by Vaglio of 10,000 infants showed 23 cases of spina bifida, or 1 in 434; 13 were females, 10 males. Most were brought to the hospital a few days after birth, less frequently in the first few months, and only 2 cases were not seen until they were 1 and 2 years old respectively. There was one case of double spina bifida. The tumor in 12 was in the lumbo-sacral region, in 3 in the sacral, in 3 in the lumbar, in 2 in the dorso-lumbar, and in 2 others in the lower dorsal and cervical regions respectively. The size varied from that of a large walnut to a large orange. In only one was the tumor considerably larger, being about the size of a fetal head. The skin was almost intact in 8 cases, intensely reddened and vascular in 5, and ulcerated in 7; in 3 there was an escape of cerebro-spinal fluid. In only one was there a tuft of hair over the tumor. Family history showed syphilis in only 3 cases, in another 6 cases its presence was shown either by the hereditary history and clinical evidence or the Wassermann reaction. In only one was the existence of other cases of spina bifida in the ascendants or collaterals noted. In one case the mother was tuberculous. Regarding concomitant affections, 7 had hydrocephalus, 6 clubfoot, 4 paralysis of the lower limbs associated in 2 with clubfoot, 1 had Little's disease, 1 cleft palate, 1 had a multiple malformation of the anus and genitals, and 1 had multiple small angiomas of the scalp.

Kimpton, A. R. TÆNIA SOLIUM IN SPINAL CORD. [Surg. Gyn. Obst., Feb., 1920.]

There is no record of a pork tapeworm being removed successfully from the spinal cord during life, and Kimpton thinks his is the first case. Mercur found a similar tumor of the spinal cord, but at necropsy. In Kingston's case a diagnosis of spinal cord tumor at the fourth or fifth dorsal segment had been made and operation was advised. A laminectomy was done on the fourth, fifth and sixth dorsal vertebræ and the cord exposed. In the tension of the cord a clear difference was noted. The tension appeared to be greater, and the cord was paler in this region. A translucent tumor was found when the cord membranes were incised. The tumor was easily shelled from the cord. It appeared as a gelatinous mass. The pathologic diagnosis was cysticercus racemosus.

Pende, N. RHIZOMYELIA. [Policlinico, Dec., 1919, Med. Sect. 12.]

In a woman of 48 who had had several attacks of acute articular rheumatism in eighteen years Pende found nothing to indicate any spe-